BOOK OF ABSTRACTS

6th INTERNATIONAL CONFERENCE ON COMPUTATIONAL MATHEMATICS AND ENGINEERING SCIENCES

20-22 May
2022, Ordu - Turkey

THE SIXTH INTERNATIONAL CONFERENCE ON COMPUTATIONAL MATHEMATICS AND ENGINEERING SCIENCES (CMES-2022), ORDU/TURKEY, MAY 20-22 2022

The **Sixth International Conference on Computational Mathematics and Engineering Sciences (CMES-2022)** will be held in Ordu University from **20- to 22 May 2022 in Ordu, Turkey.** It provides an ideal academic platform for researchers and professionals to discuss recent developments in both theoretical, applied mathematics and engineering sciences. This event also aims to initiate interactions among researchers in the field of computational mathematics and their applications in science and engineering, to present recent developments in these areas, and to share the computational experiences of our invited speakers and participants.

The Organizing Committee

©*All Rights Reserved. This conference is organized by a cooperation of several international organizations including Firat University, Final International University, Akdeniz University, Moulay Ismail University, Private University of Fes, Harran University, Atatürk University, Van Yüzüncü Yıl University. No part of this book can be reproduced or utilized in any forms or by any means, electronic or mechanical, including photocopying, recording, or by any information storage and retrieval systems, without permission from the authors.*

© The informations provided in the papers published in this book are under the responsibility of their author(s).

MESSAGE FROM THE GENERAL CHAIRS





Dear Conference Attendees,

We are honored to welcome you to the **Sixth International Conference on Computational Mathematics and Engineering Sciences (CMES-2022)** at Ordu University from 20 to 22 May 2022 in Ordu City, Turkey.

CMES, founded in 2016 at Faculty of Science and Techniques Errachidia Moulay Ismail University Morocco is an annual intarnational conference, which was very successful in the past years by providing opportunities to the participants in sharing their knowledge and informations and promoting excellent networking among different international universities. This year, the conference includes 200 extended abstracts, several submissions were received in response to the call for papers, selected by the Program Committee. The program features keynote talks by distinguished speakers such as : **Alissandra Maria Ragusa** from Catania University Italy, **Dumitru Baleanu**, from Institute of Space Sciences, Magurele-Bucharest, Romania, **Juan-Luis García Guirao** from Technical University of Cartagena, Spain, **Oscar Castillo** from Tijuana Institute of Technology, Tijuana, Mexico, **Hossein Jafari** from UNISA South Africa, **Vatan Karakaya** from Ahi Evran University, Turkey, **Ekrem Savas** from Usak University, **Muhammad Reza Safaei** from Florida University USA, **Erhan Coşkun** from Karadeniz Technical University, Turkey and **Hüseyin Merdan** from Tobb University, Turkey. The conference also comprises contributed sessions, posters sessions and various research highlights.

We would like to thank the Program Committee members and external reviewers for volunteering their time to review and discuss submitted abstracts. We would like to extend special thanks to the Honorary, Scientific and Organizing Committees for their efforts in making CMES-2022 a successful event. We would like to thank all the authors for presenting their research studies during our conference. We hope that you will find CMES-2022 interesting and intellectually stimulating, and that you will enjoy meeting and interacting with researchers around the world.

.

Hasan Bulut,

Firat University Elazig, Turkey.

Zakia Hammouch,

École Normale supérieure, Moulay Ismail University of Meknès 50000, Morocco

Thu Dau Mot University, Binh Duong Province, Vietnam

China Medical University Hospital Taichung 40402, Taiwan.

TOPICS

Applied Mathematics,
Financial Mathematics,
Control Theory,
Game Theory
Modeling of Bio-systems for Optimization and Control,
Linear and Nonlinear programming and Dynamics,
Artificial Intelligence,
Geometry and Its Applications,
Analysis and Its Applications,
Statistics and Its Applications,
Mathematics Education and Its Applications,
Algebra and Its Applications.
Engineering Sciences
Computer Science
Information technology
Electrical and Electronic Engineering
Ordinary, Partial, Stochastic and Delay Differential Equations
Chaos and Dynamical Systems
Numerical methods and scientific programming
Fractional Calculus and Applications,
Cryptography and its applications
Computational Fluids mechanics, Heat and Mass Transfers.
Economics and Econometric Studies
Topology and Its Application
Education Sciences
Economics and Econometric Studies
Topology and Its Application

COMMITTEE CHAIRS

Prof.Dr.Hasan Bulut, Firat University Elazig, Turkey

Prof.Dr.Zakia Hammouch, Ecole Normale Supérieure of Meknes, Moulay Ismail University Morocco

COMMITTEE CO-CHAIRS

Prof.Dr.Carlo Cattani, Tuscia University, Viterbo, Italy
Assoc.Prof.Dr.Haci Mehmet Baskonus, Harran University, Sanliurfa,Turkey
Prof. Dr. Ercan Çelik, Kyrgyz-Turkish Manas University, Bishkek, Kyrgyzstan
Prof. Dr. Fevzi Erdoğan, Yüzüncüyıl University, Turkey
Doç. Dr. Tolga AKTÜRK, Ordu University, Turkey

HONORARY COMMITTEE

Prof.Dr. Ali AKDOĞAN (Rector of Ordu University, Ordu, Turkey)
Prof.Dr. Fahrettin Göktaş (Rector of Fırat University, Elazığ, Turkey)
Prof.Dr. Hamdullah Şevli (Rector of Van Yüzüncü Yıl University, Van, Turkey)
Prof.Dr. Alpaslan CEYLAN (Rector of Kyrgyz-Turkish Manas University, Bishkek, Kyrgyz Republic)
Prof.Dr. İsmail Yüksek (Rector of Antalya Bilim University, Antalya, Turkey)
Prof.Dr. Hüseyin Yaratan (Rector of Final International University, Girne, Cyprus)
Prof.Dr. Ömer Çomaklı (Rector of Atatürk University, Erzurum, Turkey)
Prof.Dr. Ekrem Savaş (Rector of Uşak University, Uşak, Turkey)
Prof.Dr. Mehmet Sabri Celik (Rector of Harran University, Sanliurfa, Turkey)
Prof.Dr. Vatan Karakaya (Rector of Ahi Evran University, Kırşehir, Turkey)
Prof.Dr. Hassane Sahbi (President of Moulay Ismail University, Morocco)
Mohammed Aziz Lahlou (President of Universite Privee of Fes, Morocco)
Prof. Dr. Mohammed Ouazzani Jamil (Vice President of Universite Privee of fez, Morocco)
Prof.Dr. Omar Ossouaddi (Vice President of Moulay Ismail University, Morocco)
Prof.Dr. Mohammed Amine (Director of ENS Meknes, Moulay Ismail University, Morocco)
Prof.Dr. Selahattin Maden (Dean of Science and Literature Faculty, Ordu University, Ordu Turkey)
Prof. Dr. Sadık Kılıç (Dean of Education Faculty, Ordu University, Ordu Turkey)
Prof.Dr. Mikail Et (Dean of Science Faculty, Fırat University, Elazig, Turkey)
Prof.Dr. Etibar Penahlı (Baku State University, Baku, Azerbaijan)
Prof.Dr. Necdet Bildik (Retired Faculty Member)
Prof.Dr. Mahmut Ergüt (Namık Kemal University, Tekirdağ, Turkey)
Prof.Dr. Rifat Colak (Fırat University, Elazig, Turkey)
Prof.Dr. Vedat Asil (Firat University,Turkey)
İbrahim Taşel (Trustees Board Member, Final International University, Gime, Cyprus)
Şevket Ertem (Trustees Board Member, Final International University, Girne, Cyprus)

SCIENTIFIC COMMITTEE

Abdon Atangana (University of the Free State, South Africa; China Medical University, Taichung, Taiwan)
S.Sivasundaram (Université de Bethune-Cookman, USA)
Carlo Cattani (Tuscia University, Viterbo Italy)
Yury Luchko (Beuth Hochschule für Technik Berlin, Germany)
Juan Luis Garcia Guirao (Cartagena University, Spain)
Ouazzani Jamil Mohamed (Private University of fez, Morocco)
Fikret Aliyev (Bakû State University, Bakû, Azerbaijan)
Hüseyin Demir (Ondokuz Mayıs University, Turkey)
Ali Rostami (University of Tabriz, Tabriz, Iran)
Daniele Ritelli (University of Bologna, Bologna, Italy)
Sakthivel Rathinasamy (Bharathiar University, India)
İbrahim Alinur Büyükaksoy (Okan University, Turkey)
Mehmet Emir Köksal (Ondokuz Mayıs University, Turkey)
Mehmet Karay (Final International University, Girne, Cyprus)
A.Bashirov (Eastern Mediterranean University, Cyprus)
Vali Bashiry (Cyprus International University, Cyprus)
Sunil Purohit (Maharana Pratap University, India)
Kolade Owolabi (Federal University of Technology, Negeria)
Praveen Agarwal (ANAND International College of Engineering Jaipur, India)

Anna Sandulyak (Université de Moscow, Russia)
Robert Tchitnga (Dshang University, Cameroon)
Sare Şengül (Marmara University, Turkey)
Shilpi Jain (Poornima College of Engineering, India)
Francesco Villecco (University Salerno ,Italy)
Abdulhamit Subaşı (University of Turku, Finland)
İsmail Özgür Zembat (University of Glasgow, England)
Mohammad Mursaleen (China Medical University, Taiwan)
A.Yousef (Kuwait Col.Science and Technology. Kuwait)
Ram Jiwari (Indian Institute of Technology, Roorkee, India)
L. Stošić (College for Preschool Teachers, Aleksinac, Serbia)
Alla Belousova (Don State Technical University, Russia)
A. Barakoska (University St. Macedonia)
Vera Stojanovska (University Ss. Cyril Macedonia)
Irina Abakumova (Don State Technical University, Russian)
Sonja Velickovic (College for Preschool Teachers, Serbia)
Bashar A.Al-Talip (University of Mosul,Iraq)
M. Addam (Nat.School of Applied Sc.Hoceima, Morocco)
Kexiang XU (Nanjing University, China)
Cameron Browne (University of Louisiana at Lafayette ,USA)
Christos Volos (Aristotle University, Greece)
Fahd Jarad (Cankaya University Turkey)
Thabet Abdeljawad (Prince Sultan University KSA.)
Ahmed Elngar (Beni Suef University Egypt.)
L. M.Sánchez Ruiz (Technical University of Valencia, Spain)
Armando Ciancio (University of Messina, Italy)
R.Chapko(Ivan Franko National University Lviv , Ukraine)
Gonca Onargan (Final International University, Girne, Cyprus)
Behçet Öznacar (Near East University, Cyprus)
R. Aggarwal (Deshbandhu College, University of Delhi, India)
A. Rababah (Univ.Science and Technology Irbid, Jordan)
Fahrettin Muhtarov (Bakû State University Bakû, Azerbaijan)
S. Rashid (Gov. College University Faisalabad, Pakistan)
Dyyak Ivan (Ivan Franko Natıonal University,Lviv,Ukraine)
I. Prokodushyn (Ivan Franko Natıonal University,Lviv,Ukraine)
Yusif Gasimov (Azerbaijan University, Bakû, Azerbaijan)
El Mehdi El Khattabi (ENS Meknes, Moulay Ismail University, Morocco)
A. Elazzouzi (Mohamed Ben Abdellah University, Morocco)
Chérif Ziti (Moulay Ismail University, Morocco)
Mohamed Ouzahra (Mohamed Ben Abdellah University, Morocco)
Mustapha Ait Khellou (ENS Meknes, Moulay Ismail University, Morocco)
Youssef El Haoui (ENS Meknes, Moulay Ismail University, Morocco)
Alireza Khalili Golmankhaneh (Department of Physics, Faculty of Science, IAU Urmia Branch, Iran)
Zulqurnain Sabir (Hazara University, Pakistan)

Carla M.A. Pinto (ISEP | Instituto Superior de Engenharia do Porto, Porto – PORTUGAL)
Hossein Jafari (UNISA University Pretoria, South Africa)
Hasan Bulut (Firat University, Turkey)
Haci Mehmet Baskonus (Harran University, Turkey)
Bayram Şahin (Ege University,İzmir ,Turkey)
Dumitru Baleanu (Cankaya University, Turkey)
Eva Kaslik (University of Timisoara, Romania)
Izhar Uddin (Jamia Millia İslamia University. New Delhi)
Ayhan Şerbetçi (Ankara University, Ankara)
Hacer Sengul (Harran University, Sanlıurfa,Turkey)
Ayşe Dilek Maden (Selçuk University, Konya)
Wei Gao (Yunnan Normal University, China.)
Rauf Amirov (Cumhuriyet University, Sivas, Turkey)
Ogün Doğru (Gazi University, Ankara, Turkey)
Hanlar Reşitoğlu (Iğdır University, Turkey)
Erhan Güler (Bartın University, Turkey)
Fethi Bin Muhammad Belgacem (PAAET, Shaamyia, Kuwait)
Omer Kisi (Bartın University, Turkey)
Zhonglong Zhao (North University of China, China)
Bilal Alatas (Firat University, Turkey)
Necati Ozdemir (Balikesir University, Turkey)
Derya Avci (Balikesir University, Turkey)
Firat Evirgen (Balikesir University, Turkey)
Erol YAŞAR (Mersin University, Mersin, Turkey)
Ramadhan A. Mohammed (University of Duhok-Iraq)
Sizar A. Mohammed (University of Duhok-Iraq)
Zanyar Anwer Ameen (University of Duhok-Iraq)
Mehmet Yavuz (Necmettin Erbakan University, Turkey)
Hüseyin Yıldırım (Maraş Sütçü İmam University, K.Maraş)
Filiz Taşcan (Eskişehir Osmangazi University, Eskişehir)
Dursun Irk (Eskişehir Osmangazi University, Eskişehir)
Ali Karcı (İnönü University, Malatya)
Davut Hanbay (İnönü University, Malatya)
Burçin DOĞAN (Malatya Turgut Özal University, Turkey)
Feyza Esra ERDOĞAN (Ege University, İzmir, Turkey)
Fulya ŞAHİN (Ege University, İzmir, Turkey)
Ahmet Bedri ÖZER (Firat University, Elazığ, Turkey)
Ülkü YILMAZ (Malatya Turgut Özal University, Turkey)
Erdal EKİCİ (Çanakkale Onsekiz Mart University, Çanakkale)
Mehmet Akif AKYOL (Bingöl University, Bingöl, Turkey)
Hamza EROL (Mersin University, Mersin, Turkey)
H.Bayram Karadağ (İnönü University, Malatya)
M. Kemal Özdemir (İnönü University, Malatya)
Recep Şahin (Kırıkkale University, Kırıkkale)

M. Baki Yağbasan (Kırşehir Ahi Evran University, Kırşehir)
Mustafa Mikail Özçiloğlu (Kilis 7 Aralık University, Turkey)
Çiğdem İnan (Mersin University, Turkey)
Mustafa Yıldırım (Cumhuriyet University, Turkey)
Mehmet Küçükaslan (Mersin University, Turkey)
Okan Şakar (Bahçeşehir University, Turkey)
Hristos Anastassiu (Int. Hellenic University, Serres, Greece)
Aly Ramadan Seadawy (Taibah University, Saudi Arabia)
İpek A. Turgut (İskenderun Technical University, Turkey)
Mustafa Açıkkar (Adana Science and Tec.University, Turkey)
Bülent Tavlı (TOBB Economics and Tec.University, Turkey)
Halis Yılmaz (Dicle University,Turkey)
Yasin Kaya (Dicle University, Turkey)
Sunil Kumar (National Institute Of Tech. Jamshedpur, India)
Hatun Özlem Güney (Dicle University, Turkey)
Sedat İlhan (Dicle University, Turkey)
Arife Atay (Dicle University, Turkey)
Seçil Yalaz (Dicle University, Turkey)
Özge Kuran (Dicle University, Turkey)
Damla Barlak (Dicle University, Turkey)
Mahmut Isık (Harran University, Sanlıurfa,Turkey)
Tamer Uğur (Ataturk University /Turkey)
Muzaffer Aşkın (Munzur University,Tunceli,Turkey)
Elman Hazar (Igdır University, Turkey)
Serap Şahinkaya (Tarsus University)
Naci Genc (Van Yüzüncü Yıl University, Turkey)
Remzi Tuntas (Van Yüzüncü Yıl University, Turkey)
Kadriye Aydemir (Amasya University,Turkey)
Yadigar Sekerci (Amasya University,Turkey)
Hayati Olgar (Gazi Osman Paşa University,Turkey)
Semih Çavusoglu (Gazi Osman Paşa University,Turkey)
Nagehan Kılınc Gecer (Kırsehir Ahi Evran University,Turkey)
Cem Oguz (Kırsehir Ahi Evran University,Turkey)
Arzu Denk Oguz (Atılım University,Turkey)
Erhan Pişkin (Dicle University,Diyarbakır,Turkey)
Aynur Yalcıner (Selcuk University,Konya,Turkey)
Javid Ali (Aligarh Müslim University,India)
Hamit Mirtagiloğlu (Bitlis Eren University,Bitlis,Turkey)
Heybetkulu Mustafayev (Yüzüncüyıl University, Turkey)
Cesim Temel (Yüzüncüyıl University, Turkey)
Mehmet Mendeş (Canakkale 18 Mart University, Turkey)
Çağrı KARAMAN (Atatürk University – Erzurum, Turkey)
Ceren KINA (Malatya Turgut Özal University, Turkey)
Harun KAYA (Malatya Turgut Özal University, Turkey)

Olha Ivanyshyn (Izmir Instıtute of Technology,Izmir, Turkey)
Ozlem Ertekin (Munzur University, Tunceli, Turkey)
Onur Alp İlhan (Erciyes University, Turkey)
Danyal Soybaş (Erciyes University, Turkey)
Fatma Berna Benli (Erciyes University, Turkey)
Inan Ünal (Munzur University, Turkey)
Kemal Aydın (Selçuk University ,Turkey)
Hakan Şimşek (Kırıkkale University /Turkey)
Bilal Şenol (İnönü University, Malatya)
Sertaç GÖKTAŞ (Mersin University, Mersin, Turkey)
Derya Arslan (Bitlis Eren University, Bitlis, Turkey)
Ali Akgül (Siirt University ,Turkey)
Esra Karataş Akgül (Siirt University ,Turkey)
Abdulkadir Karakaş (Siirt Üniversity,Turkey)
Murat Sat (Erzincan Binali Yıldırım University, Turkey)
İsmail Onur Kıymaz (Ahi Evran University , Turkey)
Ayşegül Çetinkaya (Ahi Evran University , Turkey)
Nergiz (ÖNEN) POYRAZ (Çukurova University, Turkey)
Ozen Ozer (Kırklareli University, Turkey)
Sinan Çalık (Firat Univeristy,Elazıg,Turkey)
Murat Karakaş (Bitlis Eren Univeristy,Bitlis,Turkey)
Ayşe Metin Karakaş (Bitlis Eren Univeristy,Bitlis,Turkey)
Ahmet Yıldız (İnönü Univeristy,Malatya,Turkey)
Aynur Keskin KAYMAKÇI (Selcuk University, Konya)
Nejmi Cengiz (Ataturk University, Turkey)
Abdullah Kopuzlu (Ataturk University, Turkey)
Kürşat Akbulut (Ataturk University, Turkey)
Mesut Karabacak (Ataturk University, Turkey)
Yeşim Saraç (Ataturk University, Turkey)
Sıdıka Şule Şener (Ataturk University, Turkey)
Sait Taş (Ataturk University, Turkey)
Fatma Taşdelen Yeşildal (Ankara University, Turkey)
Şevket Gür (Sakarya Univercity, Turkey)
Serdal Pamuk (Kocaeli Univercity, Turkey)
Fethiye Müge Sakar (Dicle University, Turkey)
Ceren Sultan Elmalı (Erzurum Technical University, Turkey)
Muhammed Yiğider (Erzurum Technical University, Turkey)
Sibel Turanlı (Erzurum Technical University, Turkey)
Hatıra Günerhan (Kafkas University, Turkey)
Murat Çağlar (Erzurum Technical University, Turkey)
Erhan Deniz (Kafkas University, Turkey)
Nizami Mustafa (Kafkas University, Turkey)
Semra Yurttançıkmaz (Ataturk University, Turkey)
Nigar Yıldırım Aksoy (Kafkas University, Turkey)

M. Ali Dokuyucu (Agrı İbrahim Cecen University, Turkey)
Ahmet Ocak Akdemir (Agrı İbrahim Cecen University, Turkey)
Talat Körpinar(Muş Alparslan University, Turkey)
Selçuk Baş (Muş Alparslan University, Turkey)
Rıdvan Cem Demirkol (Muş Alparslan University, Turkey)
Zeliha Körpinar (Muş Alparslan University, Turkey)
Evrim Toklu (Agrı İbrahim Cecen University, Turkey)
Serhan Eker (Agrı İbrahim Cecen University, Turkey)
Kadirhan Polat (Agrı İbrahim Cecen University, Turkey)
Abdullah Çağman (Agrı İbrahim Cecen University, Turkey)
Şebnem Yıldız (Ahi Evran University, Kırşehir, Turkey)
Erhan Set (Ordu University, Turkey)
Esin Ilhan (Kırsehir Ahi Evran University / Turkey)
Hasan Toğrul (Batman University, Batman, Turkey)
İnci Toğrul (Batman University, Batman, Turkey)
Selahaddin Maden (Ordu University, Turkey)
Yıldıray Çelik (Ordu University, Turkey)
Mehmet Korkmaz (Ordu University, Turkey)
Süleyman Şenyurt (Ordu University, Turkey)
Erdal Ünlüyol (Ordu University, Turkey)
Yusuf Pandır (Bozok University – Turkey)
Asıf Yokuş (Fırat University, Elazığ, Turkey)
Hülya Durur(Ardahan University, Ardahan, Turkey)
Cemil İnan (Mardin Artuklu University, Turkey)
Çetin Kürşat Bilir (Ahi Evran University, Turkey)
Alper Ekinci (Bandırma 17 Eylul University, Turkey)
Fatma Sağsöz (Ataturk University, Turkey)
Hayriye Esra Akyüz (Bitlis Eren University,Bitlis,Turkey)
Fahrettin Özbey (Bitlis Eren University,Bitlis,Turkey)
Hatice Kuşak Samancı (Bitlis Eren University,Bitlis,Turkey)
Akbar Barati Chiyaneh (Yüzüncüyıl University,Bitlis,Van)
Ufuk Kaya (Bitlis Eren University,Bitlis,Turkey)
M. Tuncay Gençoğlu (Fırat University, Elazığ, Turkey)
Berivan Polat (Munzur University, Turkey)
Alper Polat (Munzur University, Turkey)
Şükran Konca (İzmir Bakırçay University, İzmir, Turkey)
Ali Çakmak (Bitlis Eren University, Bitlis,Turkey)
Unal İç (Firat Üniversity, Elazıg)
Seda İğreti Araz (Siirt University, Siirt, Turkey)
Burcu Gürbüz (Johannes Gutenberg-University Mainz, Germany)
Mahir Murat Cengiz (Ataturk University /Turkey)
Nguyen Huy Tuan (Thu Dau Mot university, Vietnam)
Seenith Sivasundaram (Université de Bethune-Cookman, USA)
Agamirza Bashirov (Eastern Mediterranean University, Cyprus)

Lazar Stošić (College for Preschool Teachers, Aleksinac, Serbia)
Alla Belousova (Don State Technical University, Russian Federation)
Aneta Barakoska (Faculty of Philosophy, University St. Cyril Macedonia)
Vera Stojanovska (Faculty of Philosophy, University Ss. Cyril Macedonia)
Irina Abakumova (Don State Technical University, Russian Federation)
Sonja Velickovic (College for Preschool Teachers, Aleksinac, Serbia)
Mohamed Addam (National School of Applied Sciences Hoceima, Morocco)
Luis Manuel Sánchez Ruiz (Technical University of Valencia, Spain)
Roman Chapko(Ivan Franko National University Lviv , Ukraine)
Yarema Savula (Ivan Franko National University Lviv , Ukraine)
Ihor Prokodushyn (Ivan Franko Natıonal University,Lviv,Ukraine)
Zakia Hammouch (Moulay Ismail University, Morocco)
Burçin DOĞAN (Malatya Turgut Özal University, Malatya, Turkey)
Ahmet Bedri ÖZER (Malatya Turgut Özal University, Malatya, Turkey)
Ülkü YILMAZ (Malatya Turgut Özal University, Malatya, Turkey)
Mehmet Atçeken (Tokat Gaziosmanpaşa University, Tokat)
Hristos Anastassiu (International Hellenic University, Serres, Greece)
İpek Abasıkeleş Turgut (İskenderun Technical University, Turkey)
Mustafa Açıkkar (Adana Science and Technology University, Turkey)
Bülent Tavlı (TOBB Economics and Technology University, Turkey)
Salman Khodayifar (Institue for Adv.Studies in Basic Sciences,Iran)
Mehmet Mendeş (Canakkale 18 Mart University Canakkale, Turkey)
Ceren KINA (Malatya Turgut Özal University, Malatya, Turkey)
Harun KAYA (Malatya Turgut Özal University, Malatya, Turkey)
Nergiz (ÖNEN) POYRAZ (Çukurova University, Adana, Turkey)
Mustafa Ali Dokuyucu (Agrı İbrahim Cecen University, Turkey)
Fatma Erdoğan (Firat University, Elazığ)
Gökhan Gökdere (Fırat University, Elazığ)
Yusuf PANDIR (Yozgat Bozok University, Yozgat Turkey)
Yusuf GUREFE (Uşak University, Uşak Turkey)
Şükran Konca (Bitlis Eren University, Bitlis,Turkey)
Hüseyin Merdan(TOBB University of Economics and Technology, Turkey)
Songül Kaya Merdan(Middle East Technical University,Turkey)
Engin Özkan(Erzincan Binali Yıldırım University, Erzincan)
İsrafil Okumuş(Erzincan Binali Yıldırım University, Erzincan)
Bayram Çekim(Gazi University, Turkey)
Gürhan İçöz(Gazi University, Turkey)
Fatma Ayaz(Gazi University, Turkey)
Emel Biçer (Bingöl University, Turkey)
Güven Kaya (Bingöl University, Turkey)
Ramin Najafi (Islamic Azad University, Iran)
Ebru Bozkurt (Ataturk University, Turkey)
Rıdvan Şahin(Gumushane University, Turkey)
İsa Yıldırım (Ataturk University, Turkey)

Halit Orhan (Ataturk University, Turkey)
Arzu Akbulut (Eskisehir Osmangazi University, Turkey)
Erdal Ulualan (Kutahya University, Turkey)
İbrahim Aktaş (Gumushane University, Turkey)
Numan Yalçın (Gumushane University, Turkey)
Muhammet Kamali (Kyrgyz-Turkish Manas University, Kyrgyzstan)
İsmet Altıntaş (Kyrgyz-Turkish Manas University, Kyrgyzstan)
Anargül Urdaletova (Kyrgyz-Turkish Manas University, Kyrgyzstan)
Asan Ömüraliyev (Kyrgyz-Turkish Manas University, Kyrgyzstan)
Avıt Asanov (Kyrgyz-Turkish Manas University, Kyrgyzstan)
Kalışkan Matanova (Kyrgyz-Turkish Manas University, Kyrgyzstan)
Ella Abilayeva (Kyrgyz-Turkish Manas University, Kyrgyzstan)
Peyil Esengül Kızı (Kyrgyz-Turkish Manas University, Kyrgyzstan)
Nazlı Karaca (Ataturk University, Turkey)

ORGANIZING COMMITTEE

Hasan Bulut (Chair) Fırat University, Turkey
Zakia Hammouch (Chair) Ecole Normale Supérieure of Meknes, Moulay Ismail University Morocco
Haci Mehmet Baskonus (Co-Chair) Harran University, Turkey
Carlo Cattani (Co-Chair) Tuscia University, Italy
Fevzi Erdogan (Co-Chair) (Yüzüncüyıl University, Turkey)
Mikail Et (Fırat University, Turkey)
Mehmet Bektaş (Fırat University, Turkey)
Ercan Çelik (Kyrgyz-Turkish Manas University, Kyrgyzstan)
Alaattin Esen (İnönü University, Turkey)
Mir Sajjad Hashemi (University of Bonab, Iran)
Bülent Taşel (Final International University, Girne, Cyprus)
M. Giyas Sakar (Yüzüncüyıl University, Turkey)

LOCAL COMMITTEE

Aytül GÖKÇE (Ordu University, Ordu Turkey)
Gökçe ÇAYLAK KAYATURAN (Ordu University, Ordu Turkey)
Erhan SET (Ordu University, Ordu Turkey)
Cemal BELEN (Ordu University, Ordu Turkey)
Erol TAŞ (Ordu University, Ordu Turkey)
Mehmet KORKMAZ (Ordu University, Ordu Turkey)
Yıldıray ÇELİK (Ordu University, Ordu Turkey)
Meral CANSIZ AKTAŞ (Ordu University, Ordu Turkey)
Hayal YAVUZ MUMCU (Ordu University, Ordu Turkey)
Aslıhan OSMANOĞLU (Ordu University, Ordu Turkey)
Süleyman ŞENYURT (Ordu University, Ordu Turkey)
Erdal ÜNLÜYOL (Ordu University, Ordu Turkey)
Himmet KORKMAZ (Ordu University, Ordu Turkey)
Erdinç YÜCESOY (Ordu University, Ordu Turkey)
Canan ÇİFTÇİ (Ordu University, Ordu Turkey)
İrem EROĞLU (Ordu University, Ordu Turkey)
Abdussamet ÇALIŞKAN (Ordu University, Ordu Turkey)
Fatih SAY (Ordu University, Ordu Turkey)
Ayşe GÜVEN SARIHAN (Ordu University, Ordu Turkey)
Davut CANLI (Ordu University, Ordu Turkey)
Adem TÜRKMEN (Ordu University, Ordu Turkey)
Hayrunnisa AYYILDIZ (Ordu University, Ordu Turkey)
Meral ÇELİKOĞLU (Ordu University, Ordu Turkey)
Hacı Mehmet YEŞİLTAŞ (Ordu University, Ordu Turkey)
Enes ÇİNPOLAT (Ordu University, Ordu Turkey)
Hikmet Kemaloğlu (Fırat University, Elazığ, Turkey)
Çiğdem Bektaş (Fırat University, Elazığ, Turkey)
Handan Öztekin(Fırat University, Elazığ, Turkey)
Alper Osman Öğrenmiş (Fırat University, Elazığ, Turkey)
Yavuz Altın (Fırat University, Elazığ, Turkey)
Reşat Yılmazer (Fırat University, Elazığ, Turkey)
Erdal Baş (Fırat University, Elazığ, Turkey)
Hıfsı Altınok (Fırat University, Elazığ, Turkey)
Mehmet Karay (Final Int. University, Girne, Cyprus)
Gülnür Yel (Final Int.l University, Girne, Cyprus)
Emrah Yılmaz (Fırat University, Elazığ)
Mustafa Yeneroglu (Fırat University, Elazığ, Turkey)
Münevver Tuz (Fırat University, Elazığ, Turkey)
Gülden Altay Suroglu (Fırat University, Elazığ, Turkey)
Pervin Taşel (Final Int.l University, Girne, Cyprus)
Fatma Erdogan (Fırat University, Elazığ, Turkey)
Ebru Cavlak Aslan (Fırat University, Elazığ, Turkey)
Şeyma T. Demiray (Osmaniye Korkut Ata Univ., Turkey)
M. Evren Aydın (Fırat University, Elazığ, Turkey)
Tüba Gülşen (Fırat University, Elazığ, Turkey)
Hatice Aslan (Fırat University, Elazığ, Turkey)
Ahu Ercan (Fırat University, Elazığ, Turkey)
Berat Karaagac (Adıyaman University, Adıyaman, Turkey)
Ezgi Kaya (Igdır University, Igdır, Turkey)
Mahmut İhtiyaroğlu (Van Yüzüncü Yıl Univ., Turkey)
Onur Saldır (Van Yüzüncü Yıl University, Van, Turkey)
Zelal Temel (Van Yüzüncü Yıl University, Van, Turkey)
Şakir İşleyen (Van Yüzüncü Yıl University, Van, Turkey)
Yener Altun (Van Yüzüncü Yıl University, Van, Turkey)
İbrahim Enam İnan (Fırat University, Elazığ, Turkey)
Dilek Günaydın (Fırat University, Elazığ, Turkey)
Betül Demirdağ (ODTÜ University, Ankara, Turkey)
Hajar Farhan Ismael (University of Zakho, Iraq)
Karmina K. Ali (University of Zakho, Iraq)
Sibel Şehriban Ataş (Fırat University, Elazığ, Turkey)
Aysenur Akkılıç (Assistant, Fırat University, Elazığ, Turkey)
Sibel Tarla (Assistant, Fırat University, Elazığ, Turkey)
Bahar Acay (Assistant, Fırat University, Elazığ, Turkey)
Aslı Alkan (Fırat University ,Elazığ, Turkey)
Beyhan Kemaloğlu (Fırat University, Elazığ, Turkey)
Mucahit Saydam (Fırat University, Elazığ, Turkey)
Çağrı Yılmazer (Fırat University, Elazığ, Turkey)
Merve Zeynep Gecmen (Ağrı Ibrahim Çeçen University, Ağrı, Turkey)
Necati Erdoğan (Van Yüzüncü Yıl Üniversity,Van, Turkey)
Arif Özkul (Fırat University, Elazığ, Turkey)

INTERNATIONAL COMMUNICATIONS CONSULTANT

Prof. Dr. Rengin AK (Kirklareli University, Kırklareli, Turkey)

LANGUAGE EDITORIAL COMMITTEE

Erkan Yuce (Aksaray University, Aksaray, Turkey)
Suheyla Demirkol Orak (Firat University, Elazıg, Turkey)
Fatih Mehmet Cigerci (Harran Univeristy, Sanliurfa, Turkey)
Zeki Tastan (Van Yuzuncuyil University, Van, Turkey)

PROCEEDINGS

Full version of submitted papers will be published in Special Volumes of reputed journals. Procedure, Guidelines and Checklist for the preparation and submission of papers to the Proceedings of CMES-2022 can be found in the journals websites. The journals in which selected and peer-reviewed full papers of CMES-2022 will be published are as follows:

1. FRACTAL AND FRACTIONAL JOURNAL [SCI-E],
Selected papers from CMES-2022 will be published in a special issue dedicated to the Conference entitled "**New Challenges Arising in Engineering Problems with Fractional and Integer-ll Order**". https://www.mdpi.com/journal/fractalfract/special_issues/EPFIO2022
This journal is indexed by SCI-E.

2. FRACTALS [SCI-E]
https://www.worldscientific.com/page/fractals/callforpapers

3. PROCEEDINGS OF THE INSTITUTE OF MATHEMATICS AND MECHANICS [E-SCI]
https://proc.imm.az/special/

4. FULL TEXT PROCEEDING BOOK [FREE]
At the beginning, if Authors submit FULL TEXT PAPERS, then, after getting at least two positive referee reports, FULL TEXT PAPERS will be published in FULL TEXT PROCEEDING BOOK of CMES-2022 with ISBN:77733 number. Therefore, Abstracts of these FULL TEXT PAPERS will NOT be published in ABSTRACT PROCEEDING BOOK of CMES-2022.

5. ABSTRACT [FREE]
If Authors submit ABSTRACT TEXTS, then, after getting referees evaluations for these abstracts, they will be published in ABSTRACT PROCEEDING BOOK of CMES-2022. For FULL TEXT PAPERS, Authors have to submit their FULL TEXT PAPERS online via submission system of CMES-2022. These FULL TEXT PAPERS will be published in FULL TEXT PROCEEDING BOOK of CMES-2022 after getting at least two positive reports.

6.TURKISH JOURNAL OF SCIENCE, [FREE]
"Participants of CMES 2022 can submit their good quality papers to Turkish Journal of Science. After the peer review process, the papers will be published at TJOS. The authors must write "CMES 2022" as comments to the editor. (Editor in Chief: Dr. Ahmet Ocak AKDEMİR) For on-line submission: https://dergipark.org.tr/tr/pub/tjos

7.TURKISH JOURNAL OF INEQUALITIES, [FREE]
"Participants of CMES 2022 can submit their good quality papers to Turkish Journal of Inequalities. Selected papers will be published at TJI after the peer review process. The participants can send their papers to erhanset@tjinequality.com. The authors must write "CMES 2022" as the subject. (Editor in Chief: Prof. Dr. Erhan SET) http://tjinequality.com/

8. MATHEMATICS IN NATURAL SCIENCE (MNS)
*Authors can submit their full text paper directly to the journal by using the following link https://www.isr-publications.com/mns

9. MATHEMATICS IN ENGINEERING, SCIENCE AND AEROSPACE (MESA), [FREE, SCOPUS]
"Selected papers will be published after peer review in the Journal of Mathematics in Engineering, Science and Aerospace (MESA)" (Editor in Chief: Prof. Seenith Sivasundaram) http://nonlinearstudies.com/index.php/mesa

10. APPLIED MATHEMATICS AND NONLINEAR SCIENCES, [SCOPUS]
Participants of CMES 2022 can submit their high quality full text papers to Applied Mathematics and Nonlinear Sciences by selecting CMES-2022 under the Select Article Type Menu. (https://www.editorialmanager.com/amns/default.aspx)

11.MATHEMATICAL MODELLING AND NUMERICAL SIMULATION WITH APPLICATIONS (MMNSA)
The Special Issue on "Advanced Methods of Modelling and Numerical Computation in Science and Engineering". Authors can submit their full text paper directly to the journal by using the information provided in the following link https://mmnsa.org/index.php/mmnsa/special_issues/SI-CMES2022

12.SPECIAL ISSUE "ADVANCES IN MATRIX TRANSFORMATIONS, OPERATORS AND SYMMETRY"
(SCIE) https://www.mdpi.com/journal/symmetry/special_issues/Advances_Matrix_Transformations_Operators_Symmetry

13. YUZUNCU YIL UNIVERSITY JOURNAL OF THE INSTITUTE OF NATURAL AND APPLIED SCIENCES
(The journal indexed by TR-Dizin)
https://dergipark.org.tr/tr/pub/yyufbed

14. PEDAGOGICAL PERSPECTIVE (PEDPER)
Pedagogical Perspective **(PedPer)** is an international, double blind reviwing, non-profit, professional scientific journal. PedPer is a journal that accepts manuscripts related to pedagogy and education.
http://pedagogicalperspective.com/

15. ERZURUM TECHNICAL UNIVERSITY JOURNAL OF MATHEMATICS (ETU J MATH)
https://dergi.erzurum.edu.tr/etujm

PLENARY & INVITED TALKS





The interaction of minimizers of functionals and Vanishing Mean Oscillation functions

Maria Alessandra Ragusa
University of Catania, Italy
mariaalessandra.ragusa@unict.it

Abstract We show the advances on some regularity problems related to minimizers

$$u(x) : \Omega \rightarrow R^n$$

of some quadratic and non quadratic growth functionals. About the dependence on the variable x is of the integrand function A(x, u, p) of the functionals it belongs to the Vanishing Mean Oscillation class, as a function of x. Then, is pointed out that the continuity of A(x, u, p), with respect to x, is not assumed.

Keyword: Regularity, Minimizers of functionals, Vanishing Mean Oscillation functions.

References

[1] Ragusa M.A., Tachikawa A., Regularity for minimizers for functionals of double phase with variable exponents , Adv. Nonlinear Anal., 9, 710-–728 (2020);

[2] Ragusa M.A., Tachikawa A., Boundary regularity of minimizers of p(x)-energy functionals, Annales de L'Institut Poincarè - Analyse non lineaire, 33, 451--476, (2016).



MODIFIED AND GENERALIZED FRACTIONAL OPERATORS: THEORY AND APPLICATIONS

Dumitru Baleanu
Çankaya University, Turkey and Institute of Space Sciences, Romania
dumitru@cankaya.edu.tr

Abstract

It is wel known that the fractional calculus deals with the study of so-called fractional order integral and derivative operators over real or complex domains, and their applications.In my talk I will present some new trends in the field of modified and generalized fractional operators.Illustrative examples will be presented.

Keywords: fractional calculus, modified fractional operators, generalized fractional operators

REFERENCES

[1] D. Baleanu, K. Diethelm, E. Scalas, J. J. Trujillo, Fractional calculus: models and numerical methods, World Scientific, New York, Second Edition,2016.
[2] M. Caputo and M. Fabrizio, On the singular kernels for fractional derivatives. Some applications to partial differential equations,Progress in Fractional Differentiation and Applications 7, No. 2, 79-82, 2021
[3] R. Nigmatullin, D. Baleanu, A. Fernandez, Balance equations with generalised memory and the emerging fractional kernels, Nonlinear Dynamics, 2021
[4] H. T. Nguyen Huy, D. Baleanu, N. T. Tran Ngoc,D. O'Regan,N. H. Can, Final value problem for nonlinear time fractional reaction-diffusion equation with discrete data, Journal of Computational and Applied Mathematics 376, Article Number: 112883, 2020
[5] D.Baleanu, A.Fernandez, A.Akgul, On a fractional operator combining proportional and classical differintegrals, Mathematics,8(3),Article Number 360,2020
[6] D. Baleanu,, S.S. Samaneh, J. H. Asad, Jihad, A. Jajarmi, Hyperchaotic behaviors, optimal control, and synchronization of a nonautonomous cardiac conduction system, Adv. Differ. Eq., 2021(1), Article Number: 157, 2021.



FIXED POINTS AND COUPLED FIXED POINTS IN B-METRIC SPACES VIA GRAPHICAL CONTRACTIONS

Liliana GURAN
Vasile Goldiș Western University of Arad, Arad, Romania
guran.liliana@uvvg.ro

Abstract

In this paper some existence and stability results for cyclic graphical contractions in complete metric spaces are given. Moreover, we discuss a vectorial case of our main result. Some applications to coupled fixed point problems are also derived.

Keywords: Existence and stability results for cyclic graphical contractions in complete metric spaces

REFERENCES

[1] I.A. Bakhtin, The contraction mapping principle in quasimetric spaces, Funct. Anal., Unianowsk Gos. Ped. Inst. 30(1989) 26-37.
[2] V. Berinde, Generalized contractions in quasimetric spaces, Seminar on Fixed Point Theory, 1993, 3-9.
[3] V. Berinde, Generalized coupled fixed point theorems for mixed monotone mappings in partially ordered metric spaces, Nonlinear Anal. 74 (2011) 7347-7355.
[4] M. Bota, A. Molnar, Cs. Varga, On Ekeland's variational principle in b-metric spaces, Fixed Point Theory, 12(2011), No. 1, 21–28.
[5] S. Czerwik, Contraction mappings in b-metric spaces, Acta Mathematica et Informatica Univ. Ostraviensis 1(1993) 5-11.



A COMPARISON ON SOME INTEGRAL TRANSFORMS

Hossein Jafari[1,2]

[1]Department of Mathematics, University of Mazandaran, Babolsar, Iran

[2]Department of Mathematical Sciences, University of South Africa, UNISA0003, South Africa

jafari.usern@gmail.com,

Abstract

Integral transforms are important to solve real problems. Appropriate choice of integral transforms helps to convert differential equations as well as integral equations into terms of an algebraic equation that can be solved easily.

During last two decades many integral transforms in the class of Laplace transform are introduced such as Sumudu, Elzaki, Natural, Aboodh, Pourreza, Mohand, G_transform, Sawi and Kamal transforms.

In this work, we compare some integral transforms in the class of Laplace transform which are introduced during last few decades. After that we propose a general integral transforms which is covered all of those integral transforms. Also, we discuss about combanition of these integral transforms and decomposition methods.

Keywords: Integral transform; Decomposition methods; Iterative method.

REFERENCES

1. H. Jafari, A new general integral transform for solving fractional integral equation, Journal of Advanced Research, Vol: 32, 133-138, 2021.



MATHEMATICAL METHODS IN INDUSTRIAL MATHEMATICS: A REVIEW FROM RECENT CASE STUDIES

Erhan Coşkun [1]

[1] Department of Mathematics, Karadeniz Technical University, Trabzon, Turkey

erhan@ktu.edu.tr

Abstract In this talk, we describe our experience of identification, mathematical formulation and handling of industrial problems posed by firms having active research and development units. We highlight common features of industrial problems from OECD report of industrial mathematics [1], describe basic mathematical tools and skills needed to engage in problems of industry and discuss some relevant case studies including the ones from the Euro-Asian Study Group with Industry[2] held in Trabzon, Turkey. Our aim is to convince young mathematicians that they can indeed develop their skills to tackle industrial problems, which will not only help industry to gain compatitive advantage but also lead to having original research areas that are more likely to be supported from private and government institutions.

Keywords: Industrial mathematics; mathematical modeling; study groups with industry.

REFERENCES

1. Mathematics in Industry, OECD Global Science Forum report, https://www.oecd.org/science/inno/41019441.pdf
2. Coskun, E. et al., *Gabor Filter Selection and Computational Processing for Emotion Recognition.* Study Group: ESGI 100. 7–11 April, 2014, Oxford, UK.
DOI: 10.33774/miir-2021-5q5j6.



SHANNON–WHITTAKER–KOTEL'NIKOV'S THEOREM GENERALIZED

JUANLUİS GARCÍA GUİRAO
Technical University of Cartagena, Spain
juan.garcia@upct.es
http://www.jlguirao.es

Abstract The aim of the this talk provide a generalization of the classical Shannon-Whittaker-Katel'niko's theorem for a class of non band-limited signals which plays a central role in the signal theory, the Gaussian map is the unique function which reachs the minimum of the product of the temporal and frecuential width. This solves a conjecture stated by Boas in 1972.

Keywords: Shannon–Whittaker–Kotel'nikov's Theorem; recomposi- tion of chemical products, signal theory.

REFERENCES

[1] R.P. Boas Jr., Summation formulas and band--limited signals, Tohoku Math. J., 24, (1972), 121-125.



COMPLEX DYNAMICS OF A RATIO-DEPENDENT POPULATION MODEL: STABILITY, BIFURCATIONS AND CHAOS

HUSEYIN MERDAN
TOBB University of Economics and Technology, Department of Mathematics, Ankara, Turkey

merdan@etu.edu.tr

Abstract Nonlinear dynamical behaviors of a prey-predator system with Leslie type will be presented. First, the stability analysis will be given. Second, the bifurcations and chaotic behavior of the model will be discussed. Finally, numerical simulations will be shown to support and extend the theoretical results. The results obtained will be interpreted from the biological point of view.

Keywords: Prey-predator model, stability analysis, bifurcations, chaotic behavior

References

[1] E. Karaoglu and H. Merdan, Hopf bifurcations of a ratio-dependent predator-prey model involving two discrete matutaion time delays, Chaos, Soliton and Fractals, 68, 159-168, (2014)

[2] P. Baydemir, H. Merdan, E. Karaoglu and G. Sucu, G, Complex dynamics of a discrete-time prey-predator system with Leslie type: Stability, bifucation analyses and chaos, International Journal of Bifurcation and Chaos, Vol. 30, No. 10, 2050149 (2020).

[3] S. Yildiz, S. Bilazeroglu and H. Merdan, Stability and bifurcation analyses of a discrete Lotka-Volterra type predator-prey system with refuge effect, Preprint, (2022)

[4] S. Wiggins, Introduction to Applied Nonlinear Dynamical Systems and Chaos, Springer-Verlag, New York, (2003)



TYPE-2 FUZZY SYSTEMS AND FUTURE TRENDS FOR TYPE-3: THEORY AND APPLICATIONS

Oscar CASTILLO
Tijuana Institute of Technology, Tijuana, Mexico.
ocastillo@tectijuana.mx

Abstract: Type-2 fuzzy systems are powerful intelligent models based on the theory of fuzzy sets, originally proposed by Prof. Zadeh. Most real-world applications up to now are based on type-1 fuzzy systems, which are built based on the original (type-1) fuzzy sets that extend the concept of classical sets. Type-2 fuzzy sets extend type-1 fuzzy sets by allowing the membership to be fuzzy, in this way allowing a higher level of uncertainty management. Even with the current successful applications of type-1 fuzzy systems, now several papers have shown that type-2 is able to outperform type-1 in control, pattern recognition, manufacturing and other areas. The key challenge in dealing with type-2 fuzzy models is that their design has a higher level of complexity, and in this regard the use of bio-inspired optimization techniques is of great help in finding the optimal structure and parameters of the type-2 fuzzy systems for particular applications, like in control, robotics, manufacturing and others. Finally, the prospects for the future trends and applications of type-3 fuzzy logic will be discussed.

Keywords: Type-2 Fuzzy systems, Type-1 fuzzy systems, Fuzzy Logic.



AN EVALUATION ON THE RELATIONSHIP BETWEEN HUMANITY'S ADVENTURE OF KNOWLEDGE AND HIKMAH

Vatan KARAKAYA
Department of Mathematical Engineering, Yildiz Technical University, Istanbul, Turkey
Ahi Evran University, Kirsehir, Turkey
vkkaya@yahoo.com

Abstract


The main purpose of this study is to investigate the knowledge and wisdom relationship based on the definitions of God and knowledge of certain civilizations within the context of written history. This relationship is tried to be explained under the definitions of God and knowledge derived from eminent thinkers of Ancient Greece, Islamic and Modern Western civilizations. The relationship between knowledge and wisdom in those two civilizations are examined with regard to the concept of 'hikmah' defined in Islamic civilization. After that, a general comparison was made on the topics of God and knowledge definitions of the founder thinkers of these three civilizations. As a result of this comparison, it has been concluded that the definitions of God and knowledge definitions of Ancient Greece and Modern Western civilizations are similar, and these two concepts do not match the content of 'hikmah'. Main reason behind this discrepancy is concluded as the unique conglomeration of tasawwuf, kalam and Aristotelian philosophy in Islamic civilization. In Islamic civilization, these three thought traditions assimilated the thought before them, formed their own tradition and laid the groundwork for the next civilization.

Keywords: Belief in God, definition of knowledge, knowledge and hikmah.

REFERENCES

Abdurrahman, T. (2021). Amel sorunsalı bilim ve düşüncenin pratik temelleri üzerine bir araştırma. (T. Uluç, Çev.). Pınar Yayınları.

Aristoteles. (1999). Metaphysics. A new transalation by Joe Sachs. Green Lion Press.
Descartes, R. (2019). Yöntem üzerine konuşma. (Ç. Dürüşken, Çev.). Alfa Yayınları.



A NEW SEQUENCE SPACE OF WEIGHT g

Ekrem SAVAŞ
Department of Mathematics, Usak University, Usak, Turkey
ekremsavas@yahoo.com

ABSTRACT. Recall that $f : [0,\infty) \to [0,\infty)$ is a modulus function if it satisfies the following properties.

i): $f(x) = 0$ if and only if $x = 0$,

ii): $f(x+y) \leq f(x) + f(y)$,

iii): f is increasing,

iv): f is continuous from the right at 0.

In this paper, we define a new sequence space by using modulus function and also investigate some relations the spaces between $[\hat{V}^g, \lambda, f, t]$ and $\hat{S}_\lambda^g$ where $g : [0,\infty) \times [0,\infty) \to [0,\infty)$, $g((x_{nm}) \to \infty$ for any sequence (x_{nm}) in $[0,\infty) \times [0,\infty)$ with $x_{nm} \to \infty$.

REFERENCES

[1] S. Banach, Théorie des Opérations lieaires, Chelsea Publ. Co., New York, 1955.

[2] S. Bhunia, P. Das and S. K. Pal, Restricting statistical convergence, Acta Math. Hungar. 134 (1-2) (2012), 153-161.

[3] Cakalli, H.; Savas, E. Statistical convergence of double sequences in topological groups. J. Comput. Anal. Appl. 12 (2010), no. 2, 421-426.

[4] R. Colak, Statistical convergence of order α, Modern methods in Analysis and its Applications, New Delhi, India, Anamaya Pub., (2010), 121-129.

[5] R. Colak, Y. Altin, Statistical convergence of double sequences of order α, Journal of Function Spaces, Volume 2013, Article ID 682823.

[6] M. Isik, Y. Altin, and Altin $f(\alpha,\mu)$-statistical convergence of order α for double sequences,J. Ineq. Appl. (2017) 2017:246.

[7] H. Nakano, Modular sequence spaces, Proc. Japan Acad., 27 (1953), 508 - 512.

[8] E. Savaş, Strong almost convergence and almost λ-statistical convergence, Hokkaido Math. J., 29 (3)(2000), 531 - 536.

TYPE	AUTHORS	TITLES	PAGE
Online	Esma Ulutaş	ON SOLUTIONS STOCHASTIC NONLINEAR PARTIAL DIFFERENTIAL EQUATIONS WITH CONFORMABLE DERIVATIVES	43
Online	Ilhame Amirali, Ece Eroğlu	A BRIEF SURVEY ON NUMERICAL SOLUTIONS OF SINGULARLY PERTURBED PROBLEMS	44
Oral	Rifat Çolak, Çiğdem Asma Bektaş	(□mv)u - GENERALIZED WEIGHTED STATISTICAL CONVERGENCE	45
Online	Iqbal H. Jebril, Najat M. Abed Abdelqader	HYERS-ULAM STABILITY OF QUANTUM LOGIC FUZZY IMPLICATION	46
Oral	Besik Dundua	PρLog: A LANGUAGE FOR EXACT AND APPROXIMATE REASONING	47
Oral	Esin İlhan	SOME NEW SOLITON SOLUTIONS OF TIME FRACTIONAL RESONANT DAVEY-STEWARTSON EQUATIONS	48
Oral	Mehmet Korkmaz	PERFORMANCES OF FITS WHEN SOME GROWTH MODELS ARE HYPERBOLIC	49
Oral	Arzu Aykut, Ercan Çelik	THE CONSECUTIVE SUBSTITUTION METHOD FOR THE BOUNDARY VALUE PROBLEM	50
Oral	Himmet Korkmaz , Kevser Kurucu Bayhan, Mahmut Kertil	MATHEMATICS TEACHERS' VIEWS ABOUT THE TRANSITION FROM ONLINE TEACHING TO FACETO-FACE EDUCATION	51
Oral	S.Şule Şener Kılıç Ercan Çelik Hasan Bulut	SOLUTION OF (2+1)-DIMENSIONAL QUANTUM ZAKHAROV-KUZNETSOV EQUATIONS USING THE THE MODIFIED EXP(−Ω(ξ))-EXPANSION FUNCTION METHOD	52
Oral	Özlem Defterli	ABOUT NEW MATHEMATICAL APPROACHES IN FRACTIONAL BIOLOGICAL MODELLING	53
Oral	Liliana Guran, Monica-Felicia Bota, Gabriela Petrusel	FIXED POINTS AND COUPLED FIXED POINTS IN B-METRIC SPACES VIA GRAPHICAL CONTRACTIONS	54
Online	Anass El karkri, Imane El mhamdi, Zakaria El malki, Mohamed bouachrine, Jean-Marc Sotiropoulos	SIMULATION OF A LIGHT-EMITTING DIODE BASED ON ORGANIC MATERIALS UNDER ATLAS-SILVACO	55
Oral	Esin İlhan	NOVEL SOLUTION FOR DETERMINISTIC TYPHOID FEVER MODEL WITHIN THE FRAME OF FRACTIONAL ORDER	56
Oral	Aslı Alkan, Tolga Aktürk, Hasan Bulut	THE PERIODIC WAVE SOLUTIONS OF CONFORMABLE TIME-FRACTIONAL MATHEMATICAL MODEL BY ANALYTICAL METHOD	57
Oral	Zeynep Yılmaz, Selahattin Maden, Aytül Gökçe	ANALYSING ONE PREY AND TWO PREDATORS MATHEMATICAL MODEL WITH HOLLING TYPE II RESPONSE AND FADING MEMORY	58
Online	Yusuf Ziya Yüksel, Şeref Doğuşcan Akbaş	STRESS ANALYSIS OF MULTI-LAYERED SOIL MEDIUM BY USING FINITE AND INFINITE ELEMENTS	59

Online	Kürşad Özkan, Serkan Özdemir, Ali Şenol, Ecir Uğur Küçüksille	A NEW SPECIES RICHNESS MEASURE IMPROVED FROM MARGALEF AND MENHINICK	60
Oral	M. Younus Bhat	WIGNER-VILLE DISTRIBUTION AND AMBUGUITY FUNCTION ASSOCIATED WITH QUATERNION OFFSET LINEAR CANONICAL	61
Oral	Hasan Bulut, Sümeyye Kılbitmez, Tukur A Sulaiman, Abdullahi Yusuf	SOLITARY WAVE SOLUTIONS TO A NONLINEAR MATHEMATICAL MODEL	62
Oral	Erdem Türkeli	THE WIND LOAD CALCULATION OF A MASONRY MINARET BY USING DIN1056 STANDARD	63
Oral	Hajar F. Ismael, Arif Özkul, Hasan Bulut	ON THE COMPLEX SOLITON SOLUTIONS TO THE HIROTA–SATSUMA– ITO EQUATION	64
Online	Hayatem Hamal, Pembe Sabancıgil	KANTOROVICH TYPE (p,q)-GENERALIZATION OF BERNSTEIN OPERATORS	65
Oral	Derya Deniz, Mücahit Saydam	ON THE SOLUTION OF PAVLOV EQUATION BY THE VARIATIONAL ITERATION METHOD	66
Online	Sadullah Bulut, Muhammed Yiğider	THE COMPARISON OF NUMERICAL SOLUTIONS OF COMFORMABLE AND □ COMFORMABLE FRACTIONAL DIFFERENTIAL EQUATIONS USING NEURAL NETWORK METHOD BY PYTHON	67
Online	Alican Çıvğa, Serkan Özdemir, Serkan Gülsoy	MODELING POTENTIAL DISTRIBUTION OF CAPPARIS SPINOSA IN BABADAG REGION	68
Oral	Furkan Şimşek, Şükrü Kitiş	ARDUINO BASED DIGITAL TACHOMETER WITH TRACKING PROGRAM	69
Oral	Elif Başkaya	AN APPROXIMATION METHOD FOR THE EIGENVALUE OF A STURM-LIOUVILLE PROBLEM	70
Oral	Engin Özkan, Mine Uysal	ON A GENERALIZATION OF MERSENNE HYBRID NUMBERS	71
Online	Hanife Göker, Afrah Said	COMPARATIVE ANALYSIS OF MACHINE LEARNING ALGORITHMS FOR EARLY DIAGNOSIS OF ALZHEIMER'S DISEASE	72
Online	Elif Otkun Çevik	ON THE DIFFERENCE BETWEEN NUMERICAL AND SPECTRAL RADII FOR SOME ANALYTIC FUNCTIONS OF OPERATOR	73
Online	Yusuf Ziya Yüksel, Şeref Doğuşcan Akbaş	STATIC RESPONSES OF A CAVITY IN A HALF-SPACE BY USING FINITE ELEMENT METHOD	74
Oral	Asifa Poom Kumam	A FRACTIONAL MODEL TO INVESTIGATE SHAPE EFFECTS ON THERMAL BEHAVIOR OF BRINKMAN-TYPE FERROFLUID UNDER THERMAL RADIATION AND HEAT INJECTION/CONSUMPTION	75

Online	Ayşegül Tekeş	DETERMINATION OF INDICATOR PLANT SPECIES BY INDICATOR ANALYSIS IN RED PINE (PINUS BRUTIA TEN.) AND BLACK PINE (PINUS NIGRA J.F.ARNOLD) FORESTS IN BOZDAGLAR REGION	76
Oral	Zulqurnain Sabir	SOLUTION OF SINGULAR BOUNDARY VALUE PROBLEMS USING ARTIFICIAL NEURAL NETWORKS	77
Oral	Gülnur Yel, Beyhan Kemaloglu, Hasan Bulut	A NOVEL ANALYTICAL APPROACH TO THE BENJAMIN–ONO EQUATION	78
Online	Muhammad Abubakar Isah, Asıf Yokuş	A NOVEL TECHNIQUE TO CONSTRUCT EXACT SOLUTIONS FOR THE COMPLEX GINZBURG-LANDAU EQUATION USING QUADRATIC-CUBIC NONLINEARITY	79
Oral	Zulqurnain Sabir, Ayşe Nur Akkılıç, Hasan Bulut	A STOCHASTIC COMPUTING HEURISTIC SCHEME FOR SOLVING A NOVEL THIRD ORDER PERTURBED DELAY DIFFERENTIAL EMDEN-FOWLER MODEL	80
Oral	Yusuf Gurefe, Yusuf Pandır, Tolga Akturk	ANALYSIS OF THE EFFECT OF THE CORIOLIS CONSTANT ON THE GPKDV EQUATION	81
Online	Rabia Kevser Yılmaz, Oğuz Yağcı Recep Şahin	DEGENERATE ARA TRANSFORM	82
Oral	Eren Çolak, Ali Keleş	NUMERICAL INVESTIGATION OF PARTICLE DISTRIBUTION IN A CYLINDRICAL CHANNEL	83
Oral	Talha Anwar, Asifa Poom Kumam	FRACTIONAL MODELING AND EXACT SOLUTIONS TO ANALYZE THERMAL PERFORMANCE OF FE3O4-MOS2-WATER HYBRID NANOFLUID FLOW OVER AN INCLINED SURFACE	84
Online	Kürşad Özkan, Mehmet Güvenç Negiz, İbrahim Özdemir	STRUCTURAL DIVERSITY ANALYSIS USING SHANNON'S ENTROPY AND INFORMATION MEASURES	85
Online	Afrah Said, Hanife Göker	AUTISM SPECTRUM DISORDER DETECTION USING MACHINE LEARNING ALGORITHMS	86
Online	Yücel Baltürk, Hüseyin Demir, İnci Çilingir Süngü	ON NUMERICAL SOLUTION OF LID DRIVEN PROBLEM IN A SQUARE CAVITY FILLED WITH NANOFLUID	87
Oral	Himmet Korkmaz, Serbay Duran	INVESTIGATING THE DIFFICULTIES OF REFUGEE STUDENTS IN MATHEMATICS COURSE	88
Oral	Aslı Alkan, Tolga Aktürk, Hasan Bulut	THE PERIODIC WAVE SOLUTIONS OF CONFORMABLE TIME-FRACTIONAL MATHEMATICAL MODEL BY ANALYTICAL METHOD	89
Oral	Mohammad Younus Bhat	UNCERTAINTY PRINCIPLES FOR THE SHORT-TIME NON-SEPARABLE LINEAR CANONICAL TRANSFORM	90
Online	Yasin Bozkurt, Mehmet Tahir Gulluoglu, Haci Mehmet Baskonus	SOME RESULTS FOR A NONLINEAR TRANSMISSION LINE ARISING IN THE ELECTRONIC CIRCUIT	91

Oral	Yasemin Akgün	THERMAL INSULATION PERFORMANCES OF FOAMED MORTARS CONTAINING BENTONITE BLENDED CEMENTS	92
Oral	Arran Fernandez	THINK BEFORE YOU DEFINE: ON CONJUGATION RELATIONS AND OTHER CONNECTIONS	93
Online	Fatih Coşkun, Beytullah Topçu, M. Fatih Akay	DEVELOPMENT OF A NATIONAL SMART METER SYSTEM	94
Oral	Zakaria El Malki, Anass El Karkri, Imane EL Mhamedi, Mohammed Bouachrine, Françoise SereinSpirau	COMPUTATIONAL AND SIMULATION STUDY OF ORGANIC SOLAR CELLS BASED ON DONOR – ACCEPTOR MOLECULAR MATERIAL	95
Oral	Emel Biçer	APPLICATION OF MAHGOUB TRANSFORM TO HYERS-ULAM STABILITY OF PARTIAL DIFFERENTIAL EQUATIONS	96
Online	Imane EL Mhamedi, Anass El Karkri, Zakaria El Malki, Mohammed Bouachrine, Françoise Serein-Spirauc	HIGH-PERFORMANCE OF ORGANIC SOLAR CELLS BASED ON D1- BT-EDOT-BT-D2-A/PCBM STRUCTURES	97
Oral	Guillaume Leduc	EDGEWORTH SERIES EXPANSION FOR OPTION PRICES	98
Oral	Mehmet Yavuz, Fırat Evirgen, Fatma Özköse, Necati Özdemir	AN OPTIMAL CONTROL STRATEGY FOR COVID-19 USING REAL DATA	99
Online	Kh. R. Mamedov, U. Demirbilek	SPECTRAL EXPANSIONS FOR A CLASS SECOND ORDER DIFFERENTIAL OPERATOR WITH EIGENVALUE PARAMETER CONTAINED IN THE BOUNDARY CONDITIONS	100
Oral	Onur Alp İlhan, Jalil Manafian	APPLICATION OF tan((Φ(ξ))/2)-EXPANSION SCHEME FOR SOLVING TWO NONLINEAR PARTIAL DIFFERENTIAL EQUATIONS ARISING IN CHEMICAL AND BIOLOGICAL PHENOMENA	101
Online	Özlem Kırcı	ANALYSIS OF THE SOLUTIONS OF A SPACE-TIME FRACTIONAL MATHEMATICAL MODEL WITH AN ANALYTICAL METHOD	102
Oral	Hasan Bulut, Sümeyye Kılbitmez, Tukur A Sulaiman, Abdullahi Yusuf	SOLITARY WAVE SOLUTIONS TO A NONLINEAR MATHEMATICAL MODEL	103
Online	Karmina K. Ali, Sibel Sehriban Ataş, Hasan Bulut	EXACT SOLUTIONS TO THE COUPLED-HIGGS EQUATION WITH NEUTRAL SCALAR MESONS COOPERATING IN A SYSTEM	104

Online	Mustafa Özdemir, Eralp Osman Erdogan	FORMULATION OF A 7-DOF HALF-CAR MODEL WITH NONLINEAR SPRINGS AND DAMPERS FOR RIDE COMFORT ANALYSIS OF IN-WHEEL-MOTOR ELECTRIC VEHICLES	105
Oral	Muharrem Tuncay Gencoglu	ANALYSIS OF CRYPTOGRAPHIC HASH FUNCTIONS FOR BLOCKCHAIN TECHNOLOGY	106
Oral	Enes Ata	APPLICATION OF FRACTIONAL OPERATORS TO MODIFIED SPECIAL FUNCTIONS	107
Online	Özge Erden	A STOCHASTIC MODEL FOR THE ENERGY EFFICIENT SERVER CONSOLIDATION	108
Online	Gamze Özel Kadılar, Barış Burçin Demir	A NEW GENERALIZED DISTRIBUTION FOR ESTIMATION OF FLOOD AND EARTHQUAKE SIZES IN TURKEY	109
Online	Nigar Yildirim Aksoy, Ercan Celik, Merve Zengin	AN OPTIMAL CONTROL PROBLEM FOR n-DIMENSIONAL SCHRODINGER EQUATION WITH SPECIFIC GRADIENT TERM	110
Online	Gamze Özel Kadılar, Ceren Ünal	MARKOV RENEWAL MODEL APPLIED TO THE ESTIMATION OF EARTHQUAKE OCCURRENCES	111
Oral	Sibel Özer, Selçuk Kutluay	AN EFFICIENT LINEARIZED APPROXIMATION TECHNIQUE FOR THE NON-LINEAR TWO-DIMENSIONAL COUPLED BURGERS EQUATION	112
Oral	Hasan Bulut, Ercan Çelik, Salisu Ibrahim, Tukur Abdulkadir Sulaiman, Abdullahi Yusuf	SOLUTIONS TO THE (N+1)-DIMENSIONAL NLSE EQUATION WİTH M-TRUNCATED DERIVATIVE	113
Oral	Sibel Tarla, Karmina K. Ali, Resat Yilmazer	NEW EXACT SOLUTIONS OF THE COUPLED-HIGGS EQUATİON USING THE PHI^6-MODEL EXPANSION METHOD	114
Oral	Şükrü Kitiş, Hanife Göker	PERFORMANCE EVALUATION OF MACHINE LEARNING ALGORITHMS FOR DETECTING EATING AND PHYSICAL STATUS BASED OBESITY STAGES	115
Online	Mustafa Mutlu, Çetin Kurnaz	RECTANGULAR MICROSTRIP ANTENNA DESIGN FOR WIRELESS TECHNOLOGIES	116
Online	Mehmet Çağrı Yılmazer, Emrah Yılmaz, Sertac Goktas Mikail Et	STATISTICAL CONVERGENCE ON MULTIPLICATIVE CALCULUS	117
Oral	Muhittin Evren Aydin, Seyma Kaya	USE OF FRACTIONAL DERIVATIVE IN DIFFERENTIAL GEOMETRY	118
Oral	Karmina K. Ali, Sibel Tarla, Resat Yilmazer	DYNAMIC BEHAVIOR OF THE FOKAS SYSTEM ARISING IN MONOMODE OPTICAL FIBERS THROUGH THE UNIFIED AUXILIARY EQUATION METHOD	119
Online	Melisa Bal, Gizem Topaloğlu, Sevtap Erdem, M. Fatih Akay	ARTIFICIAL INTELLIGENCE BASED DEMAND PREDICTION MODELS	120
Online	Usman Younas, Jingle Ren	DIVERSITY OF WAVE STRUCTURES TO THE CONFORMABLE FRACTIONAL DYNAMICAL MODEL	121

Oral	Muhammad Abubakar Isah, Asıf Yokuş	OPTICAL SOLITONS OF COMPLEX GINZBURG-LANDAU EQUATION HAVING DUAL POWER NONLINEAR FORM USING Q6-MODEL EXPANSION APPROACH	122
Online	Serkan Özdemir, Ecir Uğur Küçüksille, Serkan Gülsoy, Özdemir Şentürk, Mehmet Güvenç Negiz, Halil Süel, Ahmet Mert, Kürşad Özkan	MODELING OF SPECIES DISTRIBUTIONS WITH DEEP LEARNING METHOD	123
Oral	Halil Anaç	THE NEW SOLUTIONS OF CONFORMABLE FRACTIONAL WHITHAM-BROER-KAUP EQUATIONS	124
Oral	Haci Mehmet Baskonus, Wei Gao	INVESTIGATION OF OPTICAL SOLITONS TO THE NONLINEAR COMPLEX KUNDU-ECKHAUS AND ZAKHAROV-KUZNETSOV-BENJAMIN-BONAMAHONY EQUATIONS IN CONFORMABLE	125
Online	Mustafa Kara	SCATTERED GEOMETRİCAL OPTICS FIELD BY A PERFECTLY CONDUCTING HALF PLANE BETWEEN TWO ISOREFRACTIVE MEDIA	126
Oral	Tolga Aktürk, M. Kübranur Dikici	ANALYSIS OF TRAVELING WAVE SOLUTIONS OF NONLINEAR PARTIAL DIFFERENTIAL EQUATIONS	127
Oral	Nazlı Karaca	SOME CONVERGENCE RESULTS FOR A NEW GENERALIZED NONEXPANSIVE MAPPINGS IN UNIFORMLY CONVEX HYPERBOLIC SPACE	128
Oral	Ayşe Güven Sarıhan, Oscar F. Bandtlow	A FIXED POINT THEOREM IN B-METRIC SPACES AND AN APPLICATION TO SPECTRAL ASYMPTOTICS OF SOLUTIONS OF OPERATOR EQUATIONS	129
Online	Özgül İlhan	A TURBULENCE MODEL WITH SLIP WITH FRICTION BOUNDARY CONDITION	130
Online	Zeynep Özat, Bayram Çekim	A NEW GENERALIZATION OF THE APOSTOLEULER AND APOSTOL-GENOCCHI POLYNOMIALS	131
Online	Neslihan Biricik, Bayram Çekim	GENERALIZED OF HERMITE MATRIX POLYNOMIALS AND SOME PROPERTIES	132
Oral	Harun Biçer	ON THE PERIODIC SOLUTIONS OF A NONLİİNEAR DIFFERENTIAL EQUATION WITH VARIABLE DELAYS	133
Online	Muhammet Enes Durmaz, Gabil Amirali	A UNIFORM NUMERICAL APPROACH FOR SINGULARLY PERTURBED FREDHOLM INTEGRO-DIFFERENTIAL EQUATION	134
Online	Belgin Özer, Fatima Dakkak	THE EXAMINATION OF THE QUOTIENT OF NUMERICAL SEMIGROUP WITH RF-MATRICES	135
Oral	Arzu Bal, Numan Yalçın, Mutlu Dedetürk	ON SOLUTIONS OF INTEGRAL EQUATIONS IN GEOMETRIC ANALYSIS	136
Online	Numan Yalçın, Mutlu Dedetürk	ON SOLUTIONS OF FIRST AND SECOND ORDER LINEAR DIFFERENTIAL EQUATIONS IN GEOMETRIC ANALYSIS	137

Online	Kübra Çetin, Bayram Çekim	SOME PROPERTIES OF THE PARAMETRIC KIND OF FUBINI POLYNOMIALS WITH THREE VARIABLES OF ORDER r	138
Online	Bilge Zehra Sergi, Gürhan İçöz	ON A NEW TYPE OPERATORS INCLUDING THE APPELL POLYNOMIALS AND ITS APPROXIMATION PROPERTIES	139
Online	Hasan Bulut, Derya Arslan	THE NUMERICAL SOLUTION OF SINGULARLY PERTURBED DELAY INTEGRO-DIFFERENTIAL EQUATION	140
Oral	Çimen Gül Kuluşaklı, Mustafa Birinci	FINE GRINDING OF MARBLE WASTE: PRELIMINARY TEST RESULTS	141
Online	İnci Çilingir Süngü, Emre Aydın	ON THE SEMI-ANALYTICAL METHODS FOR THE FIFTH ORDER NONLINEAR KdV EQUATIONS	142
Oral	Gül Deniz Çaylı	A NEW CONSTRUCTION METHOD FOR TRIANGULAR NORMS AND TRIANGULAR CONORMS ON BOUNDED LATTICES	143
Oral	Fatma Karaca Vural, İ. Onur Kıymaz	INVESTIGATION OF THE EFFECT OF VACCINATION IN TURKEY ON PNEUMONIA CASES CAUSED BY COVID-19 WITH A SEIR TYPE MODEL	144
Oral	Abdussamet Çalışkan	NON-CANAL CIRCULAR SURFACES GENERATED BY SPECIAL CURVES WITH ALTERNATIVE FRAME AND QUATERNION	145
Online	Ömer Akgüller, Mehmet Ali Balcı	DISTRIBUTION OF DISCRETE GEODESICS ON ALPHA COMPLEXES	146
Online	Ömer Akgüller, Mehmet Ali Balcı, Sibel Paşalı Atmaca	SYMMETRIC DYNAMIC DERIVATIVES FOR QUAD REMESHING OF SIMPLICIAL SURFACES	147
Online	Halil Süel, Mahmut Tokmak, Akın Kıraç	CLASSIFICATION OF CAMERA TRAP DATA USING CONVOLUTIONAL NEURAL NETWORKS	148
Online	Mehmet Güvenç Negiz, Esra Özge Aygül, Tayfun İsa Aygül	TAXONOMIC DIVERSITY CALCULATION IN WOODY SPECIES: EXAMPLE OF DEDEGOL MOUNTAIN	149
Oral	Fernando S. Vidal Causanilles, Haci Mehmet Baskonus, Juan Luis Garcia Guirao, German Rodriguez Bermude	ON THE COMPLEX CHARACTERISTICS INTO THE LONG JOSEPHSON JUNCTION MODEL	150
Oral	Tuğçe Kunduracı, Ceren Sultan Elmalı, Tamer Uğur	TEXTURE SPACE ASSOCIATED WITH GRAPHS	151
Oral	Mehmet Korkmaz, Elif Özkurt Başustaoğlu	A STUDY ON THE GENERALIZATION OF SOME SIGMOIDAL GROWTH MODELS	152
Oral	Aziz Büyükkaragöz, Erdal Ünlüyol	GRAPH STRUCTURE OF A SPECIAL CONGRUENCE SUBGROUP OF THE EXTENDED MODULAR GROUP	153

Oral	Salisu Ibrahim, Tukur Abdulkadir Sulaiman, Abdullahi Yusuf, Ilhame Amirali, Usman Younas, Hasan Bulut	OPTICAL SOLITONS TO THE GENERALIZED HIGHER-ORDER NONLINEAR SCHRODINGER EQUATION ARISING IN OPTICAL FIBERS	154
Online	Özdemir Şentürk, Ahmet Mert	CALCULATING AND MAPPING OF TOPOGRAPHIC COMPLEXITY USING ENTROPY EQUATIONS	155
Online	Karmina K.Ali, Sibel Tarla, Reşat Yılmazer	DYNAMIC BEHAVIOR OF THE FOKAS SYSTEM ARISING IN MONOMODE OPTICAL FIBERS THROUGH THE UNIFIED AUXILIARY EQUATION METHOD	156
Online	Dilara Altan Koç, Hajar F.Ismael, Hasan Bulut	A CLASS OF SOLUTIONS FOR THE BOUSSINESQ-LIKE EQUATIONS INCLUDING SPATIO-TEMPORAL DISPERSION	157
Oral	Esma Kurban, Fatmanur Büyükbayraktar	OPINIONS OF PRE-SERVICE SCIENCE TEACHERS ABOUT VIRTUAL TOURS TO SCIENCE CENTERS	158
Oral	Nihan Güngör	ON SOLUTION OF THE NON-NEWTONIAN VOLTERRA INTEGRAL EQUATIONS WITH CONVOLUTION KERNEL	159
Online	Sibel Turanlı, Sedanur Uçan	A STUDY ON CONJUGATE CONNECTIONS ON PARA ANTI-HERMITIAN MANIFOLDS	160
Online	Beytullah Yağız, Yener Altun, Sinan Deniz	THE EXPLOSION OF THE SOLUTIONS OF STOCHASTIC WAVE EQUATIONS	161
Online	Mertcan Gülben, Emirhan Berberoğlu, Zeynep Öz, Murat Görgöz, Candan Aykurt, Kürşad Özkan, Özdemir Şentürk	ESTIMATING SAMPLE COMPLETENESS FOR BIODIVERSITY ASSESSMENT BASED ON TREE AND SHRUB SPECIES IN SARIKAYA WLDA/ANTALYA	162
Online	Yener Altun	LMI-BASED STABILITY ANALYSIS OF NONLINEAR SYSTEMS WITH TWO ADDITIVE TIME-VARYING DELAY COMPONENTS	163
Online	Ahmet Acarer, Halil İbrahim Ürkmez, Ali Şenol, Ahmet Mert	THE POTENTIAL DISTRIBUTION OF TEUCRIUM CHAMAEDRYS L. IN TURKEY	164
Online	Fatih Rıza Çelik, Gürhan İçöz	APPROXIMATION BY A NEW CONSTRUCTION OF PARAMETRIC BASKAKOV OPERATORS	165
Oral	Gökçe Çaylak Kayaturan	TO ALTERNATE THE ROUTES ENCODED BY BLOOM FILTERS IN COMPUTER NETWORKS REPRESENTED BY GRAPHS	166
Online	Oguzhan Karahan, Hasan Karcı, Ali Tangel	TIME OPTIMAL TRAJECTORY GENERATION IN JOINT SPACE FOR 6R INDUSTRIAL SERIAL ROBOTS USING CUCKOO SEARCH ALGORITHM	167
Online	Mesut Tartuk, Furkan Taha Nurdağ, Vedat Acar, Mehmet Fatih Akay, Fatih Abut	CALL ASSISTANT APPLICATION FOR CALL CENTERS	168

Online	Sinan Deniz, Şakir İşleyen, Beytullah Yağız	AN APPLICATION ON THE EXTENDED TECHNOLOGY ACCEPTANCE MODEL	169
Oral	Yıldıray Çelik	NEW OPERATION TYPES ON INTERVAL-VALUED NEUTROSOPHIC SOFT SETS	170
Oral	Yıldıray Çelik, Gül Ergün	A NEW VIEW ON INTUITIONISTIC FUZZY SOFT GRAPHS	171
Online	Şeyma Tülüce Demiray, Uğur Bayrakçı	A STUDY ON THE SOLUTIONS OF (1+1)-DIMENSIONAL MIKHAILOV NOVIKOV-WANG EQUATION	172
Online	Büşra Nur Er, Yavuz Altın	ON STATISTICAL CONVERGENCE OF GENERALIZED FUZZY NUMBER SEQUENCES	173
Oral	Güven Kaya	NEIMARK-SACKER BIFURCATION IN A PREY-PREDATOR MODEL WITH CONFORMABLE FRACTIONAL DERIVATIVE	174
Online	Gürhan İçöz, Seda Demir	APPROXIMATION PROPERTIES OF A NEW TYPE OF GAMMA OPERATOR	175
Online	Ali Farajzadeh, Alkan Özkan	ON METRIZABILITY OF TOPOLOGICAL GROUPS	176
Oral	Ömer Oruç	A COMBINED FINITE DIFFERENCE COLLOCATION METHOD FOR FITZHUGH NAGUMO EQUATION	177
Online	Fatma Bulut, Mehmet Bektaş	NONLINEAR DIFFERENTIAL EQUATIONS WITH ACCORDING TO THE DARBOUX FRAME	178
Online	Serkan Gülsoy, Candan Aykurt, Halil Süel, Ahmet Mert, Zeynep Öz, Ali Şenol	POTENTIAL DISTRIBUTION MODELING OF STYRAX OFFICINALIS L. IN MEDITERRANEAN ECOSYSTEMS	179
Oral	Güllü Esra Köse, Yusuf Uçar, Alaattin Esen	ON THE NUMERICAL SOLUTIONS OF TIME FRACTIONAL FITZHUGH NAGUMO EQUATION WITH FORCED TERM	180
Online	Tuba Dinç, Çiğdem Bektaş	THE KOTHE TEOPLITZ DUALS OF V0^λ (G)	181
Oral	Gökhan Yüce, Yıldıray Çelik	APPLICATIONS OF COMPLEX FUZZY SOFT SETS IN RING THEORY	182
Online	Hajar F.Ismael, Arif Özkul, Hasan Bulut	ON THE COMPLEX SOLUTIONS TO THE HIROTA–SATSUMA–ITO EQUATION	183
Online	Baransel Güneş, Hakki Duru	A SECOND-ORDER DIFFERENCE SCHEME ON BAKHVALOV MESH FOR THE SINGULARLY PERTURBED DELAY PSEUDO-PARABOLIC EQUATIONS	184
Online	Baransel Güneş, Musa Çakır	A ROBUST NUMERICAL METHOD FOR THE SINGULARLY PERTURBED SECOND-ORDER VOLTERRA-FREDHOLM INTEGRO-DIFFERENTIAL EQUATIONS	185
Online	Onur Zihni, Yıldıray Çelik	AN APPLICATION OF INTUITIONISTIC NEUTROSOPHIC SOFT GRAPHS IN A DECISIONMAKING PROBLEM	186
Oral	Ümmügülsüm Akbaba, Ali Hikmet Değer, Tolga Berber	AN ALGORITHM FOR THE VERTICES ON THE PATHS OF MINIMAL LENGTH IN THE SUBORBITAL GRAPHS	187

Oral	M. Ali Akbar, Md. Ekramul Islam, Fatma Berna Benli, Onur Alp İlhan	SOLITARY WAVE SOLUTIONS TO NLEES FOR HIGHER BALANCE NUMBER THROUGH THE MODIFIED SIMPLE EQUATION METHOD	188
Online	Faruk Düşünceli	NEW EXACT SOLUTIONS FOR THE IVANCEVIC OPTION PRICING MODEL	189
Online	İnci Çakmak, Aynur Şahin	ON THE PARTIAL (p , q) -DIFFERENTIAL EQUATIONS	190
Oral	Süleyman Şenyurt, Davut Canlı, Elif Çan	SOME SPECIAL SMARANDACHE RULED SURFACES ACCORDING TO FRENET FRAME IN E3 -II	191
Oral	Ümmügülsüm Akbaba, Ali Hikmet Değer	PELL AND PELL-LUCAS NUMBERS IN SUBORBITAL GRAPHS	192
Oral	Emine Özçelik, Çiğdem Bektaş	SOME RELATIONS BETWEEN THE SETS OF $\Delta_y^m(f)$ - STATISTICAL CONVERGENCE OF ORDER α	193
Oral	Gökçe İrem Kızılkaya, Elif Çil, Sevda Türkiş	DETERMINING THE SUSTAINABLE CONSUMPTION BEHAVIORS OF TEACHER CANDIDATES	194
Oral	Ayşe Kabataş	AN EIGENVALUE PROBLEM WITH A SPECTRAL PARAMETER IN BOUNDARY CONDITIONS	195
Online	Melike Karta	A NEW LOOK AT : THE NUMERİCAL SOLUTİON OF THE GENERALİZED BURGERS–HUXLEY EQUATİON	196
Online	Mehmet Açil	AN INVERSE RECONSTRUCTION PROBLEM	197
Online	Mehmet Bektaş, Esra Çiçek Çetin, Sümeyye Gür Mazlum	BIENERGIES FOR SYMPLECTIC REGULAR CURVE WITH SYMPLECTIC FRAME	198
Online	Esra Çiçek Çetin, Mehmet Bektaş	SOME NEW ASSOCIATED CURVES OF A NULL AND PSEUDO NULL CURVE IN 4-DIMENSIONAL MINKOWSKI SPACE-TIME	199
Online	Şeyda Özel, Mehmet Bektaş	BISECTOR CURVES OF COMFORMABLE CURVES IN R2	200
Online	Şeyda Özel, Mehmet Bektaş	ON THE CHARACTERİZATİONS OF CURVES WİTH MODIFIED ORTHOGONAL FRAME IN EUCLIDEAN 3-SPACE	201
Oral	Berat Karaağaç, Nuri Murat Yağmurlu, Alaattin Esen	NOVEL NUMERICAL SOLUTIONS OF TIME FRACTIONAL KORTEWEG-DE VRIES-BURGERS EQUATION	202
Oral	İrem Eroğlu	ON SOME FIXED POINT THEOREMS IN QUASIMETRIC SPACES AND ITS APPLICATIONS	203
Online	Ceren Ünal, Gamze Özel Kadılar,	AN APPLICATION ON EARTHQUAKE DATA UNDER NON-RESPONSE SCHEME IN SAMPLING THEORY	204
Online	Derya Turfan	A DIFFERENT APPROACH TO THE ASSESSMENT OF PARTICULATE MATTER (PM10) POLLUTION IN ANKARA	205
Online	Hatice Öncel Çekim, Gamze Özel Kadılar, Tuba Eroğlu Azak	COMPARISON OF AEGEAN REGION MAXIMUM EARTHQUAKE MAGNITUDES WITH TIME SERIES METHODS	206
Online	Meltem Sertbaş, Coşkun Sara	q-DIFFERENCE OPERATOR ON $L_Q^2(0,+\infty)$	207
Online	Gülnur Yel, Miraç Kayhan	NEW ANALYTICAL APPROACH TO THE GENERALIZED FISHER EQUATION	208
Online	Gülden Altay Suroğlu, Yusuf Güzel	ON MODIFIED WEINGARTEN PARALLEL RULED SURFACE	209

Oral	Ertuğrul Akgül, Elif Çil	OPINIONS OF TEACHER CANDIDATES ON WEB 2.0 TOOLS AND THEIR USAGE IN EDUCATION	210
Online	Ramazan Sarı, İnan Ünal	SEMI-INVARIANT SUBMANIFOLDS OF GENERALIZED KENMOTSU MANIFOLD ADMITTING A QUARTER SYMMETRIC NON METRIC CONNECTION	211
Online	Ramazan Sarı, İnan Ünal	SEMI-INVARIANT RIEMANNIAN SUBMERSIONS WITH RICCI SOLITON	212
Oral	Canan Ünlü	ON A HOMOGENEOUS-HETEROGENEOUS REACTIONS WITH NEWTONIAN HEATING IN FLOW OF A THIRD GRADE FLUID	213
Oral	Ali Öz, Oğuzhan Çelebi, Gökhan Kaplan, Barış Bayrak, Haluk Görkem Alcan, Abdulkadir Cüneyt Aydın	THE EFFECT OF POLYVINYL ALCOHOL FIBER ON THE MECHANICAL PROPERTIES OF HIGH STRENGTH GEOPOLYMER CONCRETES	214
Oral	Gökhan Kaplan, Oğuzhan Çelebi, Ali Öz, Haluk Görkem Alcan, Barış Bayrak, Abdulkadir Cüneyt Aydın	MECHANICAL PROPERTIES OF METAKAOLIN BASED GEOPOLYMERS USING WASTE GEOPOLYMER POWDER AND LIME	215
Oral	Şafak Nur Duman, Meral Cansız Aktaş	ANALYSIS OF PROBLEMS POSED BY PRESERVICE ELEMENTARY MATHEMATICS TEACHERS FOR THE ADDITION AND SUBTRACTION OF FRACTIONS	216
Online	Sena Halıcı, Ayşegül Çetinkaya	INTEGRAL REPRESENTATIONS OF k-SRIVASTAVA HYPERGEOMETRIC FUNCTIONS	217
Online	Sena Halıcı, Ayşegül Çetinkaya	RECURSION FORMULAS OF k-SRIVASTAVA HYPERGEOMETRIC FUNCTIONS	218
Oral	Hasan Gündüz, Ercan Çelik	COMPUTATION OF H_∞-NORM OF A TRANSFER MATRIX VIA BISECTION ALGORİTHM	219
Online	Hicran Gümüşbaş, Ulaş Vural	SEGMENTATION OF ELECTRICITY SUBSCRIBERS BASED ON THE CONSUMPTION PATTERNS	220
Oral	Noorolhuda Wyal, Erhan Coşkun	DSTABIL: A MATLAB PACKAGE FOR STABILITY ANALYSIS AND ANALYTICAL/NUMERICAL SOLUTION OF LINEAR DELAY DIFFERENTIAL EQUATION SYSTEMS	221
Online	Hakam Youness, Gaga Ahmed, Benachir El Hadadi	RENEWABLE ENERGY IN SMART GRID: PHOTOVOLTAIC POWER MONITORING SYSTEM BASED ON MACHINE LEARNING USING ANOPEN-SOURCE IOT PLATFORM	222
Online	Gülsen Kılıç, Mehmet Sezai Yıldırım	OVERVIEW OF SOME INTERVAL VALUED SEQUENCE SPACES	223
Online	Erhan Pişkin, Ayşe Fidan	BLOW UP OF SOLUTIONS FOR THE WAVE EQUATION VARIABLE COEFFICIENTS	224
Oral	Süleyman Şenyurt, Eda Öztürk	ADJOINT CURVES FROM THE MODIFIED FRAME OF INVOLUTE CURVE	225
Oral	Murat Candan	SEVERAL PROPERTIES OF COMPACT OPERATORS ON SOME GENERALIZED FIBONACCI DIFFERENCE SEQUENCE SPACES	226

Oral	Murat Candan	OPERATORS ON SOME VECTOR-VALUED ORLICZ DIFFERENCE SEQUENCE SPACES	227
Oral	Murat Candan	SOME RESULTS ON CONVERGENCE OF SOFT SEQUENCES IN SOFT TOPOLOGICAL SPACES	228
Oral	Yanlin Li, Süleyman Şenyurt, Ahmet Özduran, Davut Canlı	THE CHARACTERIZATIONS OF EQUIDISTANT RULED SURFACES GENERATED BY BINORMAL VECTORS	229
Oral	Tuğba Yazgan, Esin İlhan, Ercan Çelik, Hasan Bulut	THE SINE-GORDON EXPANSION METHOD AND THE MODIFIED EXPANSION FUNCTION METHOD FOR NEW HYPERBOLIC WAVE SOLUTIONS OF WU-ZHANG SYSTEM MODELS	230
Oral	Rıdvan Yaprak, Erhan Coşkun	A WOUND/SLOUGH MODEL ACCOUNTING FOR CELLULAR DIFFUSION AND DEBRIDEMENT	231
Online	Gözde Seven, Elif Çil	A STUDY ON THE STATUS OF RESEARCH ON SCİENCE TEACHERS' OPİNİONS ON STEM PRACTİCES	232
Online	Mahir Demir	SUSTAINABLE FISHERIES WITH OPTIMAL CONTROL THEORY	233
Online	Hajar Ahessab, Khalid Rahmani, Mohamed Al-Hattab	THE ELECTRONIC AND OPTICAL PROPERTIES OF GRAPHENE QUANTUM DOTS PYRENE FAMILY	234
Oral	Süleyman Dölek, Elif Çil, Sevda Türkiş	WEB 3.0 (SEMANTICS) APPLICATIONS IN EDUCATION	235
Online	Muhammed Çınar, Mikail Et, Koray İ. Atabey	SOME NEW SEQUENCE SPACES OF GENERALIZED FRACTIONAL ORDER	236
Online	Ayşe Metin Karakaş, Sinan Çalık, Aslıhan Demir	MARSHALL OLKIN GARIMA DISTRUBUTION AND STATISTICAL PROPERTIES	237
Oral	Hatice Karabenli, Alaattin Esen, N. Murat Yağmurlu	A NUMERICAL APPROACH FOR SAWADA-KOTERA EQUATION WITH TRIGONOMETRIC QUINTIC B-SPLINE COLLOCATION METHOD	238
Oral	Hatice Karabenli, Alaattin Esen, Yusuf Uçar	A COLLOCATION FINITE ELEMENT SOLUTION FOR THE FRACTIONAL FOKKER-PLANCK PROBLEM	239
Online	Kader Öğmen	THE RELATIONSHIP OF BILINGUALITY OF STUDENTS WITH THE ACHIEVEMENT OF SCIENCE COURSE	240
Oral	Sümeyye Gür Mazlum, Mehmet Bektaş	ON THE MODIFIED ORTHOGONAL FRAMES OF THE NON-UNIT SPEED CURVES IN EUCLIDEAN SPACE E3	241
Oral	Gülay Oğuz	SOFT TOPOLOGICAL HYPERVECTOR SPACES	242
Online	Derya Arslan	NEW NUMERICAL SCHEME FOR SOLVING OF SINGULARLY PERTURBED PROBLEM WITH NONLOCAL BOUNDARY CONDITIONS	243
Oral	Fatma Erdoğan, Sude Ay	CONTENT ANALYSIS OF THE THESES ABOUT MATHEMATICAL PATTERN IN TURKISH CONTEXT	244
Oral	Fatma Erdoğan, Neslihan Gül	A NEW TASK FOR MATHEMATICALLY GIFTED STUDENTS: CONSECUTIVE NUMBERS TRIANGLE	245
Online	Pelin Yaprakdal Uzun, Kıvanç Uzun, İlknur Koca	ANALYSIS OF THE PEER BULLYING IN SCHOOLS CONSIDERING BY FRACTIONAL ORDER METAPOPULATION MODELLING	246

Online	Megrini Meriem, Gaga Ahmed, Mehdaoui Youness	MATHEMATICAL MODELLING AND SPEED CONTROL OPTIMIZATION OF PMSM MACHINE FOR ELECTRIC VEHICLE APPLICATION	247
Online	Pelin Yaprakdal Uzun	ANALYSIS OF THE VECTORED PLANT DISEASE MODEL CONSIDERING BY FRACTIONAL OPERATOR	248
Oral	Olgun Durmaz, Büşra Aktaş, Halit Gündoğan	ON THE TOPOLOGIES IN DUAL SPACE AND THE ANALYTICITY OF DUAL FUNCTIONS	249
Oral	Serbay Duran, Himmet Korkmaz	EXAMINING SECONDARY MATHEMATICS TEACHERS' OPINIONS ABOUT THE EMERGED PROBLEMS DURING THE TEACHING OF MATHEMATICS	250
Online	Derya Avcı, Fatma Soytürk	FRACTIONAL OPTIMAL CONTROL OF AN EPIDEMIC MODELED WITH TWO TYPES OF NONLINEAR INCIDENCE RATES	251
Oral	Ayşe Kübra Demirel	FURTHER ON k-FRACTIONAL OPERATORS INVOLVING MITTAG-LEFFLER FUNCTIONS FOR STRONGLY (α,h-m)-CONVEX FUNCTIONS	252
Online	Yavuz Turan, Hayal Yavuz Mumcu	AMINATION OF 8TH GRADE STUDENTS' OPINIONS ON NEW GENERATION QUESTIONS	253
Online	Beyza Billur İskender Eroğlu, Dilara Yapışkan	AN OPTIMAL CONTROL STRATEGY FOR AN H1N1 MODEL	254
Online	Sevda Özdemir, Fevzi Erdoğan	STOCHASTIC DIFFERENTIAL EQUATION MODELING FOR GOLD DATA WITH YUIMAGUI	255
Online	Pakize Uygun, Süleyman Dırık, Mehmet Atçeken	P-SASAKIAN MANIFOLDS AND THEIR CURVATURE CLASSIFICATION	256
Online	Pakize Uygun	ON P-SASAKIAN MANIFOLDS SATISFYING CERTAIN CURVATURE CONDITIONS	257
Oral	Hüseyin Samancı, Serbay Duran	OMAR KHAYYAM'S PLACE AND IMPORTANCE IN THE HISTORY OF MATHEMATICS	258
Oral	Zeynep Kübra Kazan, Sevda Türkiş, Elif Çil	A DOCUMENT ANALYSIS ON THE EFFECT OF FINLAND AND TURKISH EDUCATION SYSTEMS ON PISA SUCCESS	259
Oral	Fatih Say	SOME ASYMPTOTIC FEATURES OF HYPER-AIRY EQUATION	260
Oral	Canan Çiftçi, Adel P. Kazemi	TOTAL DOMINATOR CHROMATIC NUMBER OF LEXICOGRAPHIC PRODUCT OF A CYCLE TO A GRAPH	261
Oral	Davut Canlı, Süleyman Şenyurt	BEZIER CURVES AND PATCHES WITH MODIFIED BERNSTEIN BASIS	262
Online	Swapnil Verma	REVISITING CHATTERJEA TYPE NONUNIQUE FIXED POINT THEOREMS VIA INTERPOLATION IN QPB - SPACE	263
Online	Zelal Temel, Musa Çakır	A NOVEL NUMERICAL APPROXIMATION FOR SINGULARLY PERTURBED VOLTERRA INTEGRO DIFFERENTIAL EQUATION WITH INTEGRAL BOUNDARY CONDITION	264

Online	Zelal Temel, Musa Çakır	CONVERGENCE ANALYSIS OF FITTED NUMERICAL METHOD FOR A SINGULARLY PERTURBED NONLINEAR VOLTERRA INTEGRODIFFERENTIAL EQUATION WITH INTEGRAL BOUNDARY CONDITION	265
Oral	Berat Gevşek, Elif Çil	EVALUATION OF TEACHER CANDIDATES' BIOTECHNOLOGY KNOWLEDGE LEVELS AND ATTITUDES TOWARDS BIOTECHNOLOGY	266
Oral	Rıdvan Şahin	A NEUTROSOPHIC CORRELATION COEFFICIENT AND ITS APPLICATION IN RECOGNITION OF PATTERNS	267
Oral	Rıdvan Şahin	NEW DISTANCE MEASURE FOR NEUTROSOPHIC SETS AND ITS APPLICATION IN MEDICAL DIAGNOSIS	268
Oral	Mustafa I.Hatim, Çiğdem A. Bektaş	A STUDY ON SOME SEQUENCE SETS BASED ON LUCAS BAND MATRIX AND MODULUS FUNCTIONS	269
Online	Elif Şenkal, Hayal Yavuz Mumcu	THE EFFECT OF MIND AND INTELLIGENCE GAMES COURSE ON STUDENTS' MATHEMATICS ACHIEVEMENT: AN EXPERIMENTAL STUDY	270
Oral	Mehmet Gıyas Sakar	A NEW REPRODUCING KERNEL APPROACH FOR NONLINEAR VARIABLE ORDER FRACTIONAL VOLTERRA INTEGRO-DIFFERENTIAL EQUATIONS	271
Oral	Büşra Uygun, Meral Cansız Aktaş	EXAMINATION OF 8th GRADE STUDENTS' OPINIONS ON MODEL ELICITING ACTIVITIES	272
Online	Ekhtiar Khodadadi, Mesut Karabacak, Ercan Çelik	NUMERICAL SOLUTION OF FUZZY DIFFERENTIAL EQUATIONS BY TWOSTEP MODIFIED SIMPSON RULE	273
Oral	Onur Saldır, Mehmet Gıyas Sakar	A NUMERICAL APPROACH FOR SINGULAR BOUNDARY VALUE PROBLEM	274
Oral	Onur Saldır	A NEW APPROACH FOR NUMERICAL SOLUTION OF FOURTH ORDER EMDEN-FOWLER EQUATION	275
Online	Fırat Çakır, Hayriye Güçkır Çakır, Musa Çakır	A FINITE DIFFERENCE METHOD FOR FREDHOLM INTEGRO-DIFFERENTIAL EQUATIONS	276
Oral	Hasen Mekki Öztürk	SPECTRA OF A LINEAR PENCIL OF PENTADIAGONAL MATRICES	277
Oral	Gizem Nur Battal, Meral Cansız Aktaş	EXAMINING OF MIDDLE SCHOOL 7TH GRADE STUDENTS' PROBLEM POSING STATUS ABOUT 'EQUALITY AND EQUATION'	278
Online	Ekhtiar Khodadadi, Mesut Karabacak, Ercan Çelik	PREDICTOR-CORRECTOR (m+1) −STEP METHOD FOR SOLVING FUZZY ORDINARY DIFFERENTIAL EQUATIONS BY USING THE INTERPOLATION OF FUZZY NUMBER	279
Online	Mohamed Hariri, Zohra Bouteffal, Nourelhouda Beghersa, Mehdi Benabdallah	ON THE STABILITY AND STABILIZATION FOR THE IMPLICIT DIFFERENTIAL CONTROL SYSTEM IN BANACH SPACES	280
Oral	Münevver Tuz	SOME NEW EXISTENCE RESULTS FOR FRACTIONAL DIFFERENTIAL INCLUSIONS WITH ANTI-PERIODIC BOUNDARY CONDITIONS	281
Oral	Münevver Tuz, Asıf Yokuş	INVESTIGATION OF SOLITARY WAVE SOLUTIONS FOR THE NONLINEAR (2 + 1) -DIMENSIONAL TYPICAL REFRACTION EQUATION	282

Online	Habiba Akter	EVOLUTIONARY COMPUTATION FOR MULTI-CONSTRAINED PATH SELECTION ALGORITHM ON THE INTERNE	283
Online	Yücel Baltürk, Hüseyin Demir, İnci Çilingir Süngü	ON NUMERICAL SOLUTION OF TWO POINT BOUNDARY VALUE PROBLEMS WITH LINEAR SHOOTING METHOD	284
Online	Sadaf Waziry, Jawad Rasheed, Mirsat Yesiltepe	PROPERLY WEARING FACE MASK DETECTION USING HYBRID DEEP AND CLASSICAL MACHINE LEARNING TECHNIQUES	285
Online	Ahmad Bilal Wardak, Jawad Rasheed, Mirsat Yesiltepe	AUTOMATIC NOISE DETECTION SYSTEM IN QR CODES	286
Oral	Zeki Taştan, Esra Alkan	THE MATHEMATICAL MIRACLE OF AN 80 MINUTE MEMORY	287
Oral	İsrafil Okumuş, Ercan Çelik	THE GROUP LAWS ON THE ALGEBRAIC CURVES	288
Online	Hülya Gültekin Çitil	ON A STURM-LIOUVILLE FUZZY PROBLEM	289
Oral	Mehmet Akif Akyol, Yılmaz Gündüzalp	SOME PROPERTIES ON POINTWISE SEMI-SLANT RIEMANNIAN MAPS (PSSRM)	290
Oral	Meltem Öğrenmiş	FRACTIONAL INVARIANTS OF CURVES IN LORENTZIAN PLANE	291
Oral	Meltem Öğrenmiş	SOME PROPERTIES OF CURVES WITH FRACTIONAL DERIVATIVE IN EUCLIDEAN SPACES	292
Oral	Zuhal Küçükarslan Yüzbaşı, Sevinç Taze	NOTES ON SURFACES WITH CONSTANT MEAN CURVATURE ALONG A GIVEN CURVE IN THE LIE GROUP	293
Oral	Bahadır Yüzbaşı	BIG DATA ANALYTICS VIA PENALIZED ESTIMATION	294
Oral	Berat Karaagac, Alaattin Esen, Ercan Çelik, Edson Pindza	NUMERICAL DYNAMICS OF VARIABLE ORDER FRACTIONAL BENNEY-LIN EQUATION VIA FINITE ELEMENT COLLOCATION METHOD	295
Oral	Elham Sefidgar, Ercan Çelik, Babak Shiri	COMPLEX TRANSFORM METHOD FOR SYSTEM OF FRACTIONAL DIFFERENTIAL EQUATIONS AND APPLICATIONS	296
Online	Hande Yüzel, Onur Toklu, Senem Şahan Vahaplar, M.Fatih Akay, Sevtap Erdem	BLOCKCHAIN-BASED SERVICE NETWORK PORTAL	297
Oral	Emine Funda Okumuş	RELATIVE TL BI-IDEAL, TL QUASI IDEAL, TL INTERIOR IDEAL	298
Oral	Bahar Açay	A MATHEMATICAL EXAMINATION OF THE FRACTIONAL HIV INFECTION MODEL	299
Oral	Zulqurnain Sabi, Haci Mehmet Baskonus, Fevzi Erdoğan	A NUMERICAL STOCHASTIC COMPUTING PARADIGMS FOR THE NONLINEAR SINGULAR DELAY DIFFERENTIAL PROBLEMS	300

Online	Saloua SENHAJI, Sanaa FAQUIR, Mohamed HAMLICH, Omar Diouri, Mohammed Ouazzani Jamil	ARTIFICIAL INTELLIGENCE FOR REAL-TIME DEFECT DETECTION AND INSPECTION OF PHOTOVOLTAIC PANELS	301
Online	Ajay Kumar, Haci Mehmet Baskonus	THE NONLINEAR FRACTIONAL GENERALISED POCHHAMMER-CHREE EQUATION ARISING AN ELECTRICAL FIELD THEORY IS DYNAMICAL ANALYSIS OF CAPUTO-FABRIZIO FRACTIONAL DERIVATIVE	302
Online	Mine Babaoğlu	EXACT SOLUTIONS OF GOURSAT PROBLEMS INVOLVING MIXED PARTIAL DERIVATIVES VIA HOMOTOPY ANALYSIS METHOD	303
Online	Nazlım Deniz Aral, Hacer Şengül Kandemir, Mikail Et	ON ρ− STATISTICAL CONVERGENCE OF DOUBLE SEQUENCES OF ORDER α IN TOPOLOGICAL GROUPS	304
Online	Oğuzhan Çelebi, Barış Bayrak, Mahmut Kılıç, Abdulkadir Cüneyt Aydın	INVESTIGATION OF THE EFFECTS OF AMBIENT VIBRATIONS FILTERED AT DIFFERENT FREQUENCY RANGE ON THE DYNAMIC BEHAVIOR OF THE BUILDING	305
Online	Oğuzhan Çelebi, Barış Bayrak, Mahmut Kılıç, Abdulkadir Cüneyt Aydın	ENERGY BASED EVALUATION OF REINFORCED CONCRETE STRUCTURE AFFORDABLE BY FAR FIELD EARTHQUAKE	306
Oral	Cemal Belen	TAUBERIAN THEOREMS FOR THE STATISTICALLY (N,P) SUMMABLE INTEGRALS	307
Online	Urdaletova A.B., Kydyraliev S. K.	DIRECT INTEGRATION OF THE SYSTEMS OF LINEAR DIFFERENTIAL EQUATIONS AND SOME APPLICATIONS IN ECONOMICS	308
Oral	Süleyman Şenyurt, Davut Canlı, Elif Çan	SOME SPECIAL SMARANDACHE RULED SURFACES ACCORDING TO FRENET FRAME IN E3 -III	309
Online	Oğuzhan Taş, Amani Yahyaoui	MACHINE LEARNING BASED INTRUSION DETECTION SYSTEM USING GRAY WOLF OPTIMIZATION FOR FEATURE SELECTION	310
Poster	Samet Uytun	EIGEN FUNCTION EXPANSIONS FOR A STRUM-LIOUVILLE OPERATOR FOR THE CASE OF AN INFINITE INTERVAL	311
Poster	Habiba Akter, Rupert Young	APPLICATION OF A GENETIC ALGORITHM FOR GENERATING REGULAR AND IRREGULAR NATURAL FRACTAL PATTERNS	312

On Solutions Stochastic Nonlinear Partial Differential Equations with Conformable Derivatives

Esma Ulutaş [1]

[1] Department of Electronics and Communication Engineering, Faculty of Technology, Karadeniz Technical University, Trabzon, Turkey

esmaates@ktu.edu.tr,eulutas1986@gmail.com

Abstract

It is well known that stochastic nonlinear partial differential equations(NPDEs) is more complicated then deterministic equations because of the additional random terms. In this study, it is considered a stochastic Wick-type nonlinear partial differential equation with conformable derivatives [1]. It is obtained exact and stochastic solutions of this equation using the extended G'/G method, white noise theory and Hermit transforms [2]. Then it is given a special example and shown how these solutions presented as Brownian motion function solutions.

Keywords: Extended G'/G-method; stochastic wick-type NPDEs; white noise theory.

REFERENCES

1. Wadati M., Stochastic Korteweg-de Vries equation, J. Phys. Soc. Jpn. 52, 2642-2648, 1983.
2. Holden H., Øksendal B., Ubøe J. and Zhang T., Stochastic partial differential equations: A Modeling, White Noise Functional Approach. Birkhäuser, Basel, 1996.

A BRIEF SURVEY ON NUMERICAL SOLUTIONS OF SINGULARLY PERTURBED PROBLEMS

Ilhame Amirali[1] and **Ece Eroğlu**[2]

[1]Department of Mathematics, University of Duzce, Duzce, Turkey

[2]Department of Mathematics, University of Duzce, Duzce, Turkey

[1]ailhame@gmail.com

[2] eceeroglu90@gmail.com

Abstract

In this work, a brief survey on numerical solutions of singularly perturbed problems is given. In the present survey contained in this paper, we don't aim at an entire complete bibliographical survey. A literature review was conducted between 2010-2021. The list of references in this paper, includes a variety of publications of singularly perturbed differential and integro-differential equations.

Keywords: Singular perturbation, Ordinary differential equations, Integro-differential equations, Initial and boundary value problems, Uniform convergence, Boundary layers, Error estimate.

REFERENCES

1. Amiraliyev, G. M. and Duru, H., A Note on A Parameterized Singular Perturbation Problem, Journal of Computational and Applied Mathematics, Vol:182, No:1, 233–242, 2005.
2. Bawa, R. K. and Natesan, S., A Computational Method for Self-Adjoint Singular Perturbation Problems Using Quintic Spline, International Journal of Computers and Mathematics with Applications, Vol:50, No:8, 1371- 1382, 2005.
3. Kumar, M. and Mishra, H. K., A Recent Survey on Computational Techniques for Solving Singularly Perturbed Boundary Value Problems, International Journal of Computer Mathematics, Vol:84, No:10, 1439 – 1463, 2007.
4. Kadalbajoo, M. K. and Gupta, V., A Brief Survey on Numerical Methods for Solving Singularly Perturbed Problems, Applied Mathematics and Computation, Vol:217, No:8, 3641–3716, 2010.
5. Kopteva, N. and O'Riordan, E., Shishkin Meshes in The Numerical Solution of Singularly Perturbed Differential Equations, International Journal of Numerical Analysis and Modeling, Vol:1, No:1, 1–18, 2009.
6. Mishra, H. K. and Saini, S., Various Numerical Methods for Singularly Perturbed Boundary Value Problems, American Journal of Applied Mathematics and Statistics, Vol:2, No:3, 129 – 142, 2014.

$(\Delta^m_v)_u$ - GENERALIZED WEIGHTED STATISTICAL CONVERGENCE

Rifat Çolak and Çiğdem Asma Bektaş [1]

[1] Department of Mathematics, University of Firat, Elazig, Turkiye

rcolak@firat.edu.tr and cbektas@firat.edu.tr

Abstract

In this study, we introduce and examine the concepts of $(\Delta^m_v)_u$ - generalized weighted statistical convergence and $(\Delta^m_v)_u$ - generalized weighted (N,t_{n})-summability. Furthermore we give some relations between $(\Delta^m_v)_u$ - generalized weighted statistical convergence and $(\Delta^m_v)_u$ - generalized weighted (N,t_n) - summability.

Keywords: Density, Cesàro summability, weighted statistical convergence, generalized difference sequence.

REFERENCES

1. M. Et and R. Çolak, On some generalized difference sequence spaces, Soochow J. Math. 21(4), 377-386 (1995).
2. M. Et and A. Esi, On Köthe-Toeplitz duals of generalized difference sequence spaces, Bull. Malaysian Math. Sc. Soc. 23, 25-32 (2000).
3. M. Et, H. Ş. Kandemir and H. Çakallı, Δ^m - weighted statistical convergence, AIP Conference Proceedings 2334, 040005 (2021).
4. S. Ghosal, Weighted statistical convergence of order α and its applications, J. Egyptian Math. Soc. 24(1), 60-67 (2016).
5. V. Karakaya and T. A. Chishti, Weighted statistical convergence, Iran. J. Sci. Technol. Trans. A Sci. 33(33), 219-223 (2009).
6. H. Kızmaz, On Certain Sequence Spaces, Canad. Math. Bull.24(2), 169-176 (1981).
7. M. Mursaleen, V. Karakaya, M. Ertürk and F. Gürsoy, Weighted statistical convergence and its application to Korovkin type approximation theorem, Appl. Math. Comput. 218(18), 9132-9137 (2012).

HYERS-ULAM STABILITY OF QUANTUM LOGIC FUZZY IMPLICATION

Iqbal H. Jebril and Najat M. Abed Abdelqader

Department of Mathematics, Faculty of Science and Technology,

Al-Zaytoonah University of Jordan, Amman Jordan

i.jebril@zuj.edu.jo

Abstract

In fuzzy logic we have four kinds of fuzzy implication which called (S, N)-Implications, R-Implications, QL-Implications, and D-Implication. Since there are many types of stability for various functional equations. In this research, we study Hyers-Ulam stability of functional equations with fuzzy implications for QL −implication.

Keywords: Fuzzy implication, QL −implication, Stability.

PρLog: A Language for Exact and Approximate Reasoning

Besik Dundua

Institute of Applied Mathematics, Tbilisi State University, Georgia
bdundua@gmail.com

Abstract

PρLog is a rule-based system [2, 3], that extends Prolog with strategy conditional transformation rules. These rules (basic strategies) define transformation steps on finite (possible empty) sequences. Strategy combinators help to combine strategies into more complex ones in a declaratively clear way. Transformations are nondeterministic and may yield several results, which fits very well into the logic programming paradigm. Strategic rewriting separates term traversal control from transformation rules. This allows the basic transformation steps to be defined concisely. The separation of strategies and rules makes rules reusable in different transformations. Transformation rules are equipped with four different kinds of variables (individual, sequence, function, and context variables) together with regular constraints. These variables allows to traverse sequences in single/arbitrary width (with individual and sequence variables) and terms in single/arbitrary depth (with functional and context variables). Regular constraints are useful to restrict possible values of sequence and context variables by regular sequence expressions and regular tree (context) expressions, respectively. These features facilitate flexibility in matching, providing a possibility to extract an arbitrary subsequence from a sequence, or to extract subterms at arbitrary depth.These capabilities enable PρLog to have highly declarative programming style that is expressive enough to support concise implementations for: specifying and prototyping deductive systems, solvers for various equational theories, tools for XML querying and transformation, etc. Recently, we extended PρLog with fuzzy proximity relation to support uncertain computation [1]. In this talk we give an overview of the PρLog system and underline its new applications.

Keywords: Fuzzy Logic Programming, Fuzzy Prolog, Proximity and Similarity Relations

Acknowledgement: This research has been supported the Shota Rustaveli National Science Foundation of Georgia under the grant YS-18-1480.

REFERENCES

[1] Besik Dundua. PρLog: a system for rule-based programming. Proceedings of the 9th International Symposium on Symbolic Computation in Software Science. RISC Technical Report 21-16. DOI 10.35011/risc.21-16. 2021

[2] Besik Dundua, Temur Kutsia, and Klaus Reisenberger-Hagmayr. An overview of PρLog. In: Y. Lierler and W. Taha, editors. Proceedings of the 19th International Symposium on Practical Aspects of Declarative Languages, PADL 2017. Volume 10137 of Lecture Notes in Computer Science. Springer, pages 34–49, 2017.

[3] Besik Dundua, Temur Kutsia, Klaus Reisenberger-Hagmayer. PρLog: Combining Logic Programming with Conditional Transformation Systems. Proceedings of the 32nd International Conference on Logic Programming, ICLP 2016. Vol. 52 of OpenAccess Series in Informatics (OASIcs). Schloss Dagstuhl, pages 10.1-10.5, 2016

SOME NEW SOLITON SOLUTIONS OF TIME FRACTIONAL RESONANT DAVEY-STEWARTSON EQUATIONS

Esin İlhan[1]

[1] Kırsehir Ahi Evran University, Kirsehir 40500, Turkey

eilhan@ahievran.edu.tr,

Abstract

In this study, via the Bernoulli sub-equation method (BS-EM), the traveling wave solution of the (2+1)-dimensional resonant Davey-Stewartson system is investigated. The nonlinear partial differential equation of the (2+1)-dimensional resonant Davey-Stewartson system is transformed into a nonlinear ordinary differential equation using a wave transformation, and then solved using the BS-EM approach. Some new solutions have been built successfully. The (2+1)-dimensional time-fractional resonant Davey-Stewartson equation is satisfied by all of the discovered solutions in this research. For all of the computations and graphic plottings in this investigation, we used Wolfram Mathematica 12 software.

Keywords: The Bernoulli sub-equation method (BS-EM), The Fractional Riemann-Liouville derivative, Resonant Davey-Stewartson equation.

REFERENCES

1. H Rezazadeh, D Kumar, A Neirameh, M Eslami, and M Mirzazadeh Pramana 94 1(2020).
2. T A Sulaiman, H Bulut, G Yel, and S S Atas Optical and Quantum Electronics (2018).
3. H F Ismael, W-X Ma, and H Bulut Physica Scripta 96 (2021).
4. H F Ismael, A Seadawy, and H Bulut International Journal of Nonlinear Sciences and Numerical Simulation (2021).
5. H F Ismael and H Bulut Mathematical Methods in the Applied Sciences (2021).
6. K K Ali, R Yilmazer, H M Baskonus, and H Bulut Physica Scripta (2020).
7. J Manafian, O A Ilhan, K K Ali, and S Abid EAST ASIAN JOURNAL ON APPLIED MATHEMATICS 10 549 (2020).

PERFORMANCES OF FITS WHEN SOME GROWTH MODELS ARE HYPERBOLIC

Mehmet Korkmaz[1]

[1] Department of Mathematics, Ordu University, Ordu, Turkey

mkorkmaz52@yahoo.com

Abstract

In this study, some growth models as well as their hyperbolic growth models were examined. In addition, the effect of these hyperbolic growth models on the selection of appropriate growth model was investigated by using some model selection criteria such as coefficient of determination and error sum of squares. Using data sets, the results of these hyperbolic growth models were found to be better than the results of these growth models.

Keywords: Hyperbolic; Growth Models; Monomolecular; Logistic; Gompertz

REFERENCES

1. Samuel Brody, Bioenergetics and Growth, Rheinhold Publishing, New York, 1945.
2. Harold E. Burkhart, Mike R. Strub, A model for simulation or planted loblolly pine stands. In growth models for tree and stand simulation. Edited by J. Fries. Royal College of Forestry, Stockholm, Sweden. Pp. 128-135. Model of forest growth. J. Ecol. 60: 849–873, 1974.
3. Anuraag R. Kansal, Salvatore Torquato, Griffith Rutherford Harsh, Ennino Antonio Chiocca, Thomas S. Deisboeck, Simulated brain tumor growth dynamics using a three dimensional cellular automaton. Journal of Theoretical Biology, Vol.203, No: 4, 367-382, 2000.
4. Cemil Kara, Ahmet Alp, Fatih Can, 2011. Growth and Reproductive Properties of Flathead Trout (Salmo platycephalus Bhenke, 1968) Population from Zamanti Stream, Seyhan River, Turkey. Turkish Journal of Fisheries and Aquatic Sciences, Vol.11, No:3, 367-375, 2011.
5. Samuel Oluwafemi Oyamakin, Angela Unna Chukwu, On the Hyperbolic Monomolecular Growth Model in Height/DiameterGrowth of PINES, British Journal of Mathematics and Computer Science, Vol:9, No:4, 320-327, 2015
6. William E. Ricker, Growth rates and models, Fish Physiology, Vol:8, 677-743, 1979.
7. Charles P. Winsor, 1932. The Gompertz curve as a growth curve, Proc. Natl. Acad. Sci., Vol:18, No:1, 1-8, 1932.
8. Abdulkadir Yıldızbakan, Analysis on mathematical models of tree growth and comparison of these models, MSc Thesis, Institute of Natural and Applied Sciences, University of Cukurova, Turkey, (in Turkish, with abstract in English), 2005.

The Consecutive Substitution Method for Boundary Value Problem with Retarded Argument

Arzu AYKUT[a,1] and Ercan ÇELİK[b]

[a] *Ataturk University, Faculty of Science, Department of Mathematics, Erzurum-TURKEY*

[b] *Kyrgyz-Turkish Manas University, Department of Applied Mathematics and Informatics, Bishkek-KYRGYZSTAN*

aaykut@atauni.edu.tr, ercan.celik@manas.edu.kg

Abstract

In this paper, we obtain the consecutive substitution method to problem for a differantial equation with retarded argument by this method, we obtain an integral equation that is equivalent to the boundary value problem. The equivalent integral equation is usually a Fredholm equation in the classicial theory. In this study we obtain a Fredholm- Volterra integral equation different from the classicial theory for the problem.

Keywords: Boundary value problem, Fredholm-Volterra integral equation,the consecutive substitution method.

REFERENCES

1. Aykut A., Yıldız B.,, On aboundary value problem for a differantial equation with variant retarded argument,, Applied Mathematics and Computation 93, 63-71, 1998.
2. Aykut A., Çelik E., Bayram M., The modified two sided approximations method and Padé approximants for solving the differantial equation with variant retarded argument, Applied Mathematics and Computation, 144, 475-482.,2003.
3. Çelik E., Aykut A., Bayram M., The modified successive approximations method and Padé approximants for solving the differantial equation with variant retarded argument, Applied Mathematics and Computation, 151, 393-400, 2004.
4. Çelik E., Aykut A., Bayram M.., The ordinary successive approximations method and Padé approximants for solving the differantial equation with variant retarded argument, Applied Mathematics and Computation, 144, 173-180., 2003.

[1] Corresponding author

MATHEMATICS TEACHERS' VIEWS ABOUT THE TRANSITION FROM ONLINE TEACHING TO FACE-TO-FACE EDUCATION

Himmet Korkmaz [1] **, Kevser Kurucu Bayhan** [3]

[1, 3] Department of Mathematics and Science Education, Ordu University, Ordu, Turkey

himmetkorkmaz@odu.edu.tr, kevserkrcbyhn@gmail.com

Mahmut Kertil [2] **,**

[2] Department of Mathematics and Science Education, Marmara University, İstanbul, Turkey

mkertil@marmara.edu.tr

Abstract

This study aims to examine the problems experienced by mathematics teachers in the transition of students from distance and online education to face-to-face education. With the return to face-to-face education, some of the challenges, problems, and what can be done to overcome them are worth investigating. There are very few studies comparing online and face-to-face as teaching approaches for mathematics teaching. In a study, it is found that there exists no significant difference between online and face-to-face math teaching in terms of the impact on the academic achievements of sixth-graders [1]. The current research was carried out by case study method from qualitative research methods and five mathematics teachers with different levels and experience in a province in the Black Sea region of Turkey were selected as participants of the study with a purposeful sampling method. Semi-structured interviews were conducted with these teachers. The data obtained from the interview were examined and the codes from the related literature and the ones resulting from the research formed the categories. The findings demonstrated that students have some behavioral and educational adaptation problems. Teachers have expressed difficulties in complying with classroom rules, listening to lessons, and communicating as behavioral adaptation problems of students. In terms of education, the lack of subject matter knowledge from previous semesters, slowing down habits such as writing and taking notes, and decreased interest and attitudes towards mathematics courses were expressed. The teachers who participated in the study stated that they observed negative attitudes in students' behavior, such as apathy, prejudice development, timidity, and lack of self-confidence. In addition, the opinions and recommendations of teachers have been examined to address or improve these problems.

Keywords: Online Teaching; Face-to-face education; Mathematics teachers.

REFERENCES

1. Edwards, Clayton, and Rule, Audrey. Attitudes of middle school students: Learning online compared to face to face. Journal of Computers in Mathematics and Science Teaching, Vol: 32, No:1, 49-66, 2013.

SOLUTION OF (2+1)-DİMENSİONAL QUANTUM ZAKHAROV-KUZNETSOV EQUATİONS USİNG THE THE MODIFIED exp(−Ω(ξ))-expansion FUNCTION METHOD

S.Şule Şener Kılıç[1*], Ercan Çelik and Hasan Bulut*****

*Department of Mathematics, Ataturk University , Erzurum, Turkey

***Kyrgyz-Turkish Manas University, Department of Applied Mathematics and Informatics, Bishkek-Kyrgyzstan*

***Department of Mathematics, Fırat University , Elazığ, Turkey

senersule@atauni.edu.tr, ercan.celik@manas.edu.kg, hbulut@firat.edu.tr

Abstract

In this paper, we obtain the (2+1)-dimensional quantum Zakharov-Kuznetsov equations are by using the the modified exp(−Ω(ξ))-expansion function approach. All the investigated solutions in this study have supported their corresponding model. Under the choice of suitable values of the parameters involved, the 2D and 3D graphs of the reported solutions are plotted.

Keywords: Expansion method, conformable derivative, Zakharov-Kuznetsov equation

REFERENCES

1. Miller K.S., and Ross B., An Introduction to the Fractional Calculus and Fractional Differential Equations, New York, Wiley, 1993.
2. Song,L.Zhang, H. Solving the fractional BBM-Burgers equation using the homotopy analysis method, Chaos Solitons Fractals, V. 40, pp. 1616-1622, 2009.
3. Zhang,X. Liu,J. Wen, J. Tang, B. He,Y. Analysis for one-dimensional time-fractional Tricomi-type equations by LDG methods, Numerical Algorithms, V.63, N.1, pp. 143-164., 2013.
4. Baleanu, D. Machado, J.A.T. Cattani, C. Baleanu, M.C. Yang, X.J. Local fractional variational iteration and decomposition methods for wave equation on Cantor sets within local fractional operators, Abstract and Applied Analysis, V. 2014, 12 pages, 2014.

[1]Corresponding author

ABOUT NEW MATHEMATICAL APPROACHES IN FRACTIONAL BIOLOGICAL MODELLING

Ozlem Defterli [1]

[1] Department of Mathematics, Cankaya University, 06790 Ankara, TURKEY
defterli@cankaya.edu.tr

Abstract

The subject of fractional calculus, as the calculus of integral and derivatives of arbitrary order, has gained popularity and importance, mainly due to its demonstrated applications in diverse fields of science and engineering (see [1-4] and references therein). This is due to the reason that these operators enable the description of the memory effect which naturally is involved in various complex processes. Hence, there is a growing need to study and use the fractional-order differential and integral equations for effective modelling of such complex phenomena.

In this study, some recent mathematical tools of fractional calculus will be presented for the mathematical modelling and anticipation of important complex biological phenomena, such as infectious diseases. The performances of new fractional operators are investigated on the considered biological models. Numerical simulations are performed with respect to the different values of the biological model parameters and order of the fractional operators.

Keywords: Complex systems; Fractional calculus; Numerical approximations.

REFERENCES

1. Diethelm K, A fractional calculus based model for the simulation of an outbreak of dengue fever, Nonlinear Dynamics, Vol: 71, No: 4, 613-619, 2013.
2. Atangana A, Baleanu, D, New fractional derivatives with non-local and non-singular kernel: theory and application to heat transfer model, Thermal Science, Vol: 20, No: 2, 757-763, 2016.
3. Defterli Ozlem, Modeling the impact of temperature on fractional order dengue model with vertical transmission, International Journal of Optimization and Control: Theories and Applications, Vol: 10, No: 1, 85-93, 2020.
4. Defterli Ozlem, Comparative analysis of fractional order dengue model with temperature effect via singular and non-singular operators, Chaos, Solitons & Fractals, Vol: 144, Article No:110654, 2021.

FIXED POINTS AND COUPLED FIXED POINTS IN B-METRIC SPACES VIA GRAPHICAL CONTRACTIONS

LILIANA GURAN[1], MONICA-FELICIA BOTA[2], GABRIELA PETRUSEL[3]
[1]Department of Pharmaceutical Sciences, Vasile Goldis Western University of Arad, Romania.
guran.liliana@uvvg.ro
[2]Department f Mathematics, Babes-Bolyai University of Cluj-Napoca, Romania.
monica.bota@ubbcluj.ro
[3]Department of Business, Babes-Bolyai University Cluj-Napoca, Romania.
gabriela.petrusel@ubbcluj.ro

Abstract. In this paper some existence and stability results for cyclic graphical contractions in complete metric spaces are given. Moreover, we discuss a vectorial case of our main result. Some applications to coupled fixed point problems are also derived.

Keywords: Fixed point, b-metric space, vector-valued b-metric space, coupled fixed point.

REFERENCES

[1] I.A. Bakhtin, The contraction mapping principle in quasimetric spaces, Funct. Anal., Unianowsk Gos. Ped. Inst. 30(1989) 26-37.
[2] V. Berinde, Generalized contractions in quasimetric spaces, Seminar on Fixed Point Theory, 1993, 3-9.
[3] V. Berinde, Generalized coupled fixed point theorems for mixed monotone mappings in partially ordered metric spaces, Nonlinear Anal. 74 (2011) 7347-7355.
[4] M. Bota, A. Molnar, Cs. Varga, On Ekeland's variational principle in b-metric spaces, Fixed Point Theory, 12(2011), No. 1, 21–28.
[5] S. Czerwik, Contraction mappings in b-metric spaces, Acta Mathematica et Informatica Univ. Ostraviensis 1(1993) 5-11.

SIMULATION OF A LIGHT-EMITTING DIODE BASED ON ORGANIC MATERIALS UNDER ATLAS-SILVACO

Anass El karkri[1], **Imane El mhamdi**[1], **Zakaria El malki**[1], **Mohamed bouachrine** [2], **Jean-Marc Sotiropoulos** [3]

[1] High School of Technology, Moulay Ismail University, Meknes, Morocco

[2] High School of Technology, Sultan Moulay Slimane University, Khenifra, Morocco

[3] Pau and Adour Countries University, UMR5254 - IPREM Chemistry-Physics Team Helioparc Pau, France

anass.elkarkri@gmail.com,

Abstract

Organic light-emitting diodes (OLEDs) represent light sources capable of providing very high luminance. OLEDs have attracted a lot of attention because of their desirable characteristics including their low consumption of electrical energy, their flexibility... Regarding the structure, OLEDs are composed of an organic semiconductor layer sandwiched between two electrodes (at least one transparent) allowing the application of a DC voltage necessary for their functioning. The objective of our work consists on the one hand in determining the electronic and optical properties of an OLED based on the organic material having the structure P3-B-EDOT-B-T3-A [molecule composed of Phenylene (P), Thiophene (T), 3,4-Ethylenedioxythiophene (EDOT) and Benzothiadiazole (B)] using the simulation software TCAD-SILVACO "ATLAS", and on the other hand, it shows the effect of varying the thickness of the emitting layer and the doping concentration of charge carriers on these properties.

Keywords: Organic materiel, OLED, donor, DFT, I-V caracteristic.

NOVEL SOLUTION FOR DETERMINISTIC TYPHOID FEVER MODEL WITHIN THE FRAME OF FRACTIONAL ORDER

Esin İlhan[1]

[1] Kırsehir Ahi Evran University, Kirsehir 40500, Turkey

eilhan@ahievran.edu.tr,

Abstract

The main aim of the present investigation is to analyse mathematical model exemplifying typhoid fever illustrated in system of five ordinary differential equations using the fractional natural decomposition method (FNDM) with the Caputo fractional operator. The projected algorithm is amalgamation of Adomian decomposition technique and natural transform. For diverse fractional order, the behaviour of the outcomes is presented in terms of plots. The present study exemplifies the importance of fractional operator and efficiency of the projected algorithm. Finally, the present study may help to examine the wild class of real-world models and also aid to predict their behaviour with respect to parameters considered in the models.

Keywords: Typhoid fever; Caputo derivative; fractional natural decomposition method.

REFERENCES

1. Liouville, J., Memoire surquelques questions de geometrieet de mecanique, etsur un nouveau genre de calcul pour resoudreces questions, J. Ecole. Polytech., 13, 1- 69, (1832).
2. Riemann,G.F.B.,Versuch Einer AllgemeinenAuffassung der Integration und Differentiation, Gesammelte Mathematische Werke, Leipzig, (1896).
3. Caputo, M., Elasticita e Dissipazione, Zanichelli, Bologna, (1969).
4. Miller,K.S. and Ross,B., An introduction to fractional calculus and fractional differential equations, A Wiley, New York, (1993).
5. Podlubny, I.,Fractional Differential Equations, Academic Press, New York, (1999).
6. Kilbas,A.A., Srivastava, H.M. and Trujillo, J.J.,Theory and applications of fractional differential equations, Elsevier, Amsterdam, (2006).

The periodic wave solutions of conformable time-fractional mathematical model by analytical method

Aslı Alkan[2],Tolga Aktürk[1], Hasan Bulut[2]
[1]Department of Mathematics and Science Education, Faculty of Education, Ordu University, Turkey
[2]Department of Mathematics, Faculty of Science, Fırat University, Turkey

[1]tolgaakturkk@gmail.com, [2]alkanasli47@gmail.com; [2]hbulut@firat.edu.tr,

Abstract

In this paper, the wave solutions of the time-fractional Zoomeron equation were get by the modified exponential function method (MEFM). When the solution functions found as a result of the calculations are analyzed, hyperbolic and trigonometric functions with periodic functions are also rational functions. Two-three-dimensional contour and density graphs representing the characteristic feature were drawn by determining the values according with the method in the parameters in the solution functions obtained.

Keywords: Conformable time-fractional Zoomeron equation; modified exponential function method; traveling wave solutions.

References

[1] R. Bertram, The development of fractional calculus 1695-1900. Hist. Math. 4(1) (1977) 75-89.

[2] I. Podlubny, Fractional Differential Equations, Academic Press, London, 1999.

[3] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, Theory and Applications of 120 Fractional Differential Equations, North-HollandMathematical Studies, Elsevier, Amsterdam, 2006.

[4] K. Mille, B. Ross, An Introduction to the Fractional Calculus and Fractional Differential Equations, Wiley, New York, 1993.

[5] R.L. Magin, Fractional Calculus in Bioengineering, BegellHouse Publisher, 125 West Redding, Conn, 2006.

[6] R. Khalil, M.A. Horani, A.M.Y. Sababheh, A new definition of fractional derivative, J. Comput. Appl. Math. 264 (2014) 65-70.

[7] T. Abdeljawad, On conformable fractional calculus, J. Comput. Appl. Math. 279 (2015) 57-66.

[8] K. Kamruzzaman, M.A. Akbar, Exact solutions of the (2+1)-dimensional cubic Klein-Gordon equation and the (3+1)-dimensional Zakharov-Kuznetsov equation using the modified simple equation method. J. Assoc. Arab Univ. Basic Appl. 15 (2014) 74-81.

[9] Odabasi, M., Traveling wave solutions of conformable time-fractional Zakharov–Kuznetsov and Zoomeron equations. Chinese Journal of Physics, 64 (2020) 194-202.

ANALYSING ONE PREY AND TWO PREDATORS MATHEMATICAL MODEL WITH HOLLING TYPE II RESPONSE AND FADING MEMORY

Zeynep Yılmaz*[1], Selahattin Maden[1], Aytül Gökçe[1]

[1]Department of Mathematics, Ordu University , Ordu, Turkey

zeynepyilmaz@odu.edu.tr*,
smaden@odu.edu.tr,
aytulgokce@odu.edu.tr,

Abstract

Studying prey-predator interactions forms the basis of ecology as it helps to understand the dynamics of population models. In this presentation, we introduce one prey and two predators mathematical model with Holling type 2 response function. The aim of this study is to investigate this model by taking the role fading memory effect and competition into account on the dynamics. The fixed points and local stability analysis of the four-component model are investigated and the findings are supported with numerical simulations.

Keywords: Population interactions; Fading memory effect; Stability analysis.

REFERENCES

1. Aytül Gökçe, The influence of past in a population system involving intraspecific competition and Allee effect , The European Physical Journal Plus, (2022) 137:200.

2. Banshidhar Sahoo, Swarup Poria**,** Dynamics of predator–prey system with fading memory, Applied Mathematics and Computation, 347 (2019) 319-333.

STRESS ANALYSIS OF MULTI-LAYERED SOIL MEDIUM BY USING FINITE AND INFINITE ELEMENTS

Yusuf Ziya Yüksel , Şeref Doğuşcan Akbaş2,

[1,2]Department of Civil Engineering, University of Bursa Technical, Bursa, Turkey

yusuf.yuksel@btu.edu.tr, seref.akbas@btu.edu.tr,

Abstract

In this study, stress analysis of in two dimensional elastic, isotropic a semi-infinite layered soil medium is by using finite element method. The soil is modelled using finite-infinite elements. In the literature, more realistic results are obtained when using infinite elements for infinite media solutions compared to finite element solutions. For solving the problem, cubic finite elements with 16 nodes and infinite elements with 8 nodes are generated. The shape functions of the infinite elements are calculated separately for each direction. The region close to the soil surface is used with finite elements, and the region far from the soil surface is used with infinite elements. Finite-infinite elements are solved by Gauss Legendre integration method with five nodes. The numerical results obtained with the Boussinesq theory are compared and verified. A parametric study is performed using different layer thicknesses and different material properties in a semi-infinite soil medium. The numerical results and graphs obtained with MATLAB. The obtained results are compared.

Keywords: Stress analysis; Finite – infinite element method; Multi-layered soil.

REFERENCES

1. Timoshenko Stephen, Goodier, J. N., Theory of Elasticity, 1951.
2. Curnier, Alain, A Static Infinite Element, International Journal for Numerical Methods in Engineering, Vol:19, No:10, 1479-1488, 1983.
3. Bettess, Peter, and Jacqueline A. Bettess, Infinite Elements for Static Problems, Engineering Computations, 1984.
4. Kumar, Prabhat, Static Infinite Element Formulation, Journal of Structural Engineering, Vol:111, No:11, 2355-2372, 1985.
5. El-Esnawy, N. A., A. Y. Akl, and A. S. Bazaraa. "A new parametric infinite domain element." Finite elements in analysis and design 19.1-2 (1995): 103-114.

A NEW SPECIES RICHNESS MEASURE IMPROVED FROM MARGALEF AND MENHINICK

Kürşad Özkan[1], Serkan Özdemir[2], Ali Şenol[1], Ecir Uğur Küçüksille[3]

[1] Isparta University of Applied Sciences, Faculty of Forestry, 32260, Isparta, Turkey

[1] Isparta University of Applied Sciences, Sutculer Prof. Dr. Hasan Gürbüz Vocational School, 32950, Isparta, Turkey

[1] Suleyman Demirel University, Faculty of Engineering, 32260, Isparta, Turkey

kursadozkan@isparta.edu.tr,

Abstract

In the present study, a new estimator $\left({}^{\alpha^*}D_{MM}\right)$ is proposed to measure species richness. Its' essential inputs are alfa (α) scale parameters of Margalef (D_{MG}) and Menhinick (D_{MN}) indices.

To evaluate the performance of ${}^{\alpha^*}D_{MM}$, a hypothetical community dataset was used. The computations were performed using a spreadsheet program created for ${}^{\alpha^*}D_{MM}$. According to estimation results, ${}^{\alpha^*}D_{MM}$ seems to be a better form rather than D_{MG} and D_{MN}. Therefore it may be employed for comparing species richness of the natural communities. To better understand the performance of ${}^{\alpha^*}D_{MM}$, further studies should be however generated using various types of real ecological data.

Keywords: Alpha diversity, assemblage, biodiversity, diversity estimators, species richness

REFERENCES

1. Margalef, S. R. (1969). Diversity and stability: a practical proposal: a model of interdependence. Book haven Symposia in Biology 22:25-37.
2. Menhinick, E.F. (1964) A comparison of some species individuals diversity indices applied to samples of field insects. Ecology 45. 859–61.
3. Özkan, K. 2016b. Biyolojik Çeşitlilik Bileşenleri (α. β ve γ) Nasıl Ölçülür (1. Basım). Süleyman Demirel Üniversitesi. Orman Fakültesi Yayını. Isparta.

WIGNER-VILLE DISTRIBUTION AND AMBUGUITY FUNCTION ASSOCIATED WITH QUATERNION OFFSET LINEAR CANONICAL

M. YOUNUS BHAT [1]

[1]Department of Mathematical Sciences, Islamic University of Science and Technology, Awantipora, India

gyounusg@gmail.com,

Abstract

This work is devoted to the development of the Wigner-Ville transform (WVD) associated with quaternion offset linear canonical transform (QOLCT) theory proposed by Bhat and Dar in 2021 that has been designated as an emerging tool in the scenario of sig- nal processing as a detection of the linear frequency modulated signal. The purpose of this work is to introduce Wigner-Ville distribution(WVD) and Ambiguity function(AF) associ- ated with the quaternion offset linear canonical transform(WVD-QOLCT/AF-QOLCT). Firstly, we propose the definition of the WVD-QOLCT and then several important prop- erties such as dialation, nonlinearity, boundedness are derived. Secondly, we focus on the ambiguity function for the proposed transform. A bunch of important properties lincluding reconstruction formula associated with the ambiguity function are studied.

.

Keywords: Quaternion algebra; Quaternion offset linear canonical transform; Wigner- Ville distribution; Ambugity function; Dalation

.

Solitary wave solutions to a nonlinear mathematical model

Hasan Bulut[1], Sümeyye Kılbitmez[1], Tukur A Sulaiman[2,3], Abdullahi Yusuf[2,3]

[1]Department of Mathematics, Firat University, Elazig, Turkey

[2]Department of Computer Engineering, Biruni University, Istanbul, Turkey

[3]Department of Mathematics, Federal University Dutse, Jigawa, Nigeria

Abstract

The solitary wave solutions to a nonlinear mathematical model arising in mathematical physics are established in this study. To construct such novel solutions, we employ robust integration technique called modified Sardar method. The novel soliton control structure has been studied in order to explain specific physical problems. Three-dimensional and contour graphs have been plotted using suitable choice of the values of parameters involved.

Keywords: Nonlinear equation; Sardar method; Solitary waves.

[1] John, F.: Partial Differential Equations. Applied Mathematical Sciences, 4th edn. Springer, Berlin (1991)

[2] Protter, M.H., Weinberger, H.F.: Maximum Principles in Differential Equations, Corrected Reprint of the1967 Original. Springer, New York (1984)

[3] Schatzman, M.: Numerical Analysis—A Mathematical Introduction. Oxford University Press, Oxford (2002)

[4] Smith, G.D.: Numerical Solutions of Partial Differential Equations, Finite Difference Methods. Oxford Uni-versity Press, New York (1985)

THE WIND LOAD CALCULATION OF A MASONRY MINARET BY USING DIN1056 STANDARD

Erdem TÜRKELİ [1]

[1] Vocational School of Technical Sciences, Construction Department, Ordu University, Ordu, Turkey

erdemturkeli@odu.edu.tr

Abstract

Masonry minarets are the most special and specific structures for Islamic Culture that should be preserved to the next generations as they can carry the memories and the traces of the past generations. Even this is the case, there are so many outer factors that adversely affects the structural load carrying masonry system of these slender structures. The most dominant outer factors that adversely affect the load carrying system of masonry minarets are earthquakes and wind storms. Therefore, the effect of wind loads on these specific structures should be carefully investigated and determined. In this study, the wind loads for a masonry minaret is calculated by using DIN1056 standard and the details of procedure is provided explicitly.

Keywords: Wind load; Masonry; Minaret; DIN1056 standard

On the complex soliton solutions to the Hirota–Satsuma–Ito equation

Hajar F. Ismael[1,2], **Arif Özkul**[1], **Hasan Bulut**[1]

Hasan Bulut[1]

[1]Department of Mathematics, University of Firat, Elazig, Turkey

[1]Department of Mathematics, University of Zakho, Zakho, Iraq

hajar.ismael@uoz.edu.krd,

Abstract

In this work, we study the Hirota–Satsuma–Ito equation that describes the propagation of unidirectional shallow-water waves and interactions of two long waves with different dispersion forms. For this study, we use the sine-Gordon expansion method for the suggested equation. Complex soliton solutions in terms of dark-bright, dark and bright are constructed. Obtained solutions are constructed in three-dimensional figures to understand these physical phenomena.

Keywords: Sine-Gordon expansion method; Hirota–Satsuma–Ito equation; Complex solutions.

REFERENCES

1. Zhou, Y., Wang, C., & Zhang, X. (2020). Rational Localized Waves and Their Absorb-Emit Interactions in the (2+ 1)-Dimensional Hirota–Satsuma–Ito Equation. Mathematics, 8(10), 1807.
2. Long, F., Alsallami, S. A., Rezaei, S., Nonlaopon, K., & Khalil, E. M. (2022). New interaction solutions to the (2+ 1)-dimensional Hirota–Satsuma–Ito equation. Results in Physics, 105475.

KANTOROVICH TYPE (p,q)-GENERALIZATION OF BERNSTEIN OPERATORS

Hayatem Hamal and Pembe Sabancıgil

Department of Mathematics, Tripoli University, Libya

hafraj@yahoo.com

Department of Mathematics, Eastern Mediterranean University, Gazimagusa, North Cyprus

pembe.sabancigil@emu.edu.tr

Abstract

In this paper, we introduce a new generalization of Kantorovich type Bernstein operators by means of (p,q)-calculus for $0<q<p\le 1$ and we derive a recurrence formula for these newly defined operators to give explicit formulas for the *m*-th order moments and central moments which play an important role in approximation theory.

Keywords: (p,q)-calculus ; Bernstein polynomials; Kantorovich type operators.

Definition 1. Let $0<q<p\le 1$. Kantorovich type (p,q)-analogue of Bernstein operators is defined as follows:

$$B^{*}_{n,p,q}(f,x)=\sum_{k=0}^{n} b_{n,k}(p,q,x)\int_{0}^{1} f\left(\frac{p^{n-k}\left([k]_{p,q}+q^{k}t\right)}{[n+1]_{p,q}}\right)d_{p,q},\quad x\in[0,1],\ n\in\mathbb{N},$$

where $b^{p,q}_{n,k}(x)=\dfrac{1}{p^{n(n-1)/2}}\begin{bmatrix} n \\ k \end{bmatrix}_{p,q} p^{k(k-1)/2}x^{k}(1-x)_{p,q}^{\,n-k}.$

Lemma 1. For all $n\in\mathbb{N}$, $x\in[0,1]$, $m\in\mathbb{Z}\cup\{0\}$ and $0<q<p\le 1$ we have

$$B^{*}_{n,p,q}(t^{m},x)=\sum_{j=0}^{m}\binom{m}{j}\frac{[n]_{p,q}^{n}}{[n+1]_{p,q}^{m}}\frac{p^{nm}}{[m-j+1]_{p,q}}\sum_{i=0}^{m-j}\frac{1}{p^{n(i+j)}}\binom{m-j}{i}\left(q^{n}-p^{n}\right)^{i}B_{n,p,q}\left(t^{i+j},x\right),$$

where $B_{n,p,q}(f,x)=\sum_{k=0}^{n} b^{p,q}_{n,k}(x)\,f\left(p^{n-k}\dfrac{[k]_{p,q}}{[n]_{p,q}}\right).$

REFERENCES

1. Nazim Mahmudov, Pembe Sabancigil, Approximation Theorems for q-Bernstein-Kantorovich Operators, Filomat, vol: 27, No:4, 721-730, 2013.

On the solution of Pavlov equation by The Variational Iteration Method

[1] Derya Deniz and Mücahit Saydam[1]

[1]Department of Mathematics, Firat University, Elazig, Turkey

deryadeniz485@yandex.com,mucahitsydm.23@gmail.com

Abstract:

In this work, the variational iteration method (VIM) that is a semi-analytical method is applied the $(2+1)$-dimensional Pavlov equation. The numerical results show that only a few terms are sufficient to obtain accurate solutions.

Keywords: The variational Iteration method, $(2+1)$-dimensional Pavlov equation

References:

1. DUNAJSKI, Maciej. A class of Einstein–Weyl spaces associated to an integrable system of hydrodynamic type. *Journal of Geometry and Physics*, 2004, 51.1: 126-137.

2. BULUT, Hasan; BASKONUS, H. Mehmet. A comparsion among homotopy perturbation method and the decomposition method with the variational iteration method for dispersive equation. *International Journal of Basic and Applied Sciences*, 2009, 9.10: 32-42.

3. BULUT, Hasan. Comparing numerical methods for Boussinesq equation model problem. *Numerical Methods for Partial Differential Equations*, 2009, 25.4: 783-796.

4. BULUT, Hasan; BASKONUS, H. Mehmet. A comparsion among homotopy perturbation method and the decomposition method with the variational iteration method for dispersive equation. *International Journal of Basic and Applied Sciences*, 2009, 9.10: 32-42.

5. INC, Mustafa; CAVLAK, Ebru. On numerical solutions of a new coupled MKdV system by using the Adomian decomposition method and He's variational iteration method. *Physica Scripta*, 2008, 78.4: 045008.

6. ALHARBI, Abdulghani; ALMATRAFI, M. B. Numerical investigation of the dispersive long wave equation using an adaptive moving mesh method and its stability. *Results in Physics*, 2020, 16: 102870.

THE COMPARISON OF NUMERICAL SOLUTIONS OF COMFORMABLE AND β − COMFORMABLE FRACTIONAL DIFFERENTIAL EQUATIONS USING NEURAL NETWORK METHOD BY PYTHON

Sadullah Bulut[1], Muhammed Yiğider[2]

[1] Department of Mathematics, University of Erzurum Technical, Erzurum, Turkey

[2]Department of Mathematics, University of Erzurum Technical, Erzurum, Turkey

sadullah.bulut@erzurum.edu.tr

myigider@erzurum.edu.tr

Abstract

ANNs are mathematical models inspired by biological nerve systems. The neural network method for solving differential equations has recently achieved significant advances in solving fractional differential equations. The Space-Time Inhomogeneous Heat Equation is solved using a neural network method in this research. The loss function comprising initial/boundary conditions is produced using changeable parameters (weights and deviations). A Space-Time Inhomogeneous Heat Equation is also expressed as an optimization problem in this article. To demonstrate the accuracy of the method utilized, numerical examples with proven analytical solutions are described, including numerical results, graphs, weights, and biases. The graphical and tabular results are also thoroughly examined. The average squared errors for various neuron, learning rate, and training steps values are shown.

Keywords: Neural Network Method, Numerical Solutions, Space-time Fractional Differential Equations

REFERENCES

1. Lee, H. Kang, I.S. 1990. Neural algorithms for solving differential equations. Journal of Computational Physics, 91(1): 110-131.
2. Lee, H. and Kang, I., Neural algorithms for solving differential equations, Journal of Computational Physics, vol. 91, pp. 110-117, 1990.
3. Meade Jr, A.J. and Fernadez, A.A., The numerical solution of Linear Ordinary Differential Equations by Feedforward Neural networks, Math. Comput. Modelling, vol. 19, no. 12, pp. 1-25, 1994.
4. Meade Jr, A.J. and Fernadez, A.A., Solution of Nonlinear Ordinary Differential Equations by Feedforward Neural networks, Math. Comput. Modelling, vol. 20, no. 9, pp. 19-44, 1994.

Modeling Potential Distribution of Capparis spinosa in Babadağ Region

Alican ÇIVĞA[1], **Serkan ÖZDEMİR**[1], **Serkan GÜLSOY**[1]

[1] Isparta University of Aplied Science, Isparta, Turkey

alicancivga@isparta.edu.tr

Abstract

Capparis spinosa is a medical plant that has both economic (food, animal breeding, medicine, etc.) and ecological (erosion control, fighting wildfires, etc.) importance and is distributed in the western and southern coastal regions of Turkey. The Maxent model was used to simulate potential distribution areas of spinosa under environmental conditions. The results showed that the potential suitable area of spinosa is 6109 hectares, mainly distributed in below 1000 meters in Babadağ Region. It was determined that the variables contributing to the model were bedrock, elevation, topographic position index and hillshade index, respectively. The acquired model presented excellent performance according to its AUC values (Training AUC: 0.909 and test AUC: 0.906). It is thought that the results revealed in the study will be important for the planning studies to be carried out for the species.

Keywords: Caper, Habitat suitability, MaxEnt, Species distribution modeling.

References

Araújo, M. B., Pearson, R. G., Thuiller, W., & Erhard, M. (2005). Validation of species–climate impact models under climate change. Global change biology, 11(9), 1504-1513.

Gassó, N., Thuiller, W., Pino, J., Vilà, M., 2012. Potential distribution range of invasive plant species in Spain. NeoBiota, 12, 25.

Kart, A. (2019). Bioactive, pharmacological effects of caper plant and its use in neurodegenerative diseases. Veterinary Journal of Mehmet Akif Ersoy University, 4(2), 101-107.

Özçelik, H., Koca, A. (2011, September). Caper (Capparis L./Capparaceae) Genus and Economic Importance in Turkey. In 2nd International Non-Wood Products Symposium (p. 32).

Yeğenoğlu, S., Ayşe, U. Z. (2011). Caper: Its importance from historical and medical perspective. Lokman Hekim Journal, 35-36.

ARDUINO BASED DIGITAL TACHOMETER WITH TRACKING PROGRAM

Furkan ŞİMŞEK[1], Şükrü KİTİŞ[1]

[1] Simav Technology Faculty, Kütahya Dumlupınar University, Kütahya, Turkey

sukru.kitis@dpu.edu.tr

Abstract

This project; It is made to be used in situations where speed measurement is required in asynchronous motors and to control the data. Arduino Nano R3 and IR Transceiver Sensor, LCD screen and various cables were used in the project.

Especially in asynchronous motors used in industry, synchronization is broken due to some unavoidable situations. The problems experienced in these engines, which are generally used in production, cause problems in operation and then cause serious problems. With this project, instantaneous measurement of engine revolutions will be possible, the received data will be sent to the tracking program created with C#, and faulty times will be monitored in the form of graphics and tables. The sent data is transmitted to the c# software with delays of 0.5 seconds in the first 10 minutes, and 1-3 seconds for 30 minutes and above. Although the deviation of the obtained result is not based on the exact measurements, it is about 3-5%.

The tachometer realized with this project costs 20-30 times cheaper than the ones offered for sale in the market and becomes more advantageous thanks to the tracking program.

Keywords: Tachometer, Arduino, C#, Asynchronous motors

AN APPROXIMATION METHOD FOR THE EIGENVALUE OF A STURM-LIOUVILLE PROBLEM

Elif BAŞKAYA [1]

[1] Department of Mathematics, Karadeniz Technical University, Trabzon, Turkey

elifbekar@ktu.edu.tr

Abstract

In this paper, we approximate the eigenvalues of a Sturm-Liouville problem includes the spectral parameter in a boundary condition by associating the problem with Riccati equation. The potential of the problem is real valued, continuous, monoton and symmetric.

Keywords: Sturm-Liouville problem; Asymptotic eigenvalues; Continuous potential.

REFERENCES

1. Başkaya Elif, Periodic and Semi-Periodic Eigenvalues of Hill' s Equation with Symmetric Double Well Potential, TWMS Journal of Applied and Engineering Mathematics, Vol: 10, No:2, 346-352, 2020.
2. Başkaya Elif, Asymptotics of Eigenvalues for Regular Sturm-Liouville Problems with Spectral Parameter-Dependent Boundary Conditions and Symmetric Single Well Potential, Turkish Journal of Mathematics and Computer Science, Vol: 13, No:1, 44-50, 2021.
3. Coşkun Haskız, Başkaya Elif, Asymptotics of Eigenvalues for Sturm-Liouville Problem with Eigenvalue in the Boundary Condition for Differentiable Potential, Annals of Pure and Applied Mathematics, Vol:16, No:1, 7–19, 2018.

ON A GENERALIZATION OF MERSENNE HYBRID NUMBERS

Engin Özkan [1] **, Mine Uysal** [2]

[1]Department of Mathematics, Erzincan Binali Yıldırım University,

Faculty of Arts and Sciences, 24100 Erzincan, Turkey

eozkan@erzincan.edu.tr

[2]Graduate School of Natural and Applied Sciences,

Erzincan Binali Yıldırım University, 24100 Erzincan, Turkey.

Corresponding author: mine.uysal@erzincan.edu.tr

Abstract

We give a generalization of Mersenne hybrid numbers. We find the Binet formula, the generating function, the sum, the character, the norm and the vector representation of the generalizition. We obtain some relations among this generalizition and well known hybrid numbers. Then we present some important identities, Cassini, Catalan, Vajda, D'ocagne, Honsberger for the generalizition.

Keywords: Mersenne hybrid numbers; Binet formula; Generating function; Cassini identity.

REFERENCES

1. Songül Çelik, İnan Durukan, Engin Özkan, New recurrences on Pell numbers, Pell-Lucas numbers, Jacobsthal numbers, and Jacobsthal-Lucas numbers, Chaos, Solitons & Fractals, 150, 111173, 2021.
2. Engin Özkan, Ali Aydoğdu, İpek Altun, Some Identities for A Family of Fibonacci and Lucas Numbers, Journal of Mathematics and Statistical Science, Vol:3, No:10, 295-303, 2017.
3. Engin Özkan, Merve Taştan, On Gauss Fibonacci polynomials, on Gauss Lucas polynomials and their applications, Communications in Algebra, Vol:48, No: 3, 952-960, 2020.
4. Mustafa Özdemir, Introduction to hybrid numbers, Advances in Applied Clifford Algebras, Vol:28, No:1, 1-32, 2018.
5. Paula Catarino, Helena Campos, Paulo Vasco, On the Mersenne sequence, CM-Centro de Matemática, Vol:46, 27-53, 2016.
6. Nabiha Sabaa, Ali Boussayouda, Kasi Viswanadh V. Kanurib, Mersenne Lucas numbers and complete homogeneous symmetric functions, Journal of mathematics and computer science, Vol: 24, No:2, 127-139, 2021.

COMPARATIVE ANALYSIS OF MACHINE LEARNING ALGORITHMS FOR EARLY DIAGNOSIS OF ALZHEIMER'S DISEASE

Hanife Göker[1] **, Afrah Said**[2]**,**

[1] Department of Electrical Electronics Engineering, Kutahya, Turkey

[2] Department of Electrical Electronics Engineering, Kutahya, Turkey

hanife.goker@dpu.edu.tr, afrah.said@ogr.dpu.edu.tr

Abstract

Alzheimer disease is a disorder that affects the neurochemical systems in the brain and it is characterized by deteriorative changes in the neurons that causes dementia or even memory loss in some patients. The disease usually affects the elderly people aged 65 and above, but it sometimes affects younger people due to the gene that is present in their family history. Therefore, early diagnosis of the disease is very important for the treatment of the disease. In this study machine learning algorithms are used to predict the disease using Alzheimer's dataset. The dataset contains clinical patient data available as open source from Kaggle. The clinical data for the Alzheimer's disease dataset has 373 records, 206 controls and 167 patients, with a set of key indicators of the health of the elderly. The dataset is divided into 70% training and 30% testing dataset. The performances of Naive Bayes, Support Vector Machine, Random Forest, k-Nearest Neighbor and Multilayer Perceptron machine learning algorithms were compared according to model performance criteria using Alzheimer's dataset. Among the algorithms whose performances were compared, the Random Forest Algorithm showed the highest performance. The performance analysis results were found to be 95.59% sensitivity, 91.38% specificity, 90.91% precision, 91.74% f1 score, 83.93% MCC and 91.96% accuracy.

Keywords: Alzheimer’s disease; Machine learning; Early diagnosis; Classification; Predictive diagnostics.

On the Difference Between Numerical and Spectral Radii for Some Analytic Functions of Operator

Elif Otkun Çevik [1]

[1] Department of Computer Engineering, University of Avrasya, Trabzon, Turkey

elifotkuncevik@gmail.com,

Abstract

In this work, some estimates for the difference number between numerical and spectral radii for polynomial, exponential, sine and cosine functions of linear bounded Hilbert space operators via same difference numbers of powers corresponding Hilbert space operators have been obtained.

Keywords: Spectral radius; Numerical radius.

REFERENCES

1. K. E. Gustafson, D. K. M. Rao, Numerical range: The field of values of linear operators and matrices, Springer, New York, 1997.
2. F. Kittaneh, Spectral radius inequalities for Hilbert space operators, Proceedings of the American Mathematical Society, Vol:134, No:2, 385–390, 2006.
3. H. Guelfen, F. Kittaneh, On numerical radius inequalities for operator matrices, Numerical Functional Analysis and Optimization, Vol:40, No:11, 1231–1241, 2019.
4. W. Bani-Domi, F. Kittaneh, Norm and numerical radius inequalities for Hilbert space operators, Linear and Multilinear Algebra, Vol:69, No:5, 934–945, 2021.
5. A. Abu-Omar, F. Kittaneh, Estimates for the numerical radius and spectral radius of the Frobenius companion matrix and bounds for the zeros of polynomials, Annals of Functional Analysis, Vol:5, No:1, 56–62, 2014.
6. A. Abu-Omar, F. Kittaneh, Upper and lower bounds for the numerical radius with an application to involution operators, Rocky Mountain Journal of Mathematics, Vol:45, No:4, 1055–1064, 2015.
7. A. Abu-Omar, F. Kittaneh, Notes on the spectral radius and numerical radius inequalities, Studia Mathematica, Vol:227, No:2, 97–109, 2015.

*Abstract Submission should be prepared only **1 page**.

STATIC RESPONSES OF A CAVITY IN A HALF-SPACE BY USING FINITE ELEMENT METHOD

Yusuf Ziya Yüksel[1], **Şeref Doğuşcan Akbaş**[2],

[1,2]Department of Civil Engineering, University of Bursa Technical, Bursa, Turkey

yusuf.yuksel@btu.edu.tr, seref.akbas@btu.edu.tr,

Abstract

In this work, static displacement analysis of two dimensional, elastic, isotropic semi-infinite soil space with a cavity are investigated. The analysis is performed by using the finite element method. The region close to the ground is modelled with finite elements, and the region far from the ground is modelled with infinite elements extending to infinity. In the literature, infinite elements used instead of finite elements for infinite media solutions. Infinite element shape functions are generated according to the direction. By using one-dimensional two-node infinite elements and 8-node infinite elements are produced. A finite element with 16 nodes is used to solve the problem. The numerical results compared with the Boussinesq theoretical results. Numerical solutions of finite-infinite elements are obtained by Gauss Legendre integration method. A single load is given from the ground surface. Effects of cavity geometry, point location and material and geometry parameters of soil on the static displacements are investigated. By using finite and infinite model. The finite-infinite element model is coded with MATLAB. Static results are compared.

Keywords: Finite – infinite element; Soil medium; Cavity, Half-Space

REFERENCES

1. Timoshenko Stephen, Goodier, J. N., Theory of Elasticity, 1951.
2. Curnier, Alain, A Static Infinite Element, International Journal for Numerical Methods in Engineering, Vol:19, No:10, 1479-1488, 1983.
3. Bettess, Peter, and Jacqueline A. Bettess, Infinite Elements for Static Problems, Engineering Computations, 1984.
4. Kumar, Prabhat, Static Infinite Element Formulation, Journal of Structural Engineering, Vol:111, No:11, 2355-2372, 1985.
5. El-Esnawy, N. A., A. Y. Akl, and A. S. Bazaraa. "A new parametric infinite domain element." Finite elements in analysis and design 19.1-2 (1995): 103-114.

A Fractional Model to Investigate Shape Effects on Thermal Behavior of Brinkman-type Ferrofluid under Thermal Radiation and Heat Injection/Consumption

Asifa[1], **Poom Kumam**[1,*]

[1] Department of Mathematics, Faculty of Science, King Mongkut's University of Technology Thonburi (KMUTT), Bangkok 10140, Thailand

asifaasifa1992@gmail.com, poom.kum@kmutt.ac.th

Abstract

The prime aim of this article is the construction of a fractional model to enhance the heat transfer rate for magnetohydrodynamic (MHD) convectional transport of ferrofluid by applying the time-fractional concept of Caputo-Fabrizio derivative on Brinkman type fluid model. Kerosene oil and water are considered to serve as carrier fluids for iron oxide (Fe_3O_4) nanoparticles of various non-spherical shapes (brick, cylinder, platelet, and blade). The non-uniform time-dependent motion of an unbounded upward plate causes to generate the flow of ferrofluid. The supplementary physical phenomena such as ramped heating, thermal radiation, and heat injection/consumption are also analyzed. In addition, the effectiveness of using differently shaped Fe_3O_4 nanoparticles for heat transfer and ferrofluid flow is evaluated. The exact solutions for non-integer order modeled differential equations are computed by employing the Laplace transform method. The comparative graphical illustrations for ramped and constant (isothermal) velocity and temperature conditions are prepared to scrutinize the impacts of several associated thermal and physical quantities.

Keywords: Brinkman-type ferrofluid; Fractional model; Shape effect

REFERENCES

1. M. Caputo, M. Fabrizio. A new definition of fractional derivative without singular kernel. *Progr. Fract. Differ. Appl.*, 1(2):1{13, 2015
2. R.L. Hamilton, O.K. Crosser. Thermal conductivity of heterogeneous two-component systems. *Ind. Eng. Chem. Fundam.*, 1(3):187{191, 1962.

Determination of Indicator Plant Species by Indicator Analysis in Red Pine (*Pinus brutia* Ten.) and Black Pine (*Pinus nigra* J.F.Arnold) Forests in Bozdağlar Region

Ayşegül Tekeş[1]

[1]Department of Forest Engineering, Faculty of Forestry, Isparta University of Applied Sciences, Isparta, Turkey

aysegultekes@hotmail.com,

Abstract

This study was performed to identify the indicator plant species of Red Pine (*Pinus brutia* Ten.) and Black Pine (*Pinus nigra* J.F.Arnold), which are among our target species distributed in Bozdağlar (İzmir-Manisa) Region. Data from sixty-seven sample fields were put to use in the study. The vegetation data matrix was organized as the presence or absence of species. Then, this vegetation data matrix was interpreted by the indicator analysis in the PC-ORD program, and eight positive and five negative indicator species for red pine and six positive indicator species for larch were defined. Among these species, it has been inferred that hairy rock-rose (*Cistus creticus* L.) and sage-leaved rock-rose (*Cistus salviifolius* L.) were the prominent positive indicators for red pine, while dog roses (*Rosa canina* L.) were the most notable negative indicator plant species in the region. For the black pine, it has been concluded *Geocaryum macrocarpum* (Boiss.&Spruner) Engstrand species is the strongest positive indicator plant species in the region.

Keywords:. Bozdağlar, Indicator Plant Species, Red Pine, Black Pine, Indicator Analysis

REFERENCES

1. Godefroid S, Koedam N, Identifying Indicator Plant Species of Habitat Quality and Invasibility as a Guide for Peri-Urban Forest Management, Biodiversity and Conversation, Vol:12, 1699-1713, 2003.
2. Gülsoy S, Şentürk Ö, Negiz M G, Identification of Indicatory Plant Species for Target Species by Using Quantitative Methods: A Case Study From Acıpayam District, SDU Faculty of Forestry Journal, Vol:14, 10-14, 2013.
3. Güner Ş T, Özkan K, Çömez A, Çelik N, Woody Indicator Species of Probable Productive Potantial Areas of Anatolian Black Pine (*Pinus nigra* subsp. *pallasiana*) in the Inner Anatolia Region, Ecology, Vol:20, No: 80, 51-58, 2011.
4. Kremen C, Assessing the Indicator Properties of Species Assemblages for Natural Areas Monitoring, Ecological Applications, Vol:2, No:2, 203-217, 1992.

Solution of singular boundary value problems using artificial neural networks

Zulqurnain Sabir[1,2]

[1]Department of Mathematical Science, College of Science, United Arab Emirates University, Al Ain, Abu Dhabi, UAE

Email: zulqurnain_s@uaeu.ac.ae

[2]Department of Mathematics and Statistics, Hazara University, Mansehra, Pakistan

[a]Email: zulqurnain_maths@hu.edu.pk

Abstract

In this study, an intelligence computational scheme is presented for solving linear and nonlinear singular models using the well-known artificial neural networks (ANNs), genetic algorithm is a global search scheme, interior-programming algorithm is a competent local search scheme and the hybrid of global and local search. The neural network provides convenient approaches to obtain valuable prototypes based on an unsubstantiated error for singular models. The incentive for awarding this research work originates a consistent structure combines with the influential geographies of ANNs to handle the challenges of the singular models. Broad numerical research is accomplished to indorse the convergence, robustness and accuracy of the suggested numerical scheme. The numerical outcomes are also compared with the true results to examine the perfection of the planned numerical structure.

Keywords: Singular systems, nonlinear, genetic algorithm, interior programming algorithm, hybrid approach

References

[1] Sabir, Z., Raja, M.A.Z., Umar, M. and Shoaib, M., 2020. Neuro-swarm intelligent computing to solve the second-order singular functional differential model. The European Physical Journal Plus, 135(6), pp.1-19.

[2] Sabir, Z., Raja, M.A.Z., Umar, M. and Shoaib, M., 2020. Design of neuro-swarming-based heuristics to solve the third-order nonlinear multi-singular Emden–Fowler equation. The European Physical Journal Plus, 135(5), p.410.

[3] Sabir, Z., Manzar, M.A., Raja, M.A.Z., Sheraz, M. and Wazwaz, A.M., 2018. Neuro-heuristics for nonlinear singular Thomas-Fermi systems. Applied Soft Computing, 65, pp.152-169.

[4] Sabir, Z., Raja, M.A.Z., Shoaib, M. and Aguilar, J.F., 2020. FMNEICS: Fractional Meyer neuro-evolution-based intelligent computing solver for doubly singular multi-fractional order Lane–Emden system. Computational and Applied Mathematics, 39(4), pp.1-18.

A NOVEL ANALYTICAL APPROACH TO THE BENJAMIN–ONO EQUATION

Gülnur Yel[1] , Beyhan Kemaloglu[2] , Hasan Bulut[2]

[1]Faculty of Education, Final International University, Kyrenia, Mersin 10, Turkey
[2]Department of Mathematics, Firat University, Elazig, Turkey

gulnur.yel@final.edu.tr, beyhanozturk1980@gmail.com, hbulut@firat.edu.tr

Abstract

In this research, we present some new wave solutions to the second-order Benjamin-Ono equation by using an analytical scheme which is the sine-Gordon expansion method. The Benjamin-Ono equation is similar to the KdV equation and it describes internal waves in stratified fluids. We achieved some travelling wave solutions in terms of hyperbolic functions. All the solutions obtained have been analyzed graphically to their physical properties. We concluded that the proposed method is an efficient method that gives analytical solutions to powerfully nonlinear partial differential models.

Keywords: The sine-Gordon expansion method; Second-order Benjamin-Ono Equation; dark-bright solitary waves

REFERENCES

1. Benjamin, T.B., Internal waves of permanent form in fluids of great depth, Journal of Fluid Mechanics, 29(3), 559–592, 1967.
2. Ewen, J.F., Gunshor, R.L. and Weston, V.H., Surface acoustic solitons, Japanese Journal of Applied Physics, 19(S1), 683, 1980.
3. N. Taghizadeh, M. Mirzazadeh and F. Farahrooz, Exact Soliton Solutions for Second-Order Benjamin-Ono Equation, Applications and Applied Mathematics: An International Journal, 6 (1), 2011.
4. Yang Y., Zhu Y, Darboux-Bäcklund transformation, breather and rogue wave solutions for Ablowitz-Ladik equation, Optik, 217, Article 164920, 2020.

A novel technique to construct exact solutions for the Complex Ginzburg-Landau equation using quadratic-cubic nonlinearity

Muhammad Abubakar Isah[1] , Asıf Yokuş [2]
[1,2]Department of Mathematics, University of Firat, Elazig, Turkey

Myphysics_09@hotmail.com [1], asfyokus@yahoo.com [2]

Abstract

The Complex Ginzburg-Landau equation (CGLE) has been suggested and investigated as a model for turbulent dynamics in nonlinear partial differential equations. It is a particularly intriguing model in this aspect since it is a dissipative form of the nonlinear Schrödinger equation. The model is studied with quadratic-cubic law nonlinear fiber by using φ6-model expansion method, bright, dark, dark-singular, singular and dark-bright optical soliton solutions are obtained. The findings of this study might assist in comprehending some of the physical implications of various nonlinear physics models. The hyperbolic sine, for example, appears in the calculation of the Roche limit and gravitational potential of a cylinder, while The hyperbolic cosine function is the shape of a hanging cable (the so-called catenary). Finally, there are some discussions regarding new complex solutions. It is explored by giving physical meaning to the constants found in traveling wave solutions, which are both physically and mathematically significant. Three-dimensional simulations and contour plots are used to enhance these discussions.

Keywords: φ6-model expansion approach, Complex Ginzburg-Landau equation with quadratic cubic nonlinearity; Jacobi elliptic functions, solitons.

REFERENCES

1. Yokus, Asıf, et al. "Role of Gilson-Pickering equation for the different types of soliton solutions: a nonlinear analysis." The European Physical Journal Plus 135.8 (2020): 1-19.
2. Sajid, Naila, and Ghazala Akram. "Novel solutions of Biswas-Arshed equation by newly φ6-model expansion method." Optik 211 (2020): 164564.
3. Arshed, Saima, et al. "Optical solitons with complex Ginzburg-Landau equation having three nonlinear forms." Physics Letters A 383.36 (2019): 126026.
4. Ali, Khaled K., Abdul-Majid Wazwaz, and M. S. Osman. "Optical soliton solutions to the generalized nonautonomous nonlinear Schrödinger equations in optical fibers via the sine-Gordon expansion method." Optik 208 (2020): 164132.

A stochastic computing heuristic scheme for solving a novel third order perturbed delay differential Emden-Fowler model

Zulqurnain Sabir[1], Ayse Nur Akkilic[2], Hasan Bulut[2]

[1]Department of Mathematics, Hazara University, Mansehra Pakistan

[2]Department of Mathemetics, Firat University, Turkey
Email: zulqurnain_maths@hu.edu.pk, akkilicaysenur@gmail.com, hbulut@firat.edu.tr

Abstract: This study shows the design of novel third order perturbed delay differential Emden-Fowler model. The comprehensive structures based on the singular-point, pantograph and perturbed terms are given along with the shape factor of the third order perturbed delay differential Emden-Fowler model. The novel singular model is simulated by using the artificial neural networks (ANNs) along with the optimization based global/local schemes, i.e., genetic algorithm (GA) and interior-point algorithm (IPA). The constructed of an activation function is provided by using the differential system based on the third order perturbed delay differential Emden-Fowler model. The fitness function's optimization is provided through the hybrid of the ANNs-GA-IPA to solve the third order perturbed delay differential Emden-Fowler model. The exactness, validation, and verification of the system is observed by taking three different cases of the novel mathematical system. The stability, robustness, convergence, and correctness of the numerical scheme is performed through the comparison of the obtainable exact results. For the consistency, the statistical outputs with necessary procedures are obtained by using the ANNs-GA-IPA.

Keywords: Perturbed; Delay; Singular; Artificial neural networks; Local scheme; Global approach; Neurons.

References

[1] P. A. Farrell, A.F. Hegarty, J.J.H. Miller, E.O'Riordan and G.I.Shishkin, (2000), Robust Computational Techniques for Boundary Layers, Chapman-Hall/CRC, New York.

[2] Doolan, E.P., Miller, J.J. and Schilders, W.H., 1980. Uniform numerical methods for problems with initial and boundary layers. Boole Press.

[3] H. G. Roos, M. Stynes, L. Tobiska, (1996), Numerical Methods for Singularly Perturbed Differential Equations, ConvectionDiffusion and Flow Problems, Springer Verlag, Berlin.

[4] T. Linss, M. Stynes, (1999), A hybrid difference scheme on a Shishkin mesh for linear convection-diffusion problems, Appl. Numer. Math. 31 255-270.

[5] Erdogan, F., Sakar, M.G. and Saldır, O., 2020. A finite difference method on layer-adapted mesh for singularly perturbed delay differential equations. Applied Mathematics and Nonlinear Sciences, 5(1), pp.425-436.

Analysis of the Effect of the Coriolis Constant on the GpKdV Equation

[1]Yusuf Gurefe, [2]Yusuf Pandır, [3]Tolga Akturk

[1] Department of Mathematics, Faculty of Science and Arts, Mersin University, Turkey

[2]Department of Mathematics, Faculty of Science and Arts, Bozok University, Turkey

[3]Department of Mathematics and Science Education, Faculty of Education, Ordu University, Turkey

ygurefe@gmail.com, yusufpandir@gmail.com, tolgaakturkk@gmail.com;

ABSTRACT

This study investigated the propagating wave solutions of a nonlinear partial differential equation (NPDE) called the geophysical Korteweg-de Vries (GpKdV) equation. First, the results obtained on the Coriolis effect due to the equation used in geophysics are mentioned. In the nonlinear equation, there is a Coriolis coefficient representing this effect. The traveling wave solution functions that provide this nonlinear mathematical model are obtained using the Modified Exponential Function Method (MEFM). There are trigonometric, hyperbolic, and rational solution functions. In order to understand the physical behavior of all these functions and interpret them effectively, simulated graphics were drawn with the help of a package program.

Keywords: Geophysics Korteweg-de Vries (GpKdV) Equaiton, Coriolis effect, Modified Exponential Function Method (MEFM).

REFERENCES

1. Rizvi, S. T. R., Seadawy, A. R., Younis, M., Ali, I., Althobaiti, S., & Mahmoud, S. F. (2021). Soliton solutions, Painleve analysis and conservation laws for a nonlinear evolution equation. Results in Physics, 23, 103999.
2. Karunakar, P., & Chakraverty, S. (2019). Effect of Coriolis constant on Geophysical Korteweg-de Vries equation. Journal of Ocean Engineering and Science, 4(2), 113-121.
3. Ak, T., Saha, A., Dhawan, S., & Kara, A. H. (2020). Investigation of Coriolis effect on oceanic flows and its bifurcation via geophysical Korteweg–de Vries equation. Numerical Methods for Partial Differential Equations, 36(6), 1234-1253.
4. Naher, H. ve Abdullah, F. A., 2013. New approach of (G′/G)-expansion method and new approach of generalized (G′/G)-expansion method for nonlinear evolution equation. American Inst of Physics Advances, 3(3),032116.
5. Pandir, Y., Gurefe, Y., & Akturk, T. (2022). New soliton solutions of the nonlinear Radhakrishnan-Kundu-Lakshmanan equation with the beta-derivative. Optical and Quantum Electronics, 54(4), 1-21.

DEGENERATE ARA TRANSFORM

Rabia Kevser Yılmaz[1], **Oğuz Yağcı**[1]

and Recep Şahin[1]

[1] Department of Mathematics, University of Kırıkkale, Kırıkkale, Turkey

rabia.kyilmaz05@gmail.com, oguzyagci26@gmail.com, recepsahin@kku.edu.tr

Abstract

Kim-Kim [1] (Russ. J. Math. Phys. 2017, 24, 241-248) defined and introduced the degenerate Laplace transform, also investigated some of their several properties. Here, we first define and introduce the degenerate Ara transform and establish several properties and relations. We attain degenerate ARA transforms of power functions, degenerate sine, degenerate cosine, degenerate hyperbolic sine, degenerate hyperbolic cosine, degenerate exponential function, and function derivatives. Furthermore, we present that the degenerate ARA transform is the theoretical dual transform to the degenerate Sumudu transform and the degenerate Laplace transform.

Keywords: Degenerate exponential function, degenerate gamma function, degenerate Laplace transform, degenerate Sumudu transform, Laplace transform, Sumudu transform, Ara transform.

REFERENCES

1. Kim, T., Kim, D. S., Degenerate Laplace transform and degenerate gamma function, Russian Journal of Mathematical Physics, 24(2), 241-248, 2017.

2. Kim, T., Kim, D. S., Note on the degenerate gamma function, Russian Journal of Mathematical Physics, 27(3), 352-358, 2020.

3. Kim, Y., Kim, B. M., Jang, L. C., & Kwon, J., A note on modified degenerate gamma and Laplace transformation, Symmetry, 10(10), 471, 2018.

4. Duran, U., Degenerate Sumudu transform and its properties, Filomat, 35(14), 4731-4741, 2021.

5. Yağcı, O., Şahin, R., Degenerate Pochhammer symbol, degenerate Sumudu transform, and degenerate hypergeometric function with applications, Hacettepe Journal of Mathematics and Statistics, 1-18, 2019.

6. Saadeh, R., Qazza, A., Burqan, A., A new integral transform: ARA transform and its properties and applications, Symmetry, 12(6), 925, 2020.

Numerical Investigation of Particle Distribution in a Cylindrical Channel

Eren ÇOLAK[1-2], Ali Keleş[2]

[1] Department of Mechanical Engineering, Hacettepe University, Ankara, Turkey

[2]GKE Energy R&D Center, Muğla, Turkey

erencolak@hacettepe.edu.tr,

Abstract

Particle distributions in a continuous media is an important research subject for thermo-fluid and computational science, because it is related with pressure drops, reaction rates, and with many other physical phenomenon. In this study particle-laden flow of the nano sized particles in a channel investigated with different flow rates (corresponds to Reynolds number 500, 1000, 1500), , different particle diameters (1e6, 5e6, 1e5), and different particle densities (1500, 2500, 3000 kg/m3). For this purpose water as a continuus phase(Eulerian) used as an inert specie to advect the nanoparticles, different particle shapes (spherical, cube, tetrahedron). Navier-Stokes equations are discretized with 2nd order numerical schemes both temporally and spatially. Structural grid is used as computational domain to represent the cylindirical channel. Distribution of these particles are investigated to understand the behavior of the bulk particle flow in a cylindrical channel.

Keywords: Computational Fluid Dynamics, Two-Way Coupling, Particle-Laden flow, OpenSource

Fractional Modeling and Exact Solutions to Analyze Thermal Performance of Fe_3O_4-MoS_2-Water Hybrid Nanofluid Flow Over an Inclined Surface

Talha Anwar[1], **Poom Kumam**[1,*]

[1] Department of Mathematics, Faculty of Science, King Mongkut's University of Technology Thonburi (KMUTT), Bangkok 10140, Thailand

anwartalha80@gmail.com, poom.kum@kmutt.ac.th

Abstract

The core determination of this article is to investigate the augmentation in the radiative heat transfer rate of Fe_3O_4-MoS_2-H_2O hybrid nanofluid specifying a flow over an inclined plate subject to ramped heating and heat generation/consumption effects. The flow of considered hybrid nanofluid is developed due to the ramped motion of the inclined plate that encounters the magnetic effects and immersed in a porous material. The fractional form of energy and momentum equations is obtained by employing the concept of the Atangana-Baleanu fractional derivative. Some unit-free quantities are introduced and the Laplace transform method is operated to construct the exact solutions of these equations. The noteworthy physical significance of involved parameters is discussed with the aid of graphical illustrations. To analyze the behavior of shear stress and heat transfer rate, numerical computations are tabulated in terms of skin friction coefficient and Nusselt number. All the figures and tables are prepared for both ramped and isothermal boundary conditions to highlight the impacts of the ramped heating condition and ramped motion of the inclined plate.

Keywords: Fractional analysis; hybrid nanofluid; ramped conditions; Laplace transform

REFERENCES

1. A. Atangana and D. Baleanu, ''New fractional derivatives with nonlocal and non-singular kernel: Theory and application to heat transfer model,'' *Thermal Sci.*, vol. 20, no. 2, pp. 763–769, 2016
2. K. K. Lawal and H. M. Jibril, ''Unsteady MHD natural convection flow of heat generating/absorbing fluid near a vertical plate with ramped temperature and motion,'' *J. Taibah Univ. Sci.*, vol. 13, no. 1, pp. 433–439, Dec. 2019.

STRUCTURAL DIVERSITY ANALYSIS USING SHANNON'S ENTROPY AND INFORMATION MEASURES

Kürşad Özkan[1], Mehmet Güvenç Negiz[2], İbrahim Özdemir[1]

[1] Isparta University of Applied Sciences, Faculty of Forestry, 32260, Isparta, Turkey

[1] Isparta University of Applied Sciences, Sutculer Prof. Dr. Hasan Gürbüz Vocational School, 32950, Isparta, Turkey

kursadozkan@isparta.edu.tr,

Abstract

The aim of the present study is to propose a holistic approach in order to estimate structural diversity for forest stands. In this context, we created a hypothetical data set composed of six plots. Each hypothetical plot consists of A (vertical layers), B (species presence and abundance values) and C (horizontal layers) factors. Shannon's entropy and information measures (totally 34 diversity partitions) were computed for each of the plots. Next, principal component analysis was employed using the data set obtained from the diversity partitions of all the plots. The first axis of PCA explained majority of total variance. According to the locations of the plots along with the first PCA axis, not only species richness but also their vertically and horizontally distribution homogeneity played important roles. The PCA results are meaningful since high structural diversity is characterized by high number of species and heterogeneity of vertical and horizontal species distribution. Holistic approach seems to be promise to evaluate structural diversity of forest stands. Further studies should therefore be done to confirm the accuracy of this novel approach using real ecological community data.

Keywords: Biodiversity, forest ecosystems, diversity components, entropy, multivariate analysis

REFERENCES

1. Orlóci, L., Anand, M., Pillar, V.D., 2002. Biodiversity analysis: issues, concepts, tecniques. Community Ecology, 3(2): 217-236.
2. Orlóci, L., 2006. Diversity partitions in 3-way sorting: functions, Venn diagram mappings, typical additive series, and examples. Community Ecology, 7(2): 253-257.
3. Shannon, C.E., "A mathematical theory of communication", Bell System Technical Journal, 27(3): 379-423, (1948).

AUTISM SPECTRUM DISORDER DETECTION USING MACHINE LEARNING ALGORITHMS

Afrah Said [1], **Hanife Göker**[2*],

[1] Department of Electrical Electronics Engineering, Kutahya, Turkey

[2] Department of Electrical Electronics Engineering, Kutahya, Turkey

afrah.said@ogr.dpu.edu.tr, hanife.goker@dpu.edu.tr,

Abstract

Autism Spectrum Disorder is a neurological condition that affects brain development. It is characterized by how the person perceives things, learns, interacts, and socializes with others. It is also said that it is a "behavioral disease" because it affects their social communication such as having poor eye contact with others and repetitive behaviors such as repeating certain words or phrases. Detection of this disorder is associated with remarkable healthcare costs and is time-consuming. However, due to the advancement of Machine Learning algorithms in the diagnosis of many diseases, predicting this disorder using these methods could be beneficial as this could result in an early intervention treatment service that can improve a child's development and hence prevent this disorder to keep going on into adolescence and adulthood stage. Therefore, this paper aims to effective detection of autism spectrum disorder using machine learning algorithms. As the signs of this disorder could be seen at an early stage of a child's development, therefore, to carry out this study we used the autism screening data for toddler's dataset that is publicly available for researchers. This dataset consists of 1054 instances and 17 attributes. Feature selection was made on the dataset using InfoGain, GainRatio, SymmetricalUncert, OneR, CfsSubSetEval, and ChiSquared feature selection algorithms, and their performance are compared. According to their performances, the features that can be used in the differential diagnosis of the disorder and the predictive power of each feature are determined. Moreover, the performances of the Support Vector Machine (SVM), Naïve Bayes, K-Nearest Neighbor, Random Forest, and MLP Classifier classification algorithms were compared.

Keywords: Autism spectrum disorder; Machine learning; Feature selection algorithms; Classification; Effective detection.

ON NUMERICAL SOLUTION OF LID DRIVEN PROBLEM IN A SQUARE CAVITY FILLED WITH NANOFLUID

Yücel BALTÜRK [1] Hüseyin DEMİR [2] İnci ÇİLİNGİR SÜNGÜ [3]

[1] Department of Mathematics, Ondokuz Mayıs University, Samsun, Turkey, **yucelbalturk@gmail.com**

[2] Department of Software Engineering, University of Samsun, Samsun, Turkey **huseyin.demir@samsun.edu.tr**

[3] Department of Mathematics and Natural Sciences Education, Ondokuz Mayıs University, Samsun, Turkey, **incicilingir@gmail.com**

Abstract

In this study the numerical solution of wall motion lid driven problem in a square cavity which filled with nanofluid is considered. The equations of motion have been discretised by Finite Difference Method. The system of equations is solved with SOR Method and a MATLAB code is developed and used for simulation. The results are in line with the literature with high compatibility. Indeed, the numerical results obtained from this study are compared with the previous results and presented graphically. The agreement appears to be pretty good. Thus, the solution procedure is implemented for various nanofluids and the graphical results are obtained for different Reynold numbers. The results show that the finite difference method is useful, reliable and accurate for solving this type problems.

Keywords: Finite Difference Method; SOR Method; Nanofluids.

REFERENCES

1. Demir, H., 1996, The Stability Properties of Some Rheological Flows, Doktora Tezi, The University of South Wales, Wales/U.K.

2. Elçioğlu E., B. , Güvenç Yazıcıoğlu, A., Kakaç, S., Nanoakışkan Viskozitesinin Karşılaştırmalı Değerlendirmesi, Isı Bilimi ve Tekniği Dergisi, 34, 1, 137-151, 2014.

3. Mahmoodi, M. et. al., "Free Convectıon of a Nanofluid in a Square Cavity With a Heat Source on the Bottom Wall and Partıally Cooled From Sides, Thermal Science, 18, 283-300, 2014

4. Süngü Çilingir, İ., 2011, Kapalı Bölgede Duvar Hareketli Akım ve Doğal Taşınımlı Isı Geçişinin Sayısal Çözümü, Doktora Tezi, Ondokuzmayıs Üniversitesi Fen Bilimleri Enstitüsü, Samsun, 73s.

5. De Vahl Davis, G., 1983. Natural Convection of Air in a Square Cavity: a Benchmark Numerical Solution. Int. J. for Numerical Methods in Fluids, 3(3), 249-264.

INVESTIGATING THE DIFFICULTIES OF REFUGEE STUDENTS IN MATHEMATICS COURSE

Himmet KORKMAZ [1]

[1] Department of Mathematics and Science Education, Ordu University, Ordu, Turkey

himmetkorkmaz@odu.edu.tr,

Serbay DURAN [2]

[2] Department of Mathematics and Science Education, Adıyaman University, Adıyaman, Turkey

sduran@adiyaman.edu.tr,

Abstract

This study aims to examine the difficulties experienced by refugee students at the middle school level in a mathematics course. In recent decades, there have been large-scale migrations due to war, economic reasons, natural disasters, etc. in several regions in the world. In this context, it is observed that problems arise in the education of the children of people who have to live as refugees in different countries. According to the density of refugees, refugee students are studied in the education system of the country of residence. It is acknowledged from the studies of different researchers that some problems appeared in the adaptation of these students [1, 2]. In this research, a case study was carried out as a qualitative research method with the participation of 85 refugee students in a middle school in the southeastern region of Turkey, where refugees live predominantly, to identify the difficulties and deficiencies experienced by refugee students in the context of mathematics course in Turkey. Data was collected by using semi-constructed interviews with refugee students and examination of students' course documents. Open coding was conducted to obtain themes as a part of qualitative analysis. Some important findings are obtained after the descriptive analysis process. The results demonstrated that refugee students have difficulties in understanding word problems and converting these problems into mathematical language. In addition, students' mathematical knowledge and readiness levels are relatively low than their Turkish counterparts. It is also observed that refugee students have a prejudice against mathematics as native students have. These findings showed that refugee students need more support from teachers and principals to overcome these barriers and difficulties.

Keywords: Mathematics course; Refugee students; Education.

REFERENCES

1. Başaran, Semra Demir. Being the Teacher of Syrian Refugee Students: Teachers' School Experiences. Egitim ve Bilim, Vol: 46, No: 206, 331-354, 2021.
2. Aydin, Hasan & Kaya, Yeliz Kaya. Education for Syrian Refugees: The New Global Issue Facing Teachers and Principals in Turkey, Educational Studies, Vol:55, No:1, 46-71, 2019.

The periodic wave solutions of conformable time-fractional mathematical model by analytical method

Aslı Alkan[2],Tolga Aktürk[1], Hasan Bulut[2]
[1]Department of Mathematics and Science Education, Faculty of Education, Ordu University, Turkey
[2]Department of Mathematics, Faculty of Science, Fırat University, Turkey

[1]tolgaakturkk@gmail.com, [2]alkanasli47@gmail.com; [2]hbulut@firat.edu.tr,

Abstract

In this paper, the wave solutions of the time-fractional Zoomeron equation were get by the modified exponential function method (MEFM). When the solution functions found as a result of the calculations are analyzed, hyperbolic and trigonometric functions with periodic functions are also rational functions. Two-three-dimensional contour and density graphs representing the characteristic feature were drawn by determining the values according with the method in the parameters in the solution functions obtained.

Keywords: Conformable time-fractional Zoomeron equation; modified exponential function method; traveling wave solutions.

References

[1] R. Bertram, The development of fractional calculus 1695-1900. Hist. Math. 4(1) (1977) 75-89.

[2] I. Podlubny, Fractional Differential Equations, Academic Press, London, 1999.

[3] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, Theory and Applications of 120 Fractional Differential Equations, North-HollandMathematical Studies, Elsevier, Amsterdam, 2006.

[4] K. Mille, B. Ross, An Introduction to the Fractional Calculus and Fractional Differential Equations, Wiley, New York, 1993.

[5] R.L. Magin, Fractional Calculus in Bioengineering, BegellHouse Publisher, 125 West Redding, Conn, 2006.

[6] R. Khalil, M.A. Horani, A.M.Y. Sababheh, A new definition of fractional derivative, J. Comput. Appl. Math. 264 (2014) 65-70.

[7] T. Abdeljawad, On conformable fractional calculus, J. Comput. Appl. Math. 279 (2015) 57-66.

[8] K. Kamruzzaman, M.A. Akbar, Exact solutions of the (2+1)-dimensional cubic Klein-Gordon equation and the (3+1)-dimensional Zakharov-Kuznetsov equation using the modified simple equation method. J. Assoc. Arab Univ. Basic Appl. 15 (2014) 74-81.

[9] Odabasi, M., Traveling wave solutions of conformable time-fractional Zakharov–Kuznetsov and Zoomeron equations. Chinese Journal of Physics, 64 (2020) 194-202.

Uncertainty Principles for the Short-time Non-separable Linear Canonical Transform

Mohammad Younus Bhat [1]

[1]Department of Mathematical Sciences, Islamic University of Science and Technology, Kashmir, India

gyounusg@gmail.com,

Abstract

The free metaplectic transformation (FMT) or the nonseparable linear canonical transformation (NSLCT) has gained much popularity in recent times because of its various application in signal processing, paraxial optical systems, digital algorithms, optical encryption and so on. However, the NSLCT is inadequate for localized analysis of non-transient signals, as such, it is imperative to introduce a unique localized trans- form coined as the short-time nonseparable linear canonical transformation (ST-NSLCT). In this paper, we investigate the ST-NSLCT. Firstly, we propose the definition of the ST-NSLCT, and provide the time-frequency analysis of the proposed transform in the NSLCT domain. Secondly, we investigate the basic properties of the proposed transform including the reconstruction formula, Moyal's formula. The emergence of the ST-NSLCT definition and its properties broadens the development of time-frequency representation of higher-dimensional signals theory to a certain extent. Finally, we extend some different uncertainty principles (UP) from quantum mechanics including Lieb's inequality, Pitt's inequality, Hausdorff-Young inequality, Heisenberg's uncertainty principle, Hardy's UP, Beurling's UP, Logarithmic UP, and Nazarov's UP.

Keywords: Short-time non-seperable transformation; Moyals formula; Uncertainty Principle; Nazarov's UP; Hardy's UP; Logarithmic's UP

Some results for a nonlinear transmission line arising in the electronic circuit

Yasin Bozkurt[1], Mehmet Tahir Gulluoglu[2], Haci Mehmet Baskonus[3]

[1]Faculty of Egineering, Harran University, Sanliurfa, Turkey

[2]Faculty of Egineering, Harran University, Sanliurfa, Turkey

[3]Faculty of Education, Harran University, Sanliurfa, Turkey

bradirlov@gmail.com, thrgll@gmail.com, hmbaskonus@gmail.com,

Abstract

In this work, we study on a nonlinear transmission line (NLTL) being an electronic circuit consisting of a high-impedance propagating medium with nonlinear capacitive devices.

***Keywords**:* Nonlinear transmission line; analytical method, Complex function solutions.

REFERENCES

[1] K.K.Ali, R.Yilmazer, H.M.Baskonus, H.Bulut, Modulation instability analysis and analytical solutions to the system of equations for the ion sound and Langmuir waves, Physica Scripta, 95(065602), 1-10, 2020.

[2] S Kumar, Optimal system, dynamical behaviors and exact solution of a nonlinear transmission line model by applying the Lie symmetry method, Indian J Phys, https://doi.org/10.1007/s12648-022-02327-x

[3] Y.M.Li, H.M.Baskonus, A.M.Khudhur, Investigations of the Complex Wave Patterns to the Generalized Calogero-Bogoyavlenskii-Schiff Equation, Soft Computing, 25(10), 6999-7008, 2021.

THERMAL INSULATION PERFORMANCES OF FOAMED MORTARS CONTAINING BENTONITE BLENDED CEMENTS

Yasemin AKGÜN[1]

[1] Department of Renewable Energy, Institue of Science, Ordu University, Ordu, Türkiye

ysmakgun@gmail.com

Abstract

The construction sector, which has intense energy consumption, has an important place in the solutions of world energy security problems. Blended cement production and/or using materials with high thermal performance in terms of insulation in buildings are some of these solutions. Bentonite is an alternative material for blended cement production. On the other hand, thermal performances of foamed mortars/concretes are quite high. The main purpose of this study is to determine how the thermal performance and mechanical properties of foamed mortar are affected by blended cements containing bentonite as alternative material. For this purpose, foamed mortar plates containing bentonite blended cements were produced. Bentonite replacement ratios of blended cements were 0, 5, 10 and 15 wt.%. In the study, the strengths and thermal performances of foamed mortars containing bentonite blended cements were determined as experimental. The experiment results were compared among themselves and with each other. From the experiment results, it was concluded that the thermal performances of the foamed mortars containing bentonite blended cements could be improved up to 10% replacement ratio without compromising the strength.

Keywords: Thermal insulation; Foamed mortar; Bentonite; Blended cement.

REFERENCES

1. Mayur Madhukar Maske1, Dr. Nandkumar Krishnarao Patil, Dr. Amey Deepak Katdare, Review of application of plain and calcined bentonite as a cement blending material in concrete and mortar, Psychology And Education, Vol: 58, No: 1, 5873-5878, 2021.
2. Yue Xie, Jun Li, Zhongyuan Lu, Jun Jiang, Yunhui Niu, Preparation and properties of ultra-lightweight EPS concrete based on pre-saturated bentonite, Construction and Building Materials, Vol: 195, 505–514, 2019.
3. Noureddine Mesboua, Khaled Benyounes, Abdelbaki Benmounah, Study of the impact of bentonite on the physicomechanical and flow properties of cement grout, Cogent Engineering, Vol: 5, 2018.

Think Before You Define: on Conjugation Relations and Other Connections

Arran Fernandez [1]

[1] Department of Mathematics, Eastern Mediterranean University, Famagusta, Northern Cyprus

arran.fernandez@emu.edu.tr

Abstract

Fractional calculus has expanded far beyond the original operators of Riemann–Liouville that it began with: now, fractional integrals and derivatives are being defined every year, with different kernels, behaviours, and applications. But, in the rush to define and use new operators, the nature of mathematics as an interconnected beast is sometimes forgotten. Sometimes, a new operator is defined which is actually the same as an old one, or almost identical to an old one up to some trivial transformation such as a constant multiplication. Sometimes, the transformation required is less trivial, such as an infinite series formula or an algebraic conjugation of operators. This talk will examine some of the many connections between different operators of fractional calculus, and how they can be used to make mathematical work easier, while urging would-be inventors to take account of such connections before declaring an operator to be a new innovation.

Keywords: Fractional integrals; Fractional derivatives; Algebraic conjugation; Philosophy of mathematics.

REFERENCES

1. O. P. Agrawal, “Some generalized fractional calculus operators and their applications in integral equations”", Fractional Calculus and Applied Analysis 15(4) (2012), pp. 700–711.
2. D. Baleanu, A. Fernandez, “On fractional operators and their classifications”, Mathematics 7(9) (2019), 830.
3. A. Fernandez, H.M. Fahad, “On the importance of conjugation relations in fractional calculus”, preprint (2021), under review.
4. M. A. Zaky, A. S. Hendy, D. Suragan, “A note on a class of Caputo fractional differential equations with respect to another function”, Mathematics and Computers in Simulation 196 (2022), pp. 289–295.

DEVELOPMENT OF A NATIONAL SMART METER SYSTEM

Fatih Coşkun[1], Beytullah Topçu[1,] M. Fatih Akay[2]

[1]Nar Sistem Teknolojileri, İstanbul, Turkey

[2]Department of Computer Engineering, Cukurova University, Adana, Turkey

fatih.coskun@nar.com.tr, beytullah.topcu@nar.com.tr, mfakay@cu.edu.tr

Abstract

The electricity distribution sector in Turkey, which has been privatized since the early 2000's, is divided into 21 regions. This distribution network includes consumer and producer subscribers as well as transformers and lighting. Although the number of meters that are still measured at its location is quite high, the rate of meters read by remote reading systems is increasing day by day. Since world standards such as The International Electrotechnical Commission (IEC) and Device Language Message Specification (DLMS) are used in communication with the meters, there is a much cleaner system, while there is no such standard in communication with gateway devices. For this reason, each gateway manufacturer can create its own protocols and some of them even keep these protocols secret. This situation causes many problems. National Smart Meter System (MASS) is a protocol developed to solve this problem. Thanks to this protocol, which is based on JSON type data and designed in accordance with the most modern communication technologies such as MQ Telemetry Transport (MQTT) and NarrowBand-Internet of Things (NB-IoT), a more efficient and reliable energy reading network will be established.

Keywords: Meter, Gateway, IEC, DLMS, MQTT, NB-IoT.

REFERENCES

1. TIMISOARA, A. (2022). A short overview of IoT based energy metering system Part I Internet of things and the energy sector. *Romanian Journal of Civil Engineering Vol*, *13*, 41-4

2. TIMISOARA, A. (2022). A short overview of IoT based energy metering system Part II IoT smart energy meters. *Romanian Journal of Civil Engineering Vol*, *13*, 48-54.

3. Al-Turjman, F., & Abujubbeh, M. (2019). IoT-enabled smart grid via SM: An overview. *Future Generation Computer Systems*, *96*, 579-590.

COMPUTATIONAL AND SIMULATION STUDY OF ORGANIC SOLAR CELLS BASED ON DONOR – ACCEPTOR MOLECULAR MATERIAL

Zakaria El Malki[1], Anass El Karkri[1], Imane EL Mhamedi[1], Mohammed Bouachrine[2], Françoise SereinSpirau[3]

[1]Moulay Ismaïl University, MEM, High School of Technology (ESTM), B.P 3103 Toulal, 50040 Meknes, Morocco.
[2]Moulay Ismaïl University, Faculty of Sciences Meknes, Morocco.
[3]Molecular and Macromolecular Heterochimy, UMR, CNRS 5076, Higher National School of Chemistry, Montpellier, France
z.elmalki@umi.ac.ma

Abstract

In this study, we have designed a series of double organic D-π-A (electron donor-π-conjugated-acceptor) dyes employed in dye-sensitized solar cells (DSSCs), the analysis of microelectronic and photonic structure in a one dimension program [AMPS-1D] has been successfully used to study organic solar cells. The program was used to optimize the performance of organic solar cells based on (carbazole-methylthiophene), benzothiadiazole and thiophene [(Cbz-Mth)- B-T]2 as electron donors, and [6,6]-phenyl-C61-butyric acid methyl ester (PCBM) as an electron acceptor. The optoelectronic properties of these dyes were investigated by using the Density Functional Theory DFT/ B3LYP/6-31G(d,p) method. We studied the influence of the variation of the thickness of the active layer, the temperature, and the density of the effective states of the electrons and the holes in the conduction and valence bands respectively on the performance of the solar cells based on [(Cbz-Mth)-BT]2–PCBM as a photoactive material, sandwiched between a transparent indium tin oxide (ITO) and an aluminum (Al) electrode. The addition of other thiophene units in the copolymer or the deposition of a layer of PEDOT between the anode (ITO) and the active layer, improves the performances of the cell, especially resulting in a remarkable increase in the value of the power conversion efficiency (PCE).

Keywords:Donor-Acceptor; solar cells;PEDOT; Thiophene; Power conversion efficiency.

REFERENCES

1. M. Gratzel, Nature 414, 338-344 (2001).
2. M. Gratzel, J. Photochem. Photobiol. C 4, 145-153 (2003).
3. K. Tanaka, K. Takimiya, T. Otsubo, K. Kawabuchi, S. Kajihara, Y. Harima. Chem Lett. 35, 592-3 (2006).

Application of Mahgoub Transform to Hyers-Ulam Stability of Partial Differential Equations

Emel Biçer [1]

[1] Department of Mathematics, University of Bingol, Bingol, Turkey

ekayabicer@gmail.com,

Abstract

In this study, we investigate Hyers-Ulam stability of some partial differential equations using Mahgoub Transform method. We also show the Hyers-Ulam constants of these partial differential equations.

Keywords: Mahgoub Transform method; Hyers-Ulam stability; partial differential equation.

REFERENCES

1. S. Aggarwal, Comparative study of Mohand and Mahgoub transforms, J. Adv. Res. Appl. Math. Stat., 4, 1–7, 2019.
2. R. Murali, A.P. Selvan, C. Park, Ulam stability of linear differential equations using Fourier Transform, AIMS Math. 5, no.2, 766-780, 2019.

High-Performance of Organic Solar Cells Based on D1-BT-EDOT-BT-D2-A/PCBM Structures

Imane EL Mhamedi [1], Anass El Karkri [1], Zakaria El Malki [1], Mohammed Bouachrine, and Françoise Serein-Spirauc

[1] Moulay Ismail University, High School of Technology, (ESTM), Modeling, Materials and Systems Control (MMSC), Computer Engineering and Intelligent Electrical Systems (2ISEI).) BP: 3103, Toulal, Meknes, Morocco.

i.elmhamedi@edu.umi.ac.ma, a.elkarkri@edu.umi.ac.ma, z.elmalki@est.umi.ac.ma

Abstract

In this paper, six Thiophene-Carbazole-based D-π-A organic dyes were reported. A series of these organic dyes containing identical donor and acceptor group but different π-system. The effect of replacing of thiophene (T) by Phenylene (P), Carbazole (C), Fluorene (F) and Antrhacene (A) as π-system on the physical properties has been focused on. The analysis of microelectronic and photonic structures-one dimension (AMPS-1D) was used to study and simulate the performance of organic heterojunction solar cells based on D1-B-Edot-B-D2-A as electron donors, and [6,6]-phenyl-C61-butyric acid methyl ester (PCBM) as an electron acceptor. The organic photovoltaic cell devices using T3-B-Edot-B-T3-A/PCBM showed the improvement open-circuit voltage Voc, short-circuit current density Jsc, fill factor FF, and power conversion efficiency PCE values for the optimum thickness of 120 nm and the effective state density of electrons and holes of 1021cm-3.The P3-B-Edot-B-T3-A/PCBM and P3-B-Edot-B-P3-A/PCBM based devices exhibited a power conversion efficiency (PCE) of 9.295% and 8.735%, respectively, which outperformed the corresponding T3-B-Edot-B-T3-A/PCBM, Cbz-B-Edot-B-T3-A/PCBM, F-B-Edot-B-T3-A/PCBM, and A-B-Edot-B-T3-A/PCBM based devices (7.330, 6.622, 7.226, and 7.327%). More importantly, the P3-B-Edot-B-T3-A/PCBM and P3-B-Edot-B-P3-A/PCBM -based device delivered the highest PCE of 14.432% and 15.031% respectively, when we deposit a layer of PEDOT between the indium tin oxide (ITO) and the active layer, which is a clear improvement over other result.

Keywords: Donor-Acceptor; Organic Solar cells; PEDOT; Efficiency.

Edgeworth series expansion for option prices

Guillaume Leduc [1]

[1] Department of Mathematics and Statistcis, American University of Sharjah, Sharjah, UAE

gleduc@aus.edu,

Abstract

Edgeworth series are a powerful tool in probability used to describe the asymptotic behavior of standardized sum of independent and identically distributed random variables. However, when evaluating options using tree methods, the standard Edgeworth expansion method does not apply directly because only triangular arrays of independent and identically distributed random variables arise. We show how the extension to triangular arrays of the Kolassa-McCullagh approach to Edgeworth series for lattice distributions combined with recent advances in Edgeworth expansion for triangular arrays yield Edgeworth series for digital and standard European put and call options. We develop closed form formula for the coefficients of $1/\sqrt{n}$ and $1/n$ under a general framework. We provide examples with the most popular trinomial models.

Keywords: Edgeworth series; Trinomial models; Option Pricing.

REFERENCES

1. Leduc, G. (2013). A European option general first-order error formula, The ANZIAM Journal, 54(4), 248-272.
2. Leduc, G. (2016). Can high-order convergence of European option prices be achieved with common CRR-type binomial trees? Bulletin of the Malaysian Mathematical Sciences Society, 39, 1329-1342.
3. Leduc, G. (2016). Option convergence rate with geometric random walks approximations, Stochastic Analysis and Applications, 34(5), 767-791.
4. Leduc, G., & Nurkanovic Hot, M. (2020). Joshi's split tree for option pricing, Risks, 8(3), 81.

AN OPTIMAL CONTROL STRATEGY FOR COVID-19 USING REAL DATA

Mehmet Yavuz [1,*], **Fırat Evirgen** [2], **Fatma Özköse** [3], **Necati Özdemir** [2]

[1] Department of Mathematics and Computer Sciences, Necmettin Erbakan University, Konya, Turkey,
[2] Department of Mathematics, Balıkesir University, Balıkesir, Turkey
[3] Department of Mathematics, Erciyes University, Kayseri, Turkey

mehmetyavuz@erbakan.edu.tr, fevirgen@balikesir.edu.tr, fpeker@erciyes.edu.tr, nozdemir@balikesir.edu.tr

Abstract

In this study, a new COVID-19 model is proposed. In this context, an integer-order pandemic model is developed to examine the spread of COVID-19 with and without Omicron variant and its relationship with heart attack using real data from Turkey. In the model, an optimal control strategy representing the social distance is considered. Then, the existence, uniqueness, positivity and boundedness of the solution are provided. Qualitative analysis of the proposed model is investigated. In order to estimate the values of the model parameters, the least squares curve fitting method is utilized. Finally, the numerical simulations are presented to examine the dynamic behaviour of the mentioned COVID-19 model of integer-order.

Keywords: COVID-19 mathematical model, optimal control, parameter estimation, numerical simulations.

Acknowledgments: This study has been supported by Necmettin Erbakan University Research Grant No. BAP 211215005.

REFERENCES

1. Özköse, F., Yavuz, M., Şenel, M. T., & Habbireeh, R. (2022). Fractional order modelling of omicron SARS-CoV-2 variant containing heart attack effect using real data from the United Kingdom. Chaos, Solitons & Fractals, 157, 111954.
2. Kaymak, Ö. Ö., Kaymak, Y., & Özdemir, N. (2021). Forecasting of the istanbul stock exchange (ISE) return with a golden ratio model in the epidemic of COVID-19. Appl. Comput. Math, 20(1), 95-107.
3. Uçar, E., Uçar, S., Evirgen, F., & Özdemir, N. (2021). A Fractional SAIDR Model in the Frame of Atangana–Baleanu Derivative. Fractal and Fractional, 5(2), 32.

* Corresponding Author

Spectral Expansions for a Class Second Order Differential Operator with Eigenvalue Parameter Contained in the Boundary Conditions

Kh. R. Mamedov[1], **U. Demirbilek**[2]

[1] Department of Mathematics, University of Iğdır, Iğdır, Turkey
[2] Department of Mathematics, University of Mersin, Mersin, Turkey
hanlar.residoglu@igdir.edu.tr, udemirbilek@mersin.edu.tr

Abstract

We consider the singular boundary value problem generated by differential equation

$$-z'' + p(x)z = \mu\gamma(x)z, \qquad 0 < x < \infty \tag{1}$$

and the boundary condition

$$\frac{z'(0)}{z(0)} = f(\mu) \tag{2}$$

where $\mu = \lambda^2$ is a spectral parameter, $p(x) \in L_1(0,\infty)$ is a real valued function and satisfying the condition

$$\int_0^{+\infty} x|p(x)|dx < \infty.$$

Here $\gamma(x)$ is real positive piecewise continuous function and $f(\mu)$ in the condition (2) is a meromorphic function.

In this work, the scattering problem of the boundary value problem (1)-(2) is studied. The expansion formula according to the eigenfunctions under certain conditions is obtained by applying Titchmarsh method [1]. Inverse problems of scattering theory were investigated in [2,3].

Keywords: Expansion formula; eigenfunction; scattering problem.

REFERENCES

1. E. C., Titchmarsh, Eigenfunction Expansions Associated with Second-Order Differential Equations, Oxford Clarendon Press, 1962.

2. Y. Yang, G.Wei, Inverse Scattering Problems for Sturm–Liouville Operators with Spectral Parameter Dependent on Boundary Conditions, Mathematical Notes, Vol: 103, No:1, 59–66,2018.

3. Kh. R. Mamedov, On an Inverse Scattering Problem for a Discontinuous Sturm-Liouville Equation with a Spectral Parameter in the Boundary Condition, Boundary Value Problem, 17, 2010.

Application of tan $\left(\frac{\Phi(\xi)}{2}\right)$-expansion scheme for solving two nonlinear partial differential equations arising in chemical and biological phenomena

Onur Alp İlhan [1,*] **and Jalil Manafian** [2,3]

[1]Department of Mathematics, Faculty of Education, Erciyes University, 38039-Melikgazi-Kayseri, Turkey

E-mail: oailhan@erciyes.edu.tr

* Corresponding author.

[2]Department of Applied Mathematics, Faculty of Mathematical Sciences, University of Tabriz, Tabriz, Iran

[3]Natural Sciences Faculty, Lankaran State University, 50, H. Aslanov str., Lankaran, Azerbaijan

E-mail: j_manafianheris@tabrizu.ac.ir

Abstract

In this paper, the reaction-diffusion Brusselator system and the Radhakrishnan, Kundu and Laskshmanan (RKL) equation. Based on the new technique namely tan $\left(\frac{\Phi(\xi)}{2}\right)$-expansion method the solution procedure of nonlinear evolution equations in chemical and biological phenomena are investigated. We obtained the plenty of exact solutions for mentioned.

Keywords: tan $\left(\frac{\Phi(\xi)}{2}\right)$-expansion technique; reaction-diffusion Brusselator system; the Radhakrishnan, Kundu and Laskshmanan (RKL) equation.

REFERENCES

[1] H. Khan, H. Jafari, R. A. Khan, H. Tajadodi and S. J. Johnston, Numerical solutions of the nonlinear fractional-order Brusselator system by Bernstein polynomials, The Sci. World J., 2014 (2014) Article ID 257484, 7 pages.

[2] A. M. Wazwaz, The decomposition method applied to systems of partial differential equations and to the Reaction-Diffusion Brusselator model, Appl. Math. Comput., 110 (2000) 251-264.

[3] M. S. H. Chowdhurya, T. H. Hassanb, S. Mawac, A new application of homotopy perturbation method to the Reaction-diffusion Brusselator model, Proc. Soc. Behavioral Sci., 8 (2010) 648-653.

[4] H. Jafari, A. Kadem, and D. Baleanu, Variational iteration method for a fractional-order Brusselator system, Abs. Appl. Anal., 2014, Article ID 496323, 6 pages, 2014.

[5] F. Khani, F. Samadi and S. Hamedi-Nezhad, New exact solutions of the Brusselator reaction diffusion model using the Exp-function method, Math. Problems Eng., 2009, Article ID 346461, 9 pages, 2009.

ANALYSIS OF THE SOLUTIONS OF A SPACE-TIME FRACTIONAL MATHEMATICAL MODEL WITH AN ANALYTICAL METHOD

Özlem Kırcı

Department of Mathematics, Faculty of Arts and Sciences, Kırklareli University, Turkey

ozlem.isik@klu.edu.tr

Abstract

In the present paper we have considered a nonlinear equation including the modified Riemann-Liouville fractional derivative of Jumarie, namely space-time fractional Calogero-Degasperis (CD) equation. The analytical solutions of CD equation which describes the (2+1)-dimensional interaction of a Riemann wave propagating along the y-axis with a long wave along the x-axis, are constructed by the modified exponential function method. The proposed method is easy to implement and powerful. Consequently three types of exact solutions are obtained as hyperbolic function solutions, trigonometric function solutions and rational solutions. The appropriate graphical results are presented to the related cases.

Keywords: Modified exponential function method; Space-time fractional Calogero-Degasperis equation; Traveling wave solutions.

REFERENCES

1. Guner, O., Exact Travelling Wave Solutions To The Space-Time Fractional Calogero-Degasperis Equation Using Different Methods, Journal of Applied Analysis & Computation, Vol:9, No:2, 428-439, 2019.
2. Sahoo, S., Saha Ray, S., Lie symmetries analysis and conservation laws for the fractional Calogero–Degasperis–Ibragimov–Shabat equation, International Journal of Geometric Methods in Modern Physics, Vol:15, No:7, 2018.
3. Kaur, L., WazWaz, A.M., Dynamical Analysis Of Soliton Solutions For Space-Time Fractional Calogero-Degasperis And Sharma-Tasso-Olver Equations, Romanian Reports in Physics, 2022(in press).
4. Arefin, M.A., Saeed, Md. A., Akbar, M.A., Uddin, M.H., Analytical behavior of weakly dispersive surface and internal waves in the ocean, Journal of Ocean Engineering and Science, 2021, ISSN 2468-0133, https://doi.org/10.1016/j.joes.2021.08.012.
5. Mohyud-Din, S.T., Nawaz, T., Azhar, E., Akbar, M.A., Fractional sub-equation method to space–time fractional Calogero-Degasperis and potential Kadomtsev-Petviashvili equations, Journal of Taibah University for Science, Volume:11, No:2, 258-263, 2017.

Solitary wave solutions to a nonlinear mathematical model

Hasan Bulut[1], Sümeyye Kılbitmez[1], Tukur A Sulaiman[2,3], Abdullahi Yusuf[2,3]

[1]Department of Mathematics, Firat University, Elazig, Turkey

[2]Department of Computer Engineering, Biruni University, Istanbul, Turkey

[3]Department of Mathematics, Federal University Dutse, Jigawa, Nigeria

Abstract

The solitary wave solutions to a nonlinear mathematical model arising in mathematical physics are established in this study. To construct such novel solutions, we employ robust integration technique called modified Sardar method. The novel soliton control structure has been studied in order to explain specific physical problems. Three-dimensional and contour graphs have been plotted using suitable choice of the values of parameters involved.

Keywords: Nonlinear equation; Sardar method; Solitary waves.

[1] John, F.: Partial Differential Equations. Applied Mathematical Sciences, 4th edn. Springer, Berlin (1991)

[2] Protter, M.H., Weinberger, H.F.: Maximum Principles in Differential Equations, Corrected Reprint of the1967 Original. Springer, New York (1984)

[3] Schatzman, M.: Numerical Analysis—A Mathematical Introduction. Oxford University Press, Oxford (2002)

[4] Smith, G.D.: Numerical Solutions of Partial Differential Equations, Finite Difference Methods. Oxford Uni-versity Press, New York (1985)

Exact solutions to the coupled-Higgs equation with neutral scalar mesons cooperating in a system

Karmina K. Ali[1,2], Sibel Sehriban Ataş[1], Hasan Bulut[1]

[1]Department of Mathematics, Faculty of Science, Firat University, Elazig, Turkey

[2]Department of Mathematics, Faculty of Science, University of Zakho, Zakho, Iraq

[1]karmina.ali@uoz.edu.krd, [2] sibel.s.atas@gmail.com, [1] hbulut@frat.edu.tr

Abstract

In the present study, through the generalized exponential rational function method (GERFM), we constitute exact wave solutions for the coupled-Higgs equation (CHE). The proposed method is one of the most powerful method to construct abundant exact solutions for nonlinear partial differential equations. The obtained wave solutions include hyperbolic, trigonometry, and exponential functions solutions. Furthermore, we draw three-dimensional surfaces and counter plots of the obtained solutions using the appropriate value for involved parameters.

Keywords: The coupled-Higgs equation, generalized exponential rational function method (GERFM), hyperbolic, trigonometry, and exponential functions solutions.

REFERENCES

[1] A. Yokus and M. Tuz, "An application of a new version of (G ′/ G)-expansion method," in *AIP Conference Proceedings*, 2017, vol. 1798, no. 1, p. 20165.

[2] T. A. Sulaiman, H. Bulut, and S. S. Atas, "Optical solitons to the fractional Schrdinger-Hirota equation," *Applied Mathematics and Nonlinear Sciences*, vol. 4, no. 2, pp. 535–542, 2019.

[3] H. F. Ismael, W.-X. Ma, and H. Bulut, "Dynamics of soliton and mixed lump-soliton waves to a generalized Bogoyavlensky-Konopelchenko equation," *Physica Scripta*, vol. 96, no. 3, 2021.

[4] K. K. Ali, R. Yilmazer, A. Yokus, and H. Bulut, "Analytical solutions for the (3+ 1)-dimensional nonlinear extended quantum Zakharov–Kuznetsov equation in plasma physics," *Physica A: Statistical Mechanics and its Applications*, vol. 548, p. 124327, 2020.

FORMULATION OF A 7-DOF HALF-CAR MODEL WITH NONLINEAR SPRINGS AND DAMPERS FOR RIDE COMFORT ANALYSIS OF IN-WHEEL-MOTOR ELECTRIC VEHICLES

Mustafa Özdemir [1], **Eralp Osman Erdoğan** [2]

[1] Department of Mechanical Engineering, Faculty of Engineering, Marmara University, Recep Tayyip Erdoğan Campus, 34854 Maltepe, Istanbul, Turkey
E-mail: mustafa.ozdemir@marmara.edu.tr
ORCID iD: https://orcid.org/0000-0002-4981-9573

[2] Department of Mechanical Engineering (English), Institute of Pure and Applied Sciences, Marmara University, Göztepe Campus, 34722 Kadıköy, Istanbul, Turkey
E-mail: eralp.osman@marun.edu.tr
ORCID iD: https://orcid.org/0000-0002-6273-5964

Corresponding author: Mustafa Özdemir

Abstract

In this paper, we formulate a seven-degree-of-freedom half-car model for ride comfort analysis of electric vehicles with in-wheel motors. All the springs and dampers in the model are considered to exhibit cubic nonlinearities in addition to their linear characteristics. In this manner the nature of the corresponding vehicle components is described realistically. The equations of motion are obtained by using the Lagrangian method. The resulting system of ordinary differential equations can be numerically integrated to simulate vibrations due to road irregularities and unbalance of rotating parts.

Keywords: Electric vehicle; Half-car model; Vehicle dynamics.

ACKNOWLEDGEMENTS

This paper is based on a part of the Master of Science Thesis of the second author (Eralp Osman Erdoğan), which is in preparation in the Department of Mechanical Engineering (English) at Marmara University Institute of Pure and Applied Sciences under the supervision of the first author (Assoc. Prof. Dr. Mustafa Özdemir).

REFERENCES

1. Mehrdad Ehsani, Yimin Gao, Ali Emadi, Modern Electric, Hybrid Electric, and Fuel Cell Vehicles: Fundamentals, Theory, and Design, 2nd Edition, CRC Press, Boca Raton, FL, USA, 2010.
2. S. Narayanan, S. Senthil, Stochastic optimal active control of a 2-DOF quarter car model with non-linear passive suspension elements, Journal of Sound and Vibration, vol. 211, no. 3, pp. 495–506, 1998.
3. Ernest O. Doebelin, System Dynamics: Modeling, Analysis, Simulation, Design, Marcel Dekker, New York, USA, 1998.
4. Reza N. Jazar, Vehicle Dynamics: Theory and Application, 2nd Edition, Springer, New York, USA, 2014.
5. Jinhui Jiang, Mohammed Seaid, M Shadi Mohamed, Hongqiu Li, Inverse algorithm for real-time road roughness estimation for autonomous vehicles, Archive of Applied Mechanics, vol. 90, no. 6, pp. 1333–1348, 2020.
6. Le Van Quynh, Bui Van Cuong, Nguyen Van Liem, Le Xuan Long, Pham Thi Thanh Dung, Effect of in-wheel motor suspension system on electric vehicle ride comfort, Vibroengineering PROCEDIA, vol. 29, pp. 148–152, 2019.
7. Peiling Wang, Effect of electric battery mass distribution on electric vehicle movement safety, Vibroengineering PROCEDIA, vol. 33, pp. 78–83, 2020.
8. Shuting Huang, Vanliem Nguyen, Influence of dynamic parameters of electric-vehicles on the ride comfort under different operation conditions, Journal of Mechanical Engineering, Automation and Control Systems, vol. 2, no. 1, pp. 1–8, 2021.
9. Edward J. Haug, Jasbir S. Arora, Applied Optimal Design: Mechanical and Structural Systems, Wiley, New York, USA, 1979.
10. Richard L. Burden, J. Douglas Faires, Numerical Analysis, 9th Edition, Brooks/Cole, Cengage Learning, Boston, MA, USA, 2011.

ANALYSIS OF CRYPTOGRAPHIC HASH FUNCTIONS FOR BLOCKCHAIN TECHNOLOGY

Muharrem Tuncay GENÇOĞLU [1]

[1] Thecnical Sciences Locational School, University of Firat, Elazig, Turkey

mtgencoglu23@firat.edu.tr,

Abstract

Blockchain is one of the most interestingly evolving technologies today. This study presents a mathematical analysis of cryptographic hash functions, which is the most important element of the security foundations of this technology.

Keywords: hash functions, blockchain, cryptographic hash functions analysis.

REFERENCES

1. PanWang, BoTian, Kun Sun, Feng-Hua Qi, Thetitle of Paper, Applied Mathematics and
2. Paar, C., Pelzl, J. Understanding Cryptography. A Textbook for Students and practitioners; Springer- Heidelberg, Germany, 2010.
3. Rosen, K. An INTRODUCTION to CRYPTOGRAPHY. Taylor&Francis Group, USA. 2007.
4. NIST. Secure Hash Standard (SHS). Federal Information Processing Standard Publication. FIPS 180-4. 2015. Available online: https://nvlpubs.nist.gov/nistpubs/FIPS/NIST.FIPS.180-4.pdf (accessed on 15 August 2021).
5. Dwork, C.; Naor, M. Pricing via Processing or Combatting Junk Mail. In Advances in Cryptology-Proceedings of Crypto'92; Brickell, E.F., Ed.; Springer: Berlin, Germany, 1993; pp. 139–147.
6. Jakobsson, M.; Juels, A. Proofs of Work and Bread Pudding Protocols (Extended Abstract). In Secure Information Networks; Preneel, B., Ed.; Springer: Boston, MA, USA, 1999; pp. 258–272.

*Abstract Submission should be prepared only **1 page**.

Application of Fractional Operators to Modified Special Functions

Enes Ata [1]

[1] Department of Mathematics, University of Kırşehir Ahi Evran, Kırşehir, Turkey

enesata.tr@gmail.com

Abstract

The motive of this paper is to apply Riemann-Liouville and Erdelyi-Kober fractional integral operators and Riemann-Liouville and Caputo fractional derivative operators to modified beta, modified Gauss hypergeometric and modified confluent hypergeometric functions.

Keywords: Gamma function; Beta function; Gauss hypergeometric function; Confluent hypergeometric function; Generalized M-series; Fractional operators.

REFERENCES

1. E. Ata, İ. O. Kıymaz, A study on certain properties of generalized special functions defined by Fox-Wright function, Applied Mathematics and Nonlinear Sciences, Vol:5, No:1, 147-162, 2020.
2. E. Ata, Generalized beta function defined by Wright function, arXiv:1803.03121v3 [math.CA], 2021.
3. E. Ata, Modified special functions defined by generalized M-series and their properties, arXiv:2201.00867v1 [math.CA], 2022.
4. A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, Theory and Applications of Fractional Differential, North-Holland Mathematics Studies 204, 2006.
5. M. Sharma, R. Jain, A note on a generalized M-series as a special function of fractional calculus, Fractional Calculus and Applied Analysis, Vol:12, No:4, 449-452, 2009.

A STOCHASTIC MODEL FOR THE ENERGY-EFFICIENT SERVER CONSOLIDATION

Özge Erden [1]

[1] Department of Industrial Engineering, Yildiz Technical University, Istanbul, Turkey

ozgerdn19@gmail.com

Abstract

Data centers form the infrastructure of information technologies (IT). Most of the Internet Protocol (IP) address, for example, traffic passes through these data centers. Moreover, the IP address data is stored at data centers. To cope with this increasing flow, many processing and storage servers are needed. However, data centers have large investment costs and high energy use. The most important data center cost factors are operating cost and energy cost. Here, there are many factors considered for energy usage in servers and different resource types like CPU, bandwidth, and memory. The CPU, on the other hand, is generally the most energy-consuming part of servers. The most important technology used to save energy in physical servers is virtualization. Virtualization enables many virtual servers to run on a single physical server. In this study, we aim to meet the resource needs of virtual servers by minimizing energy consumption in a datacenter. In other words, here we mention how much energy physical servers consume according to the amount of CPU used. Here, it is assumed that there is a linear relationship between CPU and energy usage. The workloads of the services running on virtual servers change in time. With the change of these workloads, the amount of CPU used by virtual servers' changes. For this purpose, we propose a stochastic model for server consolidation under resource demand uncertainty. This model, which presents the ambiguous workload, is a scenario-based stochastic programming model. According to the result, in the consolidation of virtual servers with uncertain resource demand, energy costs are minimized.

Keywords: Server Consolidation, Stochastic, Modelling, Data center.

A NEW GENERALIZED DISTRIBUTION FOR ESTIMATION OF FLOOD AND EARTHQUAKE SIZES IN TURKEY

Gamze Özel[1], Barış Burçin Demir[2]

[1] Department of Statistics, Hacettepe University, Ankara, Türkiye

[2]Department of Actuarial Sciences, Hacettepe University, Ankara, Türkiye

gamzeozl@hacettepe.edu.tr, barisburcin@gmail.com

Abstract

It is known that there are many types among natural disasters. Especially earthquakes and floods are among the most common natural disasters. Among the disasters that occurred in Turkey, the landslide is the most in number. In terms of the number of people affected by disasters, landslides and earthquakes are in the first place in terms of earthquakes and the number of settlements affected by disasters. Therefore, in this study, it is aimed to contribute to the minimization of the risks of these disasters by examining a new distribution that is thought to be more flexible for the estimation of the magnitudes of these two disaster types for Turkey.

Generalized distributions are probabilistic models with a wide range of applications in modeling skewed data in many different fields from science and engineering to health sciences. Since they are flexible, they have more useful statistical features in terms of real data modeling. In this study, a new generalized distribution based on the sine function was obtained, and the skewness, kurtosis, peak value, survival and hazard functions, and maximum likelihood estimates of this distribution were obtained. Then, the graphs of the distributions were drawn according to different parameter values and the flexibility of the proposed distribution was examined. Finally, the superiority of the generalized distribution over the existing distributions in the literature is demonstrated in practice on real datasets based on earthquake and landslide data. It has been shown statistically that the proposed new distribution based on the sinus function is more flexible than the distributions in the literature in estimating the magnitude of both disaster types.

Keywords: Earthquake, Flood, Maximum Likelihood Estimate, Generalized Distributions

AN OPTIMAL CONTROL PROBLEM FOR n-DIMENSIONAL SCHRODINGER EQUATION WITH SPECIFIC GRADIENT TERM

Nigar Yildirim Aksoy[1], Ercan Çelik[2] and Merve Zengin[1]

[1] Department of Mathematics, Kafkas University, Kars-Turkey

[2] Kyrgyz-Turkish Manas University, Department of Applied Mathematics and Informatics, Bishkek-Kyrgyzstan

nyaksoy55@hotmail.com, ercan.celik@manas.edu.kg, merveezengin14@gmail.com

Abstract

In this paper, we study an optimal control problem n-dimensional Schrödinger equation with specific gradient term which arise in atomic and molecular physics, pyhsical chemistry, nuclear physics and optics. The objective functional is given by an integral criterion over boundary of domain. In the present paper, we prove the solvability of considered optimal control problem in which control is a complex-valued function such that its real and imaginary parts are measurable bounded functions.

Keywords: Optimal control; Schrödinger equation; objective functional.

REFERENCES

1. Anatoliy G. Butkovskiy, Yu. I. Samoilenko, Control of quantum-mechanical processes and systems: Mathematics and its Applications. Dordrecht, Netherlands (NL): Kluwer Academic Publishers; 1990.

2. Victor Michailovich Zhuravlev, Nonlinear waves in multicomponent systems with dispersion and diffusion. Ulyanovsk: UlSU Publishing House; 2001.

3. Gökce D. Kücük, Gabil Yagub, Ercan Celik, On the existence and uniqueness of the solution of an optimal control problem for Schrödinger equation. Discrete Contin Dyn Syst-S. Vol:12, No:3, 503-512, 2019.

MARKOV RENEWAL MODEL APPLIED TO THE ESTIMATION OF EARTHQUAKE OCCURRENCES

Gamze Özel[1], Ceren Ünal[2]

[1,2] Department of Statistics, Hacettepe University, Ankara, Türkiye

gamzeozl@hacettepe.edu.tr, cerenunal@hacettepe.edu.tr

Abstract

The process of earthquake occurrences can be represented with many analytical models (Votsi et al.[1]). Some models are based on empirical observations of preliminary events, other models on physical modelling of the earthquake process and some on statistical analysis of patterns of seismicity.

The theory of Markov processes can ben applied in various fields since the Markov property is very intuitive. If we know the past and present of a system, then the future development of the system is only determined by its present state (Listwon and Saint-Pierre[2]). Hence, the history of the system does not play a role in its future development. It is called as the memoryless property. However, the Markov property has its limitations. It enforces restrictions on the distribution of the sojourn time in a state, which is exponentially distributed in the continuous case. This is a disadvantage when Markov processes are applied in real-life applications. Therefore, in this study, we use the Markov renewal process to apply a previously proposed method for seismic disaster detection and specifically to model large earthquakes consistently using high-intensity earthquake data that occurred in Turkey during the 20th century. Among the common non-Poisson models, the Markov renewal process proposed by Garavaglia and Pavani. A mixture of exponential and Weibull distributions is used for interevent times by R program

Keywords: Earthquake, Markov Process, Renewal Process, Weibull Distribution

REFERENCES

[1] Irene Votsi, Nikolaos Limnios, George Tsaklidis & Eleftheria Papadimitiou, Estimation of the Expected Number of Earthquake Occurences Based on Semi-Markov Models, Springer, 2011.

[2] Agnieszka Listwon, Philippe Saint-Pierre, SemiMarkov: An R Package for Parametric Estimation in MultiState Semi-Markov Models, Journal of Statistical Software, 2015.

An efficient linearized approximation technique for the non-linear two-dimensional coupled Burgers equation

Sibel Özer[1], **Selçuk Kutluay**[1]

[1] Department of Mathematics, University of Inönü, Malatya, Turkey

sibel.ozer@inonu.edu.tr

selcuk.kutluay@inonu.edu.tr

Abstract

In this article, the numerical solution of the non-linear two-dimensional coupled Burgers equation given by initial and boundary conditions has been carried out by linearized implicit finite difference method. Fourier stability analysis of the proposed method has been performed and four different models are applied to the problem. In order to demonstrate the accuracy and validety of the method, the error norms L_2, L_∞ and E_R with some numerical results have been calculated and compared with those available in the literatüre for the same parameters.

Keywords: Two-dimensional Burgers equation; Linearized implicit finite difference; Fourier stability analysis.

REFERENCES

1. C. A. J. Fletcher, A comparison of finite element and finite difference solutions of the one- and two-dimensional Burgers equations, Journal of Computational Physics, Vol:51,159-188, 1983.
2. A. R. Bahadır, A fully implicit finite-difference scheme for two-dimensional Burgers equations, Applied Mathematics and Computation, Vol:137, 131-137, 2003.
3. R. C. Mittal, A. Tripathi, Numerical solutions of two-dimensional Burgers equations using modified Bi-cubic B-spline finite elements, Engineering Computations, Vol:32, No:5, 1275-1306, 2015.
4. M. Tamsir, V. K. Srivastava, R. Jiwari, An algorithm based on exponential modified cubic B-spline differential quadrature method for nonlinear Burgers equation, Applied Mathematics and Computation, Vol:290, 111-124, 2016.
5. E. Ngondiep, An efficient three-level explicit time-split scheme for solving two-dimensional unsteady nonlinear coupled Burgers equations, Int. J. Numer. Meth. Fluids, Vol:92, 266-284, 2020.

Solutions to the (n+1)-dimensional NLSE equation with M-truncated derivative

Hasan Bulut[1], Ercan çelik[2], Salisu Ibrahim[3], Tukur Abdulkadir Sulaiman[4,5], Abdullahi Yusuf[4,5]

[1]Department of Mathematics, Firat University, Elazig, Turkey

[2]Department of Mathematics, Atatürk Unıversıty, Erzurum

[3]Department of Mathematics Education, Tishk International University - Erbil, Kurdistan Region, Iraq

[4]Department of Computer Engineering, Biruni University, Istanbul, Turkey

[5]Department of Mathematics, Federal University Dutse, Jigawa, Nigeria

Abstract

The solutions with M-truncated derivatives to the (n+1)-diminsional nonlinear Shrodinger equation (NLSE) with Kerr and power law nonlinearities are derived in this study. To reach such solutions, we employ the undetermined coefficient approach. The novel soliton control structure has been studied in order to explain particular physical difficulties. These discoveries have been used to the transmission of long-wave and high-power telecommunication networks, which has proven to be beneficial. The circumstances required for the existence of these solitons are provided.

References

1. M. Eslami, M. Mirzazadeh, Topological 1-soliton of nonlinear Schrodinger equation with dual power nonlinearity in optical fibers, Eur. Phys. J. Plus 128 (2013) 141-147.

2. E. Zerrad, A. Biswas, R. Kohl, D. Milovic, Optical solitons by He's variational principle in a non-kerr law media, J. Infrared Millim. Terahertz Waves 30(5) (2009) 526–537.

3. A. Biswas, P. Suarez, Exact 1-soliton solution of the Zakharov-Kuznetsov equation in plasmas with power law nonlinearity, Appl. Math. Comput. 217(17) (2011) 7372–7375.

New exact solutions of the Coupled-Higgs equation using the ϕ^6-model expansion method

Sibel Tarla[1], **Karmina K. Ali**[1,2], **Resat Yilmazer**[1]

[1] Department of Mathematics, University of Firat, Elazig, Turkey

[2]Department of Mathematics, University of Zakho, Zakho, Iraq

sibeltarla@gmail.com, karmina.ali@uoz.edu.krd, ryilmazer@firat.edu.tr,

Abstract

In this study, we investigate the coupled-Higgs equation (CHE), which represents a system of preserved scalar nucleons collaborating with unbiased scalar mesons with the use of symbolic computation, the ϕ^6-model expansion method. The trigonometric and hyperbolic functions are obtained based on $m \to 1$ and $m \to 0$ are degenerated state of the Jacobi elliptic functions. All obtained solutions verify the partial differential equation under the given constrained conditions. To shed more light on the dynamic behavior of the obtained solutions, we illustrated 3D surfaces and 2D graphics of some solutions constructed with the help of a computer program.

Keywords: Exact solutions; Coupled-Higgs equation; The ϕ^6-model expansion method.

REFERENCES

[1] Rizvi, S. T. R., Seadawy, A. R., Ashraf, M. A., Bashir, A., Younis, M., & Baleanu, D. Multi-wave, homoclinic breather, M-shaped rational and other solitary wave solutions for coupled-Higgs equation. The European Physical Journal Special Topics, 230(18), 3519-3532, 2021.

[2] Singh, S. S. Soliton solutions of coupled Higgs field equation via the trial equation method. Int. J. Phys. Res., 7, 106-110, 2019.

[3] Islam, M., Barman, H. K., & Akbar, M. A. Search for interactions of phenomena described by the coupled-Higgs field equation through analytical solutions. Optical and Quantum Electronics, 52(11), 1-19, 2020.

[4] Zayed, E. M., Al-Nowehy, A. G., Elshater, M. E. New ϕ^6-model expansion method and its applications to the resonant nonlinear Schrödinger equation with parabolic law nonlinearity. The European Physical Journal Plus, 133(10), 417, 2018.

[5] Zayed, E. M., Shohib, R. M., El-Horbaty, M. M., Biswas, A., Yıldırımm, Y., Alshomrani, A. S., Belic, M. R. Optical solitons in birefringent filbers with quadratic-cubic refractive index by ϕ^6 -model expansion. Optik, 202, 163620, 2020.

PERFORMANCE EVALUATION OF MACHINE LEARNING ALGORITHMS FOR DETECTING EATING AND PHYSICAL STATUS BASED OBESITY STAGES

Şükrü KİTİŞ[1], Hanife GÖKER[1]

[1] Simav Technology Faculty, Kütahya Dumlupınar University, Kütahya, Turkey

sukru.kitis@dpu.edu.tr

Abstract

Obesity is the excess of body weight relative to height above the desired level as a result of excessive increase in the ratio of body fat mass to lean mass. It causes many health problems due to its negative effects on body systems (endocrine system, cardiovascular system, respiratory system, gastrointestinal system, skin, genitourinary system, musculoskeletal system) and psychosocial status. In this study is aimed to effective detection of the eating and physical condition based obesity stages using machine learning algorithms. The dataset contains data for the estimation of obesity levels in individuals from the countries of Mexico, Peru and Colombia, available as open source from UCI Machine Learning Repository. The data contains 17 attributes and 2111 records. The records are labeled with the class variable obesity stages, that allows classification of the data using the values of insufficient weight, normal weight, overweight level I, overweight level II, obesity type I, obesity type II and obesity type III. The 10-fold cross-validation method was used to validate the model and the performances of the Support Vector Machine, Random Forest, and Multilayer Perceptron classification algorithms were compared. It has been determined that the highest performance among the algorithms whose performances are compared belongs to the Random Forest Algorithm (95.78%).

Keywords: Machine learning, obesity, UCI, Random Forest

RECTANGULAR MICROSTRIP ANTENNA DESIGN FOR WIRELESS TECHNOLOGIES

Mustafa Mutlu[1] and Çetin Kurnaz[2]

[1]Vocational School of Technical Sciences

Ordu University, Ordu, Turkey

mustafamutlu@odu.edu.tr

[2]Department of Electrical and Electronics Engineering

Ondokuz Mayıs University, Samsun, Turkey

ckurnaz@omu.edu.tr

Abstract

A rectangular microstrip antenna operating for wireless technologies (Wi-Fi, Bluetooth and ISM) is designed which is made of PF-4 having dielectric constant of 1.06, loss tangent of 0.0001 and antenna patch and ground part consist of copper band whose one side is sticky. Operating frequency is 2.45 GHz. Its operating band is between 2.4136 GHz and 2.4791 GHz. Its gain at the critical frequency is 5.109 dB.

Keywords: Wifi; Bluetooth; ISM; Microstrip Antenna.

REFERENCES

1. Taiji Kondo, Hiroshi Fujita, Makoto Yoshida and Tamio Saito, Technology for WiFi/Bluetooth and WiMAX coexistence, 2010.
1. Palandöken, M. ve Kaya, A., Biyotelemetri Uygulamaları için Kompakt Implant Anten Tasarımı, 1. Ulusal Biyomedikal Cihaz Tasarımı-Üretimi Sempozyumu (UBİCTUS 2016), sayfa. 16-19, İstanbul, 2016.
2. Görgün, A.R. Çömlekçi, S. ve KAYA, A, Mikrodalga Tümör Ablasyonu için Hüzme Yönlülüğü Yüksek Kompakt Anten ve Kompakt çok Bandlı Düzlemsel Anten Tasarımı ve Analizi, Mühendislik Bilimleri ve Tasarım Dergisi 7(3), 662 – 670, e-ISSN: 1308-6693 Araştırma Makalesi, 2019.
3. Ferdous, N. Chin Hock, G. and Hamidah, S., Design of an Elliptical Patch Antenna for RF Energy Harvesting Application in 2.4 GHz Frequency band, International Journal of Engineering and Advanced Technology (IJEAT) ISSN: 2249 – 8958, Volume-9, Issue-2, December, 2019.

STATISTICAL CONVERGENCE ON MULTIPLICATIVE CALCULUS

Mehmet Çağrı Yılmazer[1] , Emrah Yılmaz[2] , Sertac Goktas[3] & Mikail Et[4]

[1,2,4] Department of Mathematics, University of Firat, Elazig, Turkey

[3] Department of Mathematics, University of Mersin, Mersin, Turkey

m.cagri.yilmazer@gmail.com, emrah231983@gmail.com

srtcgkts@gmail.com, mikailet68@gmail.com

Abstract

In this study, our aim is to give concepts of density, statistical convergence, statistically Cauchy sequence, Cesaro summability and λ-statistical convergence on multiplicative calculus as a special case of non-Newtonian calculus. Especially, if a generator on non-Newtonian calculus is determined as identity function, all results in the classical case are reached.

Keywords: Statistical convergence; Statistical Cauchy sequence; Cesaro convergence; Non-Newtonian calculus; Multiplicative calculus.

REFERENCES

1. Grossman, M. (1979). An introduction to non-Newtonian calculus. *International Journal of Mathematical Educational in Science and Technology*, *10*(4), 525-528.
2. Steinhaus, H. (1951). Sur la convergence ordinaire et la convergence asymptotique. *Colloquium Mathematicum*, 2(1), 73-74.
3. Stanley, D. (1999). A multiplicative calculus. *Problems, Resources, and Issues in Mathematics Undergraduate Studies*, *9*(4), 310-326.
4. Tekin, S., and Başar, F. (2013). Certain sequence spaces over the non-Newtonian complex field. *Abstract and Applied Analysis,* Article ID 739319, 1-11.
5. Zygmund, A. (2002). *Trigonometric series* (Vol. 1). Cambridge University Press.

USE OF FRACTIONAL DERIVATIVE IN DIFFERENTIAL GEOMETRY

Muhittin Evren Aydin[1], **Seyma Kaya**[2]

[1,2]Department of Mathematics, University of Firat, Elazig, Turkey

meaydin@firat.edu.tr, seymakayaa23@hotmail.com

Abstract

This talk discusses how to use the tools of fractional calculus in differential geometry of curves. Although there are many fractional derivative operator, we consider the fractional derivative of Caputo type among others because it takes a constant function to zero. Caputo fractional derivative, however, has some disadvantages where the Leibniz rule and the derivative of composite functions are given by infinite series, which these are essential for the study of differential geometry of parametrized objects. In order to overcome these troubles, we will be interested in a simplification for Caputo fractional derivative of composite functions that was first proposed in [1]. Via this simplification, we interpret the differential geometry of curves. In addition, by using the gradient in terms of Caputo fractional derivative (see [2]), we will investigate the geometry of level curves, that is, those curves given in an implicit equation.

Keywords: Caputo fractional derivative; Affine geometry; Euclidean geometry, level curves, parametrized curves.

REFERENCES

1. Yajima, T., Oiwa, S. ve Yamasaki, K., Geometry of curves with fractional-order tangent vector and Frenet-Serret formulas, Fract. Calc. Appl. Anal. Vol:21, No: 6, 1493-1505, 2018.
2. Tarasov, V.E., Fractional vector calculus and fractional Maxwell‟s equations, Annals of Physics 323, 2756-2778, 2018.

Dynamic behavior of the Fokas system arising in monomode optical fibers through the unified auxiliary equation method

Karmina K. Ali[1], Sibel Tarla[2], Resat Yilmazer[2]

[1] Department of Mathematics, University of Zakho, Zakho, Iraq

2 Department of Mathematics, University of Firat, Elazig, Turkey

karmina.ali@uoz.edu.krd

sibeltarla@gmail.com

rstyilmazer@gmail.com

Abstract

In this study, we investigate the Fokas system which represents the spread of irregular pulse in monomode optical fibers by using the unified auxiliary equation method, which is the most powerful method for exploring exact solutions for nonlinear models. The wave transform is used in order to convert a partial differential equation into an ordinary differential equation. As a result, several distinct solutions are obtained such as dark, bright, dark–bright, singular, Jacobi elliptic function, and exponential solutions. Furthermore, 3D surfaces and 2D graphics for the obtained solutions are drawn by choosing appropriate values for the involved free parameters. particular values to the parameters involved.

Keywords: Fokas system; the unified auxiliary equation method; Exact solutions.

REFERENCES

1. Tarla, S., Ali, K. K., Sun, T. C., Yilmazer, R., & Osman, M. S. (2022). Nonlinear pulse propagation for novel optical solitons modeled by Fokas system in monomode optical fibers. Results in Physics, 105381.
2. Zayed, E. M. E., & Shohib, R. M. A. (2019). Solitons and Other Solutions for Two Higher-Order Nonlinear Wave Equations of KdV Type Using the Unified Auxiliary Equation Method. Acta Physica Polonica, A., 136(1).
3. Zhang, S., & Xia, T. (2007). A generalized new auxiliary equation method and its applications to nonlinear partial differential equations. Physics Letters A, 363(5-6), 356-360.

*Abstract Submission should be prepared only **1 page**.

ARTIFICIAL INTELLIGENCE BASED DEMAND PREDICTION MODELS

Melisa BAL[1], Gizem TOPALOĞLU[1], Sevtap ERDEM[2], M. Fatih Akay[2]
[1]Trendyol, Data Science, Istanbul, Turkey
[2]Çukurova University, Department of Computer Engineering, Adana, Turkey

gizem.topaloglu@trendyol.com, melisa.bal@trendyol.com, sevtaperdem10@gmail.com, mfakay@cu.edu.tr

Abstract

Accurate demand forecasting allows companies to efficiently serve their customers' needs without investing large amounts of capital, helping to effectively lower overall operational costs. The purpose of this research is to predict the future demand of products sold in the online stores so that stocking of the product can be managed correctly. For this purpose, two different approaches were developed to accurately predict the sales of a product in six different stores in the online marketplace. The first approach utilizes deep learning based univariate time series Temporal Convolutional Networks (TCN) method whereas the second approach applies supervised machine learning based Extreme Gradient Boost (XGBoost) method to develop prediction models. In the second approach, models were created with and withouth minimum Redundancy Maximum Relevance (mRMR) feature selection algorithm. The dataset includes 5675 rows of data and includes sales counts of a product along with some other features. The sales counts in the last month of the dataset were predicted. Mean Absolute Error (MAE) was used for evaluating the performance of the prediction models. The results show that for six different stores, MAE values ranged from 6.37 to 21.89 for models developed with XGBoost, from 5.68 to 13.08 for models developed with XGBoost after mRMR and from 5.79 to 13.37 for models developed with TCN.

Keywords: : Demand forecasting, Machine learning, Deep learning

REFERENCES

1. Shilong, Z. (2021, January). Machine learning model for sales forecasting by using XGBoost. In 2021 IEEE International Conference on Consumer Electronics and Computer Engineering (ICCECE) (pp. 480-483). IEEE.
2. Mahmoudyan, M., & Zeqiri, A. (2021). Time Series Forecasting Using Neural Networks Minimizing Food Waste By Forecasting Demand in Retail Sales. Master Theses.

Diversity of wave structures to the conformable fractional dynamical model

Usman Younas, Jingle Ren

Department of Mathematics and Statistics, Zhengzhou University, China

Abstract

This manuscript examines the recently developed conformable three-dimensional Wazwaz-Benjamin-Bona-Mahony (3D-WBBM) equation's dynamical behavior in terms of its spatial and temporal variables. The governing equation is stretch for the Korteweg-de-Vries equation that represents the unidirectional propagation of small amplitude long waves on the surface of hydro magnetic and acoustic waves. Solitary wave solutions of various types, such as kink and shock, as well as singleton, combined solitons, and complex solitons, are all retrieved. Additionally, solutions to hyperbolic, exponential, and trigonometric functions are obtained through the use of recently developed methods, namely the Kudryashov method (KM), the modified Kudryashov method (MKM), and the new extended direct algebraic method (NEDAM). The study conducts a comparison of our findings to well-known findings, and concludes that the solutions reached here are novel. Additionally, the earned results are sketched in different shapes to demonstrate their dynamics as a function of parameter selection. We can assert from the obtained results that the applied techniques are simple, vibrant, and quite well, and will be helpful tool for addressing more highly nonlinear issues in various of fields.

Keywords: Integrability; soliton solutions; 3D-WBBM equation; Kudryashov and modified Kudryashov method; NEDAM.

Optical solitons of Complex Ginzburg-Landau equation having dual power nonlinear form using φ6-model expansion approach

Muhammad Abubakar Isah[1] , Asıf Yokuş [2]

[1,2]Department of Mathematics, University of Firat, Elazig, Turkey

Myphysics_09@hotmail.com[1], asfyokus@yahoo.com [2]

Abstract

The Ginzburg-Landau equation is a well-known partial differential equation in mathematics and physics, and it is applied to the research of superconducting phenomenology theory, which is widely used to describe the propagation of optical solitons over optical fibers over long distances. This paper employs a novel φ6-model expansion approach to get dark, bright, periodic, dark-bright and singular soliton solutions to the complex Ginzburg-Landau equation with dual power-law nonlinearity. The dual-power law found in photovoltaic materials is used to explain nonlinearity in the refractive index. The results of this paper may assist in comprehending some of the physical effects of various nonlinear physics models. For example, the hyperbolic sine arises in the calculation of the Roche limit and the gravitational potential of a cylinder, the hyperbolic tangent arises in the calculation of the magnetic moment and the rapidity of special relativity and the hyperbolic cotangent arises in the Langevin function for magnetic polarization. Finally, there are some discussions regarding new complex solutions. It is explored by giving physical meaning to the constants found in traveling wave solutions, which are both physically and mathematically significant. Three-dimensional simulations are used to enhance these discussions.

Keywords: φ6-model expansion approach, Complex Ginzburg-Landau equation; Jacobi elliptic functions, solitons.

REFERENCES

1. Yokus, Asif, Tukur Abdulkadir Sulaiman, and Hasan Bulut. "On the analytical and numerical solutions of the Benjamin-Bona-Mahony equation." Optical and Quantum Electronics 50.1 (2018): 31.

2. Zayed, Elsayed ME, Abdul-Ghani Al-Nowehy, and Mona EM Elshater. "New φ6-model expansion method and its applications to the resonant nonlinear Schrödinger equation with parabolic law nonlinearity." The European Physical Journal Plus 133.10 (2018): 417.

3. Arshed, Saima, et al. "Optical solitons with complex Ginzburg-Landau equation having three nonlinear forms." Physics Letters A 383.36 (2019): 126026.

MODELING OF SPECIES DISTRIBUTIONS WITH DEEP LEARNING METHOD

Serkan ÖZDEMİR[1*], Ecir Uğur KÜÇÜKSİLLE[2], Serkan GÜLSOY[3], Özdemir ŞENTÜRK[4], Mehmet Güvenç NEGİZ[1], Halil SÜEL[1], Ahmet MERT[3], Kürşad ÖZKAN[3]

[1]Isparta University of Applied Sciences, Sutculer Prof. Dr. Hasan Gürbüz Vocational School, 32950, Isparta, Turkey

[2]Suleyman Demirel University, Faculty of Engineering, 32260, Isparta, Turkey

[3]Isparta University of Applied Sciences, Faculty of Forestry, 32260, Isparta, Turkey

[4]Burdur Mehmet Akif Ersoy University, Gölhisar Vocational School, 15400, Burdur, TurkeyTurkey

serkanozdemir@isparta.edu.tr,

Abstract

There are many methods used in species distribution modeling. In recent years, the deep learning method has started to be included in these methods due to its modeling performance. In the present study, potential distribution modeling of Turkish red pine was carried out using the deep learning method with Python. In addition to having distinctive site conditions within its wide distribution, the fact that it is an ecologically and economically important species is the main reason for the Turkish red pine being the target species in this study. Considering that they may be effective in the potential distribution of the Turkish red pine, the following independent variables for modeling were chosen: elevation, topographic position index, bio1, bio8, bio12, bio14, distance from the sea, latitude and longitude. As a result of this deep learning application, it has been determined that the model is successful in terms of AUC values. According to the model map, it was observed that the determined potential areas coincided with the results from the studies conducted on the ecology of the same taxon. The results from this study showed that the deep learning method can be an effective tool for species distribution modeling.

Keywords: Deep learning, Neural network, Python, Species distribution modeling, Turkish red pine.

REFERENCES

1. Christin, S., Hervet, É., Lecomte, N. Applications for deep learning in ecology. Methods in Ecology and Evolution, 10(10), 1632–1644, 2019.

2. Keten İ., Gülsoy S. Research on site index in the red pine (Pinus brutia (Ten.)) forests. Bilge International Journal of Science and Technology Research, 4, 88–102, 2020.

THE NEW SOLUTIONS OF CONFORMABLE FRACTIONAL WHITHAM-BROER-KAUP EQUATIONS

Halil Anaç[1]

[1] Torul Vocational School, Gumushane University, Gumushane, Turkey

halilanac0638@gmail.com

Abstract

In this study, I have studied conformable q-homotopy analysis transform method for obtaining the new solutions of conformable fractional Whitham-Broer-Kaup equations. The graphs of the numerical solutions for conformable fractional Whitham-Broer-Kaup equations are plotted in MAPLE software.

Keywords: Conformable q-homotopy analysis transform method; conformable fractional Whitham-Broer-Kaup equations; conformable fractional derivative.

REFERENCES

1. Khalil R, Al Horani M, Yousef A, Sababheh M, "A new definition of fractional derivative", Journal of Computational and Applied Mathematics, 264, 65-70, 2014.
2. Abdeljawad T, On conformable fractional calculus. *Journal of computational and Applied Mathematics*, *279*, 57-66, 2015.
3. Shah R, Khan H, Baleanu D, Fractional Whitham–Broer–Kaup equations within modified analytical approaches. *Axioms*, *8*(4), 125, 2019.
4. Sherriffe D, Behera D, Nagarani P, Analytical new soliton wave solutions of the nonlinear conformable time-fractional coupled Whitham–Broer–Kaup equations. *Modern Physics Letters B*, *35*(32), 2150492, 2021.

INVESTIGATION OF OPTICAL SOLITONS TO THE NONLINEAR COMPLEX KUNDU-ECKHAUS AND ZAKHAROV-KUZNETSOV-BENJAMIN-BONA-MAHONY EQUATIONS IN CONFORMABLE

Haci Mehmet Baskonus[1] **Wei Gao**[2]

[1] Faculty of Education, Harran University, Sanliurfa, Turkey

[2] School of Information Science and Technology, Yunnan Normal University, Yunnan, China

hmbaskonus@gmail.com, gaowei@ynnu.edu.cn

Abstract This research focuses on the extraction of dark-bright solitons and periodic wave distributions of two models, namely, the Zakharov-Kuznetsov-Benjamin-Bona-Mahony equation and complex Kundu-Eckhaus equation with conformable derivative. To reach these important properties, the generalized exponential rational function method is considered. To observe wave distributions in periodic and singular sense, dynamical behaviour modulus of solutions are also archived. Strain conditions for validity of results obtained in this paper are also reported.

Keywords: Zakharov-Kuznetsov-Benjamin-Bona-Mahony equation, Nonlinear complex Kundu-Eckhaus equation with conformable, The generalized exponential rational function method,

REFERENCES

1. Khodadad FS, Nazari F, Eslami M, Rezazadeh H. Soliton solutions of the conformable fractional Zakharov–Kuznetsov equation with dual-power law nonlinearity. Optical and Quantum Electronics. 2017 Nov;49(11):1-2.
2. M.M.A. Khater, A. R. Seadawy, D. Lu, Optical soliton and rogue wave solutions of the ultra-short femto-second pulses in an optical fber via two different methods and its applications, Optik 158 (2018) 434-450.
3. B. Hu, T. Xia, N. Zhang, A Riemann–Hilbert Approach to the Kundu–Eckhaus Equation on the Half-Line, 2017 arXiv preprint arXiv:1711.02516.

Scattered Geometrical Optics Field by a Perfectly Conducting Half Plane Between Two Isorefractive Media

Mustafa Kara [1]

[1] Department of Electronics and Automation, Ordu University University TBMYO, Ordu, Türkiye

mustafa.kara@odu.edu.tr,

Abstract

Scattered geometrical optics field of plane electromagnetic waves by a perfectly conducting half plane between two isorefractive media is derived by means of the method of transition boundary. The interface between the two media is considered as a resistive surface and affects the reflection and transmission coefficients of the half plane, and it has both reflection and transmission coefficients. Initial field is determined first by excluding the half plane causing diffraction from the geometry. Secondly total geometrical optics field is determined. Finally, scattered geometrical optics field is obtained by subtracting the initial field from the total geometrical optics field.

Keywords: Scattered geometrical optics; Conducting half plane; isorefractive media

REFERENCES

1. J.S. Tyo, A Class of Artificial Materials Isorefractive With Free Space, IEEE Trans. Antennas Propag., Vol:51, No:5, 1093–1099, 2003.
2. J.S. Tyo, C.J. Buchenauer, J.S.H. Schoenberg, Use of Isorefractive Media to Improve Prompt Aperture Efficiency in a Lens IRA, IEEE Trans. AntennasPropag., Vol:46, No:7, 1114–1115, 1998.
3. P.L.E. Uslenghi, Scattering by a Half-Plane at the Interface Between Isorefractive Media, IEEE Antennas Wirel. Propag. Lett., Vol:12, 1662–1664, 2013.
4. P.L.E. Uslenghi, Radiation of a Line Source Over a Half-Plane Located at the Interface Between Isorefractive Media, IEEE Antennas Wirel. Propag. Lett., Vol:15, 356–357, 2016.
5. Y.Z. Umul, Scattering by a conductive half-screen between isorefractive media, Appl. Opt., Vol:54, 10309–10313, 2015.
6. Y.Z. Umul, Scattering of waves by a soft-hard half-plane between isorefractive medium, J. Mod. Opt., Vol:64, No:4, 329–333, 2016.
7. Y.Z. Umul, Diffraction of line source field by a resistive half-plane between isorefractive media, Opt. Quant. Electron., Vol:49, No:1, 23, 2017.
8. H.D. Basdemir, Scattering of waves by a perfectly conducting wedge residing at the interface between two isorefractive media, IEEE Antennas Wirel.Propag. Lett., Vol:15, 1361–1364, 2016.
9. H.D. Basdemir, Scattering of evanescent plane waves by a black wedge at the interface between two isorefractive media, IEEE Trans. AntennasPropag., Vol:64, No:8, 3661–36642016.

ANALYSIS OF TRAVELING WAVE SOLUTIONS OF NONLINEAR PARTIAL DIFFERENTIAL EQUATIONS

Tolga AKTÜRK[1] , MahşureKübranur DİKİCİ[2]

[1]Department of Mathematics and Science Education, Faculty of Education, Ordu University, Turkey

[2]Department of Mathematics, Faculty of Science, Ordu University, Turkey

[1]tolgaakturkk@gmail.com,[2]kubradikici.55@hotmail.com;

ABSTRACT

In this article, traveling wave solutions of the nonlinear partial differential equations have been analyzed using the modified exponential function method (MEFM). This method was applied to analytical traveling wave solutions of the Calogero Bogoyavlenskii Schiff (CBS) and Whitham Broer Kaup (WBK) equations. Solution functions were obtained through a mathematical program. Two, three- dimensional and contour graphs expressing the behavior of solution functions representing the mathematical model were plotted with the help of the Mathematica package program by determining the appropriate parameters.

Keywords: Calogero Bogoyavlenskii Schiff (CBS) and Whitham Broer Kaup (WBK) Equaiton, Nonlinear Partial Differential Equations, Modified Exponential Function Method.

REFERENCES

1. Baskonus, H. M., Bulut, H. ve Süleyman, T. A. (2017). Investigation of various travelling wave solutions to the extended (2+1)-dimensional quantum ZK equation. The European Physical Journal Plus, 132(11), 482.
2. Bulut, H., Aktürk, T. ve Yel, G. (2018). An Application of the Modified Expansion Method to Nonlinear Partial Differential Equation. *Turk. J. Math.* Comput. Sci., 10,202–206.
3. Gurefe, Y., Aktürk, T. ve Günaydın, E. (2018). On the kink type and singular solitions solitions to the nonlinear partial differential equation. International Conference on Pure and Applied Mathematics (ICPAM-2018), 11-13 Eylül, Van yüzüncüyıl üniversitesi, Van.
4. Xu, G., & Li, Z. (2005). Exact travelling wave solutions of the Whitham–Broer–Kaup and Broer–Kaup–Kupershmidt equations. *Chaos, Solitons & Fractals*, *24*(2), 549-556.
5. Wazwaz, A. M. (2008). Multiple-soliton solutions for the Calogero–Bogoyavlenskii–Schiff, Jimbo–Miwa and YTSF equations. *Applied Mathematics and Computation*, *203*(2), 592-597.
6. Naher, H. ve Abdullah, F. A., 2013. New approach of (G′/G)-expansion method and new approach of generalized (G′/G)-expansion method for nonlinear evolution equation. American Inst of Physics Advances, 3(3),032116.
7. Yel, G. ve Aktürk, T. (2019). Modified Expansion Function Method for the KP-BBM Equation. CMES 2019, 2-3.

SOME CONVERGENCE RESULTS FOR A NEW GENERALIZED NONEXPANSIVE MAPPINGS IN UNIFORMLY CONVEX HYPERBOLIC SPACE

Nazlı Karaca [1]

[1] Department of Mathematics, Ataturk University, Erzurum, Turkey

nazli.kadioglu@atauni.edu.tr,

Abstract

In this paper, we introduce a new generalized class of nonexpansive mappings. Also we establish some strong and Δ-convergence theorems of an iteration process for approximating fixed point of these mappings in a uniformly convex hyperbolic space

Keywords: Fixed point; Generalized nonexpansive mappings; Uniformly convex hyperbolic space.

REFERENCES

1. Adeyemi, Tijesunimi Abraham, et al. The Existence of a Solution of the Nonlinear Integral Equation Via The Fixed Point Approach, Advances in Mathematics: Scientific Journal, 10, no.7, 2977–2998, 2021.
2. Sahin, Aynur; Basarir, Metin, Some convergence results for nonexpansive mappings in uniformly convex hyperbolic spaces, Creat. Math. Inform, 26, No. 3,331 – 338, 2017.
3. W. A. Kirk, B. Panyanak: A concept of convergence in geodesic spaces, Nonlinear Anal. Theory Methods Appl., Ser. A, Theory Methods, 68 (12), 3689–3696, 2008.
4. Khan, S. H. and Abbas, M., Strong and Δ-convergence of some iterative schemes in CAT(0) spaces, Comput. Math. Appl., 61, No. 1, 109–116, 2011.
5. Kohlenbach, U., Some logical metatheorems with applications in functional analysis, Trans. Am. Math. Soc., 357, No. 1, 89–128, 2005.

A FIXED POINT THEOREM IN B-METRIC SPACES AND AN APPLICATION TO SPECTRAL ASYMPTOTICS OF SOLUTIONS OF OPERATOR EQUATIONS

Ayşe GÜVEN SARIHAN [1], Oscar F. BANDTLOW [2]

[1] Department of Mathematics, Ordu University , Ordu, Turkey

[2] School of Mathematical Sciences, Queen Mary University of London, London, UK

ayseguvensarihan@odu.edu.tr
o.bandtlow@qmul.ac.uk

Abstract

In this paper, we prove a fixed point theorem for eventually contracting maps on b-metric spaces and use it to study eigenvalue bounds for solutions of operator equations in quasi-Banach operator ideals.

Keywords: Fixed point; B-metric space; Quasi-Banach operator ideals.

REFERENCES

1. Faezeh Bazazi, Kourosh Nourouzi, Donal O'Regan, Two order theoretic proofs for a fixed point theorem in partially ordered metric spaces and its application to trace class operator equations, Fixed Point Theory, Vol:15, No:2, 393–398, 2014.
2. Ayşe Güven Sarıhan, Oscar F. Bandtlow. Quantitative spectral perturbation theory for compact operators on a Hilbert space, Linear Algebra Appl., 610, 169–202, 2021.
3. A. C. M. Ran, M. C. B. Reurings, L. Rodman, A perturbation analysis for nonlinear selfadjoint operator equations, SIAM J. Matrix Anal. Appl., Vol:28, No:1, 89–104, 2006.
4. Ştefan Cobzaş, B-metric spaces, fixed points and Lipschitz functions, 2018.

A TURBULENCE MODEL WITH SLIP WITH FRICTION BOUNDARY CONDITION

Özgül İLHAN[1]

[1] Department of Mathematics, University of Muğla Sıtkı Koçman, Muğla, Turkey

oilhan@mu.edu.tr,

Abstract

In this study, the slip with friction boundary condition is applied to Navier-Stokes-ω model [1]. Numerical tests are performed using this boundary condition on two-dimensional channel flows across a step and on the bottom wall. The influence of the friction parameters are investigated for different Reynolds numbers. For this reason, the reattachment points are calculated and compared with the reference solutions for different Reynolds numbers and friction coefficients. The reference solutions are obtained with Navier-Stokes equation with no-slip boundary conditions on fine grid [2]. Finally, with the help of the results obtained the effect of the friction coefficient on the flow is explained.

Keywords: Navier-Stokes-ω model, turbulence model, reattachment point, slip with friction boundary condition.

REFERENCES

1. Hill, G.R., (2010). \emph{Benchmark Testing the α-models of Turbulence.}, Master of Science, the Graduate School of Clemson University.
2. John, V., Layton, W.J. and Sahin, N., "Derivation and analysis of near wall models for channel and recirculating flows.", Comput. Math. Appl., 28: 1135-1151, 2004.

A NEW GENERALIZATION OF THE APOSTOL-EULER AND APOSTOL-GENOCCHI POLYNOMIALS

Zeynep Özat[1], **Bayram Çekim**[1]

[1] Department of Mathematics, University of Gazi, Ankara, Turkey

zeynepozat95@gmail.com, bayramcekim@gazi.edu.tr

Abstract

In the present study, we introduce new type Apostol-Euler polynomials and Apostol-Genocchi polynomials with three variables via generating functions using the Euler formula and operational methods. By means of these generating functions, some identities, multiplicative and derivative operators are obtained. We also examine Gould-Hopper-Apostol-Euler and Genocchi polynomials, and Hermite-Appell-Apostol-Euler and Genocchi polynomials by considering special cases.

Keywords: Apostol-Euler polynomials and numbers; Apostol-Genocchi polynomials and numbers; Generating function.

REFERENCES

1. Hari Mohan Srivastava, Rekha Srivastava, Abdulghani Muhyi, Ghazala Yasmin, Hibah Islahi, Serkan Aracı, Construction of a new family of Fubini-type polynomials and its applications, Advances in Difference Equations, Vol:36, 1-25, 2021.
2. Hari Mohan Srivastava, Mohammad Masjed-Jamei, Mohammad Reza Beyki, A parametric type of the Apostol–Bernoulli, Apostol–Euler and Apostol–Genocchi polynomials, Appl. Math. Inf. Sci. Vol:12, No:5, 907—916, 2018.
3. Hari Mohan Srivastava, Can Kızılateş, A parametric kind of the Fubini-type polynomials. RACSAM, 113, 3253–3267, 2019.

GENERALIZED OF HERMITE MATRIX POLYNOMIALS AND SOME PROPERTIES

Neslihan BİRİCİK[1], **Bayram ÇEKİM**[1]

[1]Department of Mathematics, University of Gazi, Ankara, Turkey

neslihanbiricik@gazi.edu.tr, bayramcekim@gazi.edu.tr

Abstract

In this study, we introduce generalized Hermite matrix polynomials defined with the help of generalized exponential function. For this family of polynomials, some properties such as recurrence relation, inversion formula are obtained. In addition, the relationship of generalized Hermite matrix polynomials with the Beta function is emphasized.

Keywords: Hermite matrix polynomials; Dunkl generalization; Recurrence relation.

REFERENCES

1. Gabor Szego, Orthogonal Polynomials, New York: American Mathematical Society Colloquium Publications, 1-440, 1939.
2. Theodore S. Chihara, An Introduction to Orthogonal Polynomials, New York: Gordon and Breach, 43, 157, 1978.
3. Marvin Rosenblum, Generalized Hermite polynomials and the Bose-like oscillator calculus, In Nonselfadjoint Operators and Related Topics, Operator Theory: Advances and Applications, Vol: 73, 369-396, 1994.
4. Lucas Jodar and Rafael Company, Hermite matrix polynomials and second order matrix differential equations, Journal of Approximation Theory and Applied, Vol:12, No:2, 20–30, 1996.

On The Periodic Solutions of a Nonlinear Differential Equation with Variable Delays

Harun Biçer[1]

[1] Department of Mathematics, University of Bingol, Bingol, Turkey

hbicer23@gmail.com,

Abstract

We obtain new results for existense of periodic solutions of a nonlinear differential equation with variable delays. Our findings have a contribution to the topic and complete that in the relevant literature.

Keywords: Nonlinear differential equation; periodic solutions.

REFERENCES

1. L. Wang, J. Shao, New results of periodic solutions for a kind of forced Rayleigh-type equations, Nonlinear Anal. Real World Appl. 11, 99–105, 2010.
2. B. Liu, C. Tunc, Pseudo almost periodic solutions for a class of first order differential iterative equations. Appl. Math. Lett. 40 29-34, 2015.

A UNIFORM NUMERICAL APPROACH FOR SINGULARLY PERTURBED FREDHOLM INTEGRO-DIFFERENTIAL EQUATION

Muhammet Enes Durmaz[1*], **Gabil Amirali**[2]

[1] Department of Information Technology, Kırklareli University, Kırklareli, Turkey

[2] Department of Mathematics, Faculty of Art and Science, Erzincan Binali Yıldırım University, Erzincan, Turkey

menesdurmaz025@gmail.com, gabilamirali@yahoo.com

Abstract

This article deals with a finite difference method for solving a singularly perturbed Fredholm integro-differential equation. For the numerical solution of the problem, we use an exponentially fitted difference scheme on a uniform mesh which is succeeded by the method of integral identities with the use of exponential basis functions and interpolating quadrature rules with the weight and remainder terms in integral form. The method is shown to be first-order convergent in the discrete maximum norm. Parameter uniform error estimates for the approximated solutions are established. The numerical example validates the theoretical results.

Keywords: Finite difference scheme; Fredholm integro-differential equation; Singular perturbation; Uniform convergence.

REFERENCES

1. Roos HG, Stynes M, Tobiska L., Numerical methods for singularly perturbed differential equations convection-diffusion and flow problems, Berlin: Springer-Verlag; 2014.

2. Amiraliyev GM, Durmaz ME, Kudu M., A numerical method for a second order singularly perturbed Fredholm integro-differential equation, Miskolc Mathematical Notes 2021; 22 (1): 37-48.

The examination of the quotient of numerical semigroup with RF-matrices

Belgin Özer[1], Fatima Dakkak[2]

[1]Department of Mathematics, University of Gaziantep, Gaziantep, Turkey

emirhan@gantep.edu.tr

[2] Department of Mathematics, University of Gaziantep, Gaziantep, Turkey

fatimadakkak94@gmail.com

Abstract

In this paper, we study quotients of a numerical semigroups with RF (Row-Factorization) matrices. We prove a formula for the Frobenius number of quotients of some families of numerical semigroups. Moreover, we examine half of the numerical semigroups, pseudo-symmetric numerical semigroups.

Keywords: *Quotient of A Numerical Semigroup; Pseudo-Frobenious Number; RF (Row Factorization) Matrices.*

REFERENCES

1.J. C. Rosales, J. M. Urbano-Blanco, Proportionally modular Diophantine inequalities and full semigroups, Semigroup Forum 72 (2006), 362-374 https://doi.org/10.1007/s00233-005-0527-8.
2. A. M. Robles-P´erez, J. C. Rosales, Equivalent proportionally modular Diophantine inequalities, Archiv der Mat
3. J.C. Rosales, P. A. Garcia-Sanchez, Every numerical semigroup is one half of infinitely many symmetic numerical semigroups, to appear in Comm. Algebra.
4. J. C. Rosales, One half of a pseudo-symmetric numerical semigroup, preprint.
5.J.C. Rosales, P. A. Garcia-Sanchez, J.I.Garcia-Garcia, J. M. Urbano-Blanco, proportionally modular Diophantine inequalities,J.Number Theory 103 (2000),281-294. https://doi.org/10.1016/j.jnt.2003.06.002.
6. J.C. Rosales, P. A. Garcia-Sanchez, Every numerical semigroup is one half of infinitely many symmetic numerical semigroups, Commun .Algebra 36 (2008) , 2910-2916. https://doi.org/10.1080/00927870802108171.
7.J.C. Rosales, P. A. Garcia-Sanchez, Every numerical semigroup is one half of a symmetic numerical semigroups, Proc.Amer.Math.Soc.136 (2008) , no.2,477-457. https://doi.org/10.1090/S0002-9939-07-09098-3.
8. A. Moscariello , On the type of an almost gorenstein monomial curve .Journal of Algebra 456,266-277 (2016) . https://doi.org/10.1016/j.jalgebra.2016.02.019.
9.J.C. Rosales, P. A. Garcia-Sanchez, Numerical Semigroup ,Springer Developments in Mathematies ,Volume 20,(2009).

ON SOLUTIONS OF INTEGRAL EQUATIONS IN GEOMETRIC ANALYSIS

Arzu Bal[1], Numan Yalçın[2] and Mutlu Dedetürk[1]

[1] Department of Mathematical Engineering, University of Gumushane, Gümüşhane, Turkey

arzucavusculubal@gmail.com,

mutludedeturk@gumushane.edu.tr,

[2]Gumushane Vocational School, Department of Electronics and Automation, University of Gumushane, Gümüşhane, Turkey

numan@gumushane.edu.tr,

Abstract

In this paper, we will give definitions of types of integral equations on geometric analysis. And solutions of different types of integral equations in geometric analysis will be examined. These studies will be reinforced with examples.

Keywords: Multiplicative integral equation; Multiplicative Fredholm integral equation; Multiplicative Volterra integral equation; Multiplicative Volterra-Fredholm integral equation.

REFERENCES

1. Fredholm, I. (1903). Sur une classe d'équations fonctionnelles. *Acta mathematica*, *27*, 365-390.
2. Volterra, V., & Hostinsky, B. Opérations Infinitésimales linéares, Herman, 1938.
3. Lovitt, W.V.: Linear Integral Equations. Dover Publications Inc.,New York, 1950.
4. Smithies, F.: Integral Equations. Cambridge University Press, London, 1958.
5. Krasnov, M., Kiselev, K., Makarenko, G.: Problems and Exercises in Integral Equation. Mır Publishers, Moscow, 1971.
6. Kiselev, A., Makeronko, G., Krasnov, M., (Cevdet Cerit trans.), Problems and Exercises in Integral Equations, Alfa Yayınları, İstanbul, 1976.
7. Delves, L.V., Mohamed, J.L.: Computational Methods for Integral Equations. Cambridge University Press, New York, 1985.
8. Upadhyay, S., Integral equation. 10.13140/RG.2.1.2854.8569, 2015.

ON SOLUTIONS OF FIRST AND SECOND ORDER LINEAR DIFFERENTIAL EQUATIONS IN GEOMETRIC ANALYSIS

Numan Yalçın [1] and Mutlu Dedetürk[2]

[1] Gumushane Vocational School, Department of Electronics and Automation, University of Gumushane, Gümüşhane, Turkey

numan@gumushane.edu.tr,

[1] Department of Mathematical Engineering, University of Gumushane, Gümüşhane, Turkey

mutludedeturk@gumushane.edu.tr,

Abstract

In this work, the solutions of first and second order linear differential equations in geometric analysis will be focused on. For this, multiplicative power series will be used. These studies will be reinforced with examples.

Keywords: Geometric analysis; Linear differential equation; Multiplicative power series.

REFERENCES

1. Aniszewska, D.: Multiplicative Runge-Kutta method, Nonlinear Dynamics 50, 265–272 (2007) Volterra, V., & Hostinsky, B. Opérations Infinitésimales linéares, Herman, 1938.
2. Bashirov, A. E., Kurpinar, E. M., Ozyapici, A.: Multiplicative calculus and its applications, J. Math. Anal. Appl. 337, 36–48 (2008)
3. Bashirov, A.E., Misirli, E., Tandogdu, Y., Ozyapici, A.: On modeling with multiplicative differential equations, Applied Mathematics-A Journal of Chinese Universities 26:4, 425–438 (2011).
4. Bhat, A.H., Majid, J., Shah, T.R., Wani, I.A., Jain, R.: Multiplicative Fourier Transform and Its Applications to Multiplicative Differential Equations, Journal of Computer and Mathematical Sciences, 10(2),375–383 (2019)
5. Grossman, M., Katz, R.: Non-Newtonian calculus Lee Press, Pigeon Cove, MA, (1972)
6. Stanley, D.: A multiplicative calculus, PRIMUS 9:4, 310–326 (1999)

SOME PROPERTIES OF THE PARAMETRIC KIND OF FUBINI POLYNOMIALS WITH THREE VARIABLES OF ORDER r

Kübra Çetin [1] **, Bayram Çekim** [1]

[1] Department of Mathematics, University of Gazi, Ankara, Turkey

ccetinkubraa@gmail.com, bayramcekim@gazi.edu.tr

Abstract

In this study, we introduce Fubini polynomials with three variables of order r with the help of generating functions. We derive addition formulas, summation formulas, trigonometric relation, differential formulas. Also we prove some relation between our polynomials and Stirling numbers.

Keywords: Fubini polynomials; Summation formulas; Trigonometric relation.

REFERENCES

1. Dae San Kim, Taekyun Kim, Hyuck-In Kwon, Jın- Woo Park, Two variable higher order Fubini polynomials, Journal of the Korean Mathematical Society, Vol:55, No:4, 975–986, 2018.
2. Sunil Kumar Sharma, Waseem A. Khan, Cheon Seoung Ryoo, A parametric kind of Fubini polynomials of a complex variable, Mathematics, Vol:8, No:4, 643, 2020.
3. Stephen M. Tanny, On some numbers related to the Bell numbers, Canadian. Mathematical Bulletin, Vol:17, No:5, 733-738, 1975.

ON A NEW TYPE OPERATORS INCLUDING THE APPELL POLYNOMIALS AND ITS APPROXIMATION PROPERTIES

Bilge Zehra Sergi, Gürhan İçöz

Department of Mathematics, University of Gazi, Ankara, Turkey

bilgezsergi@gazi.edu.tr, gurhanicoz@gazi.edu.tr

Abstract

The purpose of this paper is to define a new sequence of linear positive operators including the Appell polynomials. We have convergence properties of this operators such as the Korovkin's theorem, modulus of continuity, Lipschitz class and Voronovskaja-type theorem. Then, a Kantorovich form of this operators and its some approximation properties are given.

Keywords: Appell polynomials; Szàsz operators; Voronovskaja-type theorem.

REFERENCES

1. Gürhan İçöz, Serhan Varma, Sezgin Sucu, Approximation by operators including generalized Appell polynomials, Filomat, Vol:2, No:30, 429–440, 2016.
2. Ammon Jakimovski, Dany Leviatan, Generalized Szàsz operators for the approximation in the infinite interval, Mathematica, Vol:34, No:11, 97–103, 1969.
3. Mehmet Ali Özarslan, Banu Yaşar, Δ_h -Gould Hopper Appell polynomials, Acta Mathematica Scientica, Vol:4, No:41B, 1196–1222, 2021.
4. Sezgin Sucu, Serhan Varma, Approximation by sequence of operators involving analytic functions, Mathematics, No:7, 188, 2019.

THE NUMERICAL SOLUTION OF SINGULARLY PERTURBED DELAY INTEGRO-DIFFERENTIAL EQUATION

Hasan Bulut[1], **Derya Arslan**[2]

[1] Department of Mathematics, University of Firat, Elazig, Turkey

[2]Department of Mathematics, University of Bitlis Eren, Bitlis, Turkey

[1]hbulut@firat.edu.tr,

[2]ayredlanu@gmail.com.,

Abstract

In this study, we use differential transform method to numerical solve delay integro-differential equations. Both theory and application of integro-differential equation are a very important in applied mathematics. These equations are used to mathematical models for many physical situations. The differential transformation method is an iterative procedure to obtain serial solutions of ordinary and partial differential equations. For this reason, it is used to solve the delay integro-differential equations in the study. The reliability, effectively, and simplicity of the proposed method are proved by the results of applying the method on thesingularly perturbed delay integro-differential equations.

REFERENCES

1. Zhou, J.K., Differential Transformation and Its Applications for Electrical Circuits. Huazhong University Press, Wuhan, 1986.
2. Ayaz, F., Application of Differential Transform Method to Differential-Algebraic Equations, Appl. Math. Comput., Vol:152, 649-657, 2004.
3. Arikoglu, A. and Ozkol, I., Solution of Boundary Value Problems for Integro-Differential Equations by Using Differential Transform Method, Applied Mathematics and Computation, Vol:168, 1145-1158, 2005.
4. Biazar, J. and Eslami, M., Differential Transform Method for Systems of Volterra Integral Equations of the Second Kind and Comparison with Homotopy Perturbation Method, International Journal of Physical Sciences, Vol:6, 1207-1212, 2011.

Fine Grinding of Marble Waste: Preliminary Test Results

Çimen Gül Kuluşaklı[1], Mustafa Birinci[2]*,

[1]Graduate School of Natural and Applied Sciences, İnönü University, Malatya, Turkey

[2]Department of Mining Engineering, İnönü University, Malatya, Turkey

mustafa.birinci@inonu.edu.tr, cimenaras@gmail.com

Abstract

The aim of this study was to investigate the production of ultrafine material from marble wastes. For this purpose, marble wastes were ground to fine sizes using planetary ball mill with porcelain milling media. The milling tests were performed at a constant mill speed of 150 rpm with three different ball to powder ratios (2.5, 5 and 10) and variable grinding times up to 240 min. The particle size analyses showed that the mean particle size (d_{50}) of the milled products was significantly reduced compared to the feed size. Finally, the preliminary test results from planetary milling indicated that marble wastes can be ground to the desired fine particle sizes for industrial utilization.

Keywords: Marble waste; Ultrafine grinding; Planetary ball mill.

ON THE SEMI-ANALYTICAL METHODS FOR THE FIFTH ORDER NONLINEAR KdV EQUATIONS

İnci Çilingir Süngü [1] **Emre Aydın** [2]

[1] Department of Mathematics Education, University of Ondokuz Mayis, Samsun, Turkey

incicilingir@gmail.com,

[2]Department of Mathematics, University of Ondokuz Mayis, Samsun, Turkey,
emr_aydn_55@outlook.com

Abstract

In this study, the Caudrey-Dodd-Gibbon (CDG), Sawada-Kotera (SK) and Caudrey-Dodd-Gibbon-Sawada-Kotera (CDGSK) equations being from the class of fifth-order nonlinear KdV equations are investigated. Adomian Decomposition Method (ADM) and modified Variational Iteration Method (mVIM) are proposed semi-analytical solutions for the CDG, SK and CDGSK equations. ADM and mVIM solutions of these equations are compared with both their exact solutions and each other by using tables and graphs. Also, it is visualized using curve and surface graphs. It is shown that the semi-analytical solutions are calculable, convenient and highly compatible with the analytical solutions.

Keywords: Caudrey-Dodd-Gibbon (CDG) equations; Sawada-Kotera (SK) equations; Caudrey-Dodd-Gibbon-Sawada-Kotera (CDGSK) equations; Adomian decomposition method (ADM); modified variational iteration method (mVIM).

REFERENCES

1. Hasan Bulut, H. Mehmet Baskonus, Seyma Tuluce, Tolga Akturk, A Comparison Between HPM and ADM for the Nonlinear Benjamin-Bona-Mahony Equation, International Journal of Basic & Applied Sciences, Vol:11, No:3, 117-127, 2011.
2. Tamer A. Abassy, Modified Variational Iteration Method (Nonlinear Homogeneous Initial Value Problem), Computers and Mathematics with Applications, Vol: 59, No: 2, 912-918, 2010.
3. Yongyi Gu, Analytical Solutions to the Caudrey-Dodd-Gibbon-Sawada-Kotera Equation via Symbol Calculation Approach, Journal of Function Spaces, Vol:2020, article id: 5042724, 8 pages, 2020.
4. Hasibun Naher, Farah Aini Abdullah, M. Ali Akbar, The (G'/G)- Expansion Method for Abundant Travelling Wave Solutions of Caudrey-Dodd-Gibbon Equation, Mathematical Problems in Engineering, Vol: 2011, article id: 218216, 11 pages, 2011.

A NEW CONSTRUCTION METHOD FOR TRIANGULAR NORMS AND TRIANGULAR CONORMS ON BOUNDED LATTICES

Gül Deniz Çaylı

Department of Mathematics, Karadeniz Technical University, Trabzon, Turkey

guldeniz.cayli@ktu.edu.tr

Abstract

A triangular norm T (resp. triangular conorm S) is a binary operation on a bounded lattice L such that it is commutative, associative, increasing in each variable and satisfies the boundary condition $T(1,x)=x$ (resp. $S(0,x)=x$) for all $x \in L$. A triangular subnorm F (resp. triangular superconorm Q) is obtained by changing the boundary condition of triangular norm (resp. triangular conorm) into the condition that $F(x,y) \leq \inf\{x,y\}$ (resp. $Q(x,y) \geq \sup\{x,y\}$) for all $x,y \in L$. In this paper, we demonstrate that it is possible to describe a construction method for triangular norms and triangular conorms on a bounded lattice L derived from triangular subnorms and triangular superconorms defined on the subinterval of L, respectively.

Keywords: Triangular norm; Triangular conorm; Triangular subnorm; Triangular superconorm.

REFERENCES

1. Garrett Birkhoff, Lattice Theory, American Mathematical Society Colloquium Publishers, Providence, RI, 1967.
2. Gül Deniz Çaylı, On a new class of t-norms and t-conorms on bounded lattices, Fuzzy Sets and Systems, Vol:332, 129-143, 2018.
3. Jesus Medina, Characterizing when an ordinal sum of triangular norms is a triangular norm on bounded lattices, Fuzzy Sets and Systems, Vol:202, 75-88, 2012.
4. Susanne Saminger, On ordinal sums of triangular norms on bounded lattices, Fuzzy Sets and Systems, Vol:157, 1403-1416, 2006.
5. Berthold Schweizer, Abe Sklar, Statistical metric spaces, Pacific Journal of Mathematics, Vol:10, 313-334, 1960.

INVESTIGATION OF THE EFFECT OF VACCINATION IN TURKEY ON PNEUMONIA CASES CAUSED BY COVID-19 WITH A SEIR TYPE MODEL

Fatma Karaca Vural, İ. Onur Kıymaz

Department of Mathematics, University of Kırşehir Ahi Evran, Kırşehir, Turkey

fatmakaraca_1987@hotmail.com , iokiymaz@ahievran.edu.tr

Abstract

In this study, a SEIR type compartmental disease model was developed to investigate the impact of vaccination on the evolution of disease to pneumonia during the COVID-19 epidemic. The stability of the developed model and the basic reproduction number was also calculated. To arrive at numerical solutions of the model two periods of 63 days, the unvaccinated-first vaccinated and the first vaccinated-second vaccinated period, were taken as basis. Data for these periods were obtained from the website of TR Ministry of Health. The graphs of the numerical solutions were drawn and their compatibility with the real data was shown.

Keywords: COVID-19; SEIR epidemic model; local stability; basic reproduction number; numerical solution.

REFERENCES

1. Brauer, F., Van den Driessche, P., Wu J., (Eds), (2008), Mathematical Epidemiology, Springer-Verlag, Germany .

2. Diekmann, O., Heesterbeek, J. A. P., Metz, J.A.J.,(1990), On the definition and thecomputation of the basic reproduction ratio R0 in models for infectious diseases inheterogeneous populations, J. Math. Biol. 28,365.

3. Allen, L. J. S., An Introduction to Stochastic Epidemic Models, Department of Mathematics and Statistics, Texas Tech University, Lubbock, TX 79409-1042, USA.

4. Qureshi, S., Yusuf, A., (2019), Fractional derivatives applied to mseir problems: comparative study with real world data, Eur. Phys. J. Plus, 134 (4), p. 171.

NON-CANAL CIRCULAR SURFACES GENERATED BY SPECIAL CURVES WITH ALTERNATIVE FRAME AND QUATERNION

Abdussamet Çalışkan[1]

[1]Fatsa Vocational School, Accounting and Tax Applications, University of Ordu, Ordu, Turkey

abdussamet65@gmail. com,

Abstract

In this paper, we show that non-canal circular surfaces can be obtained by Alternative frames of the special curves and obtain some corollaries. Besides, these surfaces are obtained by quaternion and homothetic motion.

Keywords: Non- Canal Circular Surface; Alternative Frame; Bertrand curve; Involute-Evolute curve; Quaternion; Rotation matrices; Homothetic motion.

REFERENCES

1. B. Uzunoğlu, İ. Gök, ,Y. Yaylı, A new Approach on Curves of Constant Precession. Applied Mathematics and Computation, Vol:275, 317-323, 2016.
2. E. Li, D. Pei, Involute-evolute and pedal contrapedal curve pairs on S^2, Mathematical Methods in the Applied Sciences, 1-15, 2020.
3. İ. Gök, Quaternionic approach of Canal Surfaces constructed by some new ideas, Advances in Applied Clifford Algebras, Vol. 27, No. 2, 1175-1190, 2017.
4. S. Aslan, Y. Yaylı, Split Quaternions and Canal Surfaces in Minkowski 3-Space, International Journal of Geometry, Vol. 5, No. 2, 51-61, 2016.
5. Z. Xu, R. Feng, J.G. Sun, Analytic and algebraic Properties of Canal Surfaces, Journal of Computational and Applied Mathematics, Vol. 195, No. 1-2, 220-228, 2006.

SYMMETRIC DYNAMIC DERIVATIVES FOR QUAD REMESHING OF SIMPLICIAL SURFACES

Mehmet Ali Balcı[1] **, Ömer Akgüller**[1], **Sibel Paşalı Atmaca**[1]

[1] Department of Mathematics, Mugla Sitki Kocman University, Mugla, Turkey

mehmetalibalci@mu.edu.tr, oakguller@mu.edu.tr, sibela@mu.edu.tr

Abstract

In this study, we introduce concepts from time scale calculus emerging discrete differential geometry and an algorithm for quadrilateral surface parametrization. This algorithm converts simplicial surfaces to a global quad-parametrization. The discrete integrability of time scale operators defined in symmetric sense may serve an example to increase reliability along the geometry processing.

Keywords: Time scale calculus, Discrete normal fields, geometry processing

REFERENCES

1. Paşalı Atmaca, S., Akgüller, Ö.. Curvature of curves parameterized by a time scale. Advances in Difference Equations, Vol 2015, No 1, 1-7., 2015

2. Akgüller, Ö., Atmaca, S. P. Discrete normal vector field approximation via time scale calculus. Applied Mathematics and Nonlinear Sciences, Vol 5, No 1, 349-360, 2020

DISTRIBUTION OF DISCRETE GEODESICS ON ALPHA COMPLEXES

Ömer Akgüller[1] **, Mehmet Ali Balcı**[1]

[1] Department of Mathematics, Mugla Sitki Kocman University, Mugla, Turkey

oakguller@mu.edu.tr, mehmetalibalci@mu.edu.tr

Abstract

In this study, the shortest paths calculated on the graph structures expressed by alpha complex skeleton of the submanifold from which the point cloud is sampled will provide an approximation to the geodetic curves of the manifold. It should also be taken into account that this sampling will contain noise. Although the graph structures formed by the simplex complex skeleton are affected by this noise, the approaches to the geodesics of the submanifold are not affected much by the distributions of the geodesics. Thus, a kernel function to be defined by the Wasserstein similarity of the distributions of discrete geodesics in the skeletons of 3D point clouds will be an effective tool in the point cloud similarity measurement process.

Keywords: Discrete geodesics; Alpha Complex; Point cloud process.

REFERENCES

1. Li, Xin, Hsinchun Chen. Recommendation as link prediction in bipartite graphs: A graph kernel-based machine learning approach. Decision Support Systems, Vol 54, No 2, 880-890, 2013.

2. Ma, T., Shao, W., Hao, Y., Cao, J. Graph classification based on graph set reconstruction and graph kernel feature reduction. Neurocomputing, Vol 296, 33-45, 2018

3. Boulch, A. ConvPoint: Continuous convolutions for point cloud processing. Computers & Graphics, Vol 88, 24-34, 2020

CLASSIFICATION OF CAMERA TRAP DATA USING CONVOLUTIONAL NEURAL NETWORKS

Halil SÜEL[1], Mahmut TOKMAK[2], Akın KIRAÇ[3]

[1] Sütçüler Prof.Dr.Hasan Gürbüz Vocational School, Isparta University of Applied Sciences, Isparta, Turkey halilsuel@isparta.edu.tr,

[2]Bucak Zeliha Tolunay Applied Technology and Business Administration School, Burdur Mehmet Akif Ersoy University, Burdur Turkey mahmuttokmak@mehmetakif.edu.tr

[3]Çanakkale Technical Science Vocational School, Çanakkale Onsekiz Mart University, Çanakkale, Turkey akinkirac@comu.edu.tr

Abstract

Camera trap data which is a lot of and big data is considered to be important an inventory method. In this study; In order to develop an alternative method, 353 camera trap data including Brown bear (*Ursus arctos*), Eurasian lynx (*Lynx lynx*), Gray wolf (*Canis lupus*), European hare (*Lepus europaeus*), and Wild goat (*Capra aegagrus*) species were classified using ResNet-18 architecture, which is one of the convolutional neural network models. These data are primarily organized according to the 224x224 resolution size. ResNet ESA architecture was used as a method; It was trained on the ImageNet dataset and achieved a 3.57% error rate, ranking first in the classification success of the ILSVRC-2015 competition with this success rate. Pre-trained ResNet-18 architecture is used as standard ESA. The model was built using the Python programming language and the fast.ai library was used for training and testing the model. In the model, 80% of the data is randomly allocated for training and 20% for testing. The batch_size parameter for the input data is set to 64 and the epoch value used in the model is set to 50. The performance criteria of the model, precision, sensitivity, and f1 score, were obtained as 0.76, 0.75, and 0.75, respectively. As a result of the analysis, the accuracy value of the success of separating the species in the camera trap images was determined as 0.75. Although this success is not at the desired level, it is an important result for camera traps with this method. When examining which camera trap images cannot be separated as a result of the test, more image processing, and image quality come to the fore. Although the method has shortcomings, the results are of great importance in terms of being a good alternative for camera trap data processing problems.

Keywords: Wildlife Animals; Machine Learning; Data Management.

REFERENCES

1. He, K., Zhang, X., Ren, S., Sun, J. (2016). Deep residual learning for image recognition. Proceedings of the IEEE conference on computer vision and pattern recognition, 770-778.
2. Young, S., Rode-Margono, J., & Amin, R. (2018). Software to facilitate and streamline camera trap data management: A review. Ecology and Evolution, 8(19), 9947-9957.

Taxonomic Diversity Calculation in Woody Species: Example of Dedegöl Mountain

Mehmet Güvenç NEGİZ[1], Esra Özge AYGÜL[2], Tayfun İsa AYGÜL[2]

[1] Applied Sciences University of Isparta, Sütçüler Prof. Dr. Hasan Gürbüz Vocational School, Forestry Department, Sütçüler / Isparta, Turkey

[2]Yozgat Bozok University, Boğazlıyan Vocational School, Forestry Department Boğazlıyan / Yozgat, Turkey

mehmetnegiz@isparta.edu.tr

Abstract

Biodiversity includes all species living in a habitat, all genotypic and phenotypic variations within each species, and all spatial and temporal variability in the communities and ecosystems of those species. Taxonomic diversity, which is one of the important parts of biological diversity, allows interpretation by considering the kinship status of species and obtaining information about ecosystem functioning. However, today, there are very few studies evaluating the taxonomic characteristics of species. This study was carried out to determine the taxonomic diversity of woody species in the Dedegöl Mountain region in order to set an example for the studies to be carried out within the scope of taxonomic diversity.

In the study, the data of 48 woody plant species taken from 103 sample areas were used. The new taxonomic diversity index (pTO) based on Deng Entropy, proposed by Özkan (2018), was used to determine the taxonomic diversity in the region. According to the taxonomic diversity values calculated with pTO, the highest taxonomic diversity in the Dedegöl Mountain region was determined in the 39th sample plot, which contains 18 different woody plant species and is located at an altitude of 1367 meters. While Shannon diversity index was determined as 2,60 in the subject area, weighted taxonomic diversity measurement value (TO) 15,28, unweighted taxonomic diversity measurement value (uTO) 10,495, weighted taxonomic distance measurement value (TO^{+}) 15,68, unweighted the taxonomic distance measurement value was calculated as (uTO^{+}) 10,895. When the results obtained in the study are examined in detail, as the altitude of the area increases, the species diversity decreases, while the taxonomic diversity and taxonomic distance values increase.

Keywords: Biodiversity; Shannon Diversity Index; Deng Entropy

REFERENCES

1.Özkan K., A new proposed measure for estimating taxonomic diversity. Turkish Journal of Forestry, Vol:19, No: 336-346, 2015.

2. Deng, Y., Deng entropy. Chaos, Solitons & Fractals, Vol:91, No:1, 549-553, 2016.

ON THE COMPLEX CHARACTERISTICS INTO THE LONG JOSEPHSON JUNCTION MODEL

Fernando S. Vidal Causanilles [1], Haci Mehmet Baskonus [2], Juan Luis Garcia Guirao [3,4], German Rodriguez Bermude [5]

[1] Departamento de Matematica Aplicada y Estadistica. Universidad Politecnica de Cartagena, Hospital de Marina, 30203-Cartagena, Region de Murcia, Spain,
fervicau@gmail.com

[2] Department of Mathematics and Science Education, Faculty of Education, Harran University, Sanliurfa, Turkey
hmbaskonus@gmail.com

[3] Departamento de Matematica Aplicada y Estadistica, Universidad Politecnica de Cartagena, Hospital de Marina, Murcia, 30203, Spain
juan.garcia@upct.es

[4] Department of Mathematics, Faculty of Science, King Abdulaziz University, P.O. Box 80203, Jeddah, 21589, Saudi Arabia

[5] University Centre of Defence at the Spanish Air Force Academy, UPCT-MDE Calle Coronel Lopez Pen~a, s/n, Santiago de la Ribera, Murcia 30720, Spain
german.rodriguez@cud.upct.es

Abstract We investigate the perturbed sine-Gordon equation. We determine the coefficients for the test function of solution formula given by a powerful scheme. The governing perturbation model is used to symbolize the long Josephson junction which is separated by two superconductors.

Keywords: Nonlinear perturbed sine-Gordon equation, Complex travelling wave solutions, Singularly perturbed results.

REFERENCES

1. P.W.Anderson; J.M.Rowell, Probable Observation of the Josephson Tunnel Effect. Phys. Rev. Lett. 10 (6): 230, 1963.
2. J.Guckenheimer, P.Holmes, Nonlinear Oscillations, Dynamical Systems and Bifurcations of Vector Fields, Springer, New York, 1983.
3. A.Barone, G.Patern`o, Physics and Applications of the Josephson Effect, Wiley/Interscience, New York, 1982.
4. T.A.Sulaiman, H.Bulut, H.M.Baskonus, On the exact solutions to some system of complex nonlinear models, Applied Mathematics and Nonlinear Sciences, 6(1), 29-42, 2020.
5. H.M.Baskonus, H.Bulut, T.A.Sulaiman, New Complex Hyperbolic Structures to the Lonngren-Wave Equation by Using Sine-Gordon Expansion Method, Applied Mathematics and Nonlinear Sciences, 4(1), 141-150, 2019.
6. E.I.Eskitascioglu, M.B.Aktas, H.M.Baskonus, New Complex and Hyperbolic Forms for Ablowitz-Kaup-Newell-Segur Wave Equation with Fourth Order, Applied Mathematics and Nonlinear Sciences, 4(1), 105-112, 2019.

TEXTURE SPACE ASSOCIATED WITH GRAPHS

Tuğçe Kunduracı [1], Ceren Sultan Elmalı [2], Tamer Uğur [3]

[1] Department of Mathematics, University of Atatürk, Erzurum, Turkey

[2] Department of Mathematics, University of Erzurum Teknik, Erzurum, Turkey

[3] Department of Mathematics, University of Atatürk, Erzurum, Turkey

tugcek@atauni.edu.tr, ceren.elmali@erzurum.edu.tr, tugur@atauni.edu.tr

Abstract

In this paper, texture graphs are defined as associated with graphs and ditopologies. Also, some properties of texture graphs are investigated.

Keywords: Texture space, ditopology, graph.

REFERENCES

1. Brown, L.M., Ditopological fuzzy structures I, Fuzzy Systems A. I M. 3 (1), (1993).12-17.
2. Brown, L.M., Ditopological fuzzy structures II, Fuzzy Systems A. I M. 3 (2), (1993).5-14.
3. Brown, L.M, Ertürk, R.,Dost, S., Ditopological texture spaces and fuzzy topology, I. Basic concepts, Fuzzy Sets and Systems 147 (2004) 171-199.
4. Brown, L.M, Ertürk, R.,Dost, S., Ditopological texture spaces and fuzzy topology II Topological Considerations , Fuzzy Sets and Systems 147, Issue:2 (2004) 201-231.
5. Brown, L.M, Ertürk, R.,Dost, S., Ditopological texture spaces and fuzzy topology III:Seperation Axioms Fuzzy Sets and Systems 157, (2006) 1886-1912.
6. Evans J.W., Harary F. and Lynn M.S.. On Computer enumeration of finite topologies, Comm. Assoc.Comp.Mach. 10 (1967), 295-298.
7. Bhargav T.N. and Ahlborn T.J., On topological spaces associated with digraphs. Acta, Math.Acad.Sc.Hungar.19(1968)47-52.
8. Kannan, K., A note on some generalized closed sets in bitopological spaces associated to digraphs., journal of app. math. 2012 (2012), 1-5.
9. Sampathkumar E., Transitive Digraphs and topologies on a set, Journal of the Karnatak Univ-Science, 266-273 (1972).
10. Özçağ, S., Brown, L.M.,Di-uniform Texture Space,Applied General Topology, Vol:4, No:1, (2003) 157-192.
11. Diestel, R.: Graph Theory. Springer–Verlag (2005).
12. Bang-Jensen, j. & Gutin, G.: Digraphs: Theory, Algorithms and Applications. Springer–Verlag (2002).
13. Girija B., Pilakkat R., *Bitopological spaces associated with digraphs*, South Asian Journal of Mathematics, Vol.3 (1), (2013) 56-65.

A STUDY ON THE GENERALIZATION OF SOME SIGMOIDAL GROWTH MODELS

Mehmet Korkmaz[1,*], Elif ÖZKURT BAŞUSTAOĞLU[1]

[1] Department of Mathematics, Ordu University, Ordu, Turkey

mkorkmaz52@yahoo.com elif_ozkurtt@hotmail.com

Abstract

In this study, the generalization to some sigmoidal growth models is given in detail. For this purpose, the generalized transformation values and relations of the models are presented with a flow chart. It is stated that the models used are the solutions of the velocity-state differential equations. Their solutions are examined.

Keywords: Sigmoidal Growth Models; Koya-Goshu Growth Model

Acknowledgements: This work is supported by Ordu University, Scientific Research Project (BAP), Project Number: B-1812.

REFERENCES

1. S. Brody, Bioenergetics and Growth, Reinhold Publishing Corporation, New York, 1945.
2. L. von Bertalanffy, Quantitative Laws in Metabolism and Growth, The Quarterly Review of Biology, 3(2), 218, 1957.
3. F. J. Richards, *A* Flexible Growth Function for Empirical Use, Journal of Experimental Botany, 10, 290-300, 1959.
4. C. P. Winsor, *The Gompertz Curve as a Growth C*urve, Proceedings of National Academy of Science, 18(1), 1-8, 1932.
5. J. A. Nelder, *The Fitting of a Generalization of the Logistic Curve*, Biometrics, 17(7),89-110, 1961.
6. L.L. Eberhardt, J.M. Breiwick, Models for Population Growth Curves, ISRN Ecology, 2012.
7. J. O. Rawlings, W. W. Cure, The Weibull Function as a Dose Response Model for Air Pollution Effects on Crop Yields, Crop Science, 25, 807-814, 1985.
8. A. T. Goshu, Simulation Study of the Commonly Used Mathematical Growth Models, Journal of the Ethiopian Statistical Association, 17, 44-53, 2008.
9. Y.C. Lei, S.Y. Zhang, Features and Partial Derivatives of Bertalanffy- Richards Growth Model in Forestry. Nonlinear Analysis: Modelling and Control, 9(1), 65-73, 2004.
10. P. R. Koya, A. T. Goshu, Generalized Mathematical Model for Biological Growths, Open Journal of Modelling and Simulation, 1, 42-53, 2013.

GRAPH STRUCTURE OF A SPECIAL CONGRUENCE SUBGROUP OF THE EXTENDED MODULAR GROUP

Aziz Büyükkaragöz[1] **, Erdal Ünlüyol**[1]

[1] Department of Mathematics, University of Ordu, Ordu, Turkey

azizbk@windowslive.com, erdalunluyol@odu.edu.tr

Abstract

In this paper, firstly we define the congruence subgroup $\widehat{\Lambda}_n(N)$, i.e.

$$\widehat{\Lambda}_n(N) \coloneqq \langle \Lambda_n(N), R(z) \rangle = \Lambda_n(N) \cup R\,\Lambda_n(N).$$

It is obtained by the expanding the

$$\Lambda_n(N) \coloneqq \{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(N):\ a^2 \equiv d^2\ mod\ N, N \in \mathbb{Z}^+, n|N \}$$

congruence subgroup with the reflection $R(z) = -\bar{z}$. Secondly, we research the graph structure of the congruence subgroup $\widehat{\Lambda}_n(N)$ on $\widehat{\mathbb{Q}}_n(N)$,

$$\widehat{\mathbb{Q}}_n(N) \coloneqq \{ \frac{a}{cN} \in \widehat{\mathbb{Q}} : (a, cN) = 1\ \ and\ \ a^4 \equiv 1\ mod\ n \}.$$

Later on, we establish edge condition and be a triangle at the suborbital graphs $\hat{F}_{u,n,N}$. These suborbital graphs are on the set $\widehat{\mathbb{Q}}_n(N)$ of the group $\widehat{\Lambda}_n(N)$. And finally, we have state and prove under what conditions the suborbital graph $\hat{F}_{u,p,N}$ is a forest for $\widehat{\Lambda}_p(N)$ on $\widehat{\mathbb{Q}}_n(N)$. Here p is a prime number.

Keywords: Extended modular group; Congruence subgroup; Edge condition; Triangle, Forest.

REFERENCES

1. G. A. Jones, D. Singerman, K. Wicks, The modular group and generalized Farey graphs, London Math. Soc. Lecture Notes Series 160, CUP, Cambridge, 1991.
2. M. Akbaş, On Suborbital Graphs for Modular Group, Bull. London Math. Soc., Vol: 33, 647-652, 2001.
3. A. Büyükkaragöz, Signatures and graph connections of some subgroups of extended modular group, Ph. D. Thesis, Ordu University, Science Institute, Ordu, Turkey, 2019.

Optical solitons to the generalized higher-order nonlinear Schrodinger equation arising in optical fibers

Salisu Ibrahim[1], Tukur Abdulkadir Sulaiman[2,3], Abdullahi Yusuf[2,3], Ilhame Amirali[4], Usman Younas[5], Hasan Bulut[6]

[1]Department of Mathematics Education, Tishk International University - Erbil, Kurdistan Region, Iraq

[2]Department of Computer Engineering, Biruni University, Istanbul, Turkey

[3]Department of Mathematics, Federal University Dutse, Jigawa, Nigeria

[4]Henan Academy of Big Data/School of Mathematics and Statistics, Zhengzhou University, Zhengzhou 450001, China

[5]Duzce University, Faculty of Science, Duzce, Turkey

[6]Department of Mathematics, Firat University, Elazig, Turkey

Abstract

The optical solitons for the higher order nonlinear Schrodinger equation (NLSE) are established in this study. To construct such novel solutions, we employ robust integration technique called extended rational sinh-cosh. The novel soliton control structure has been studied in order to explain specific physical problems. These findings have been used for the transmission of long-wave and high-power telecommunication networks, which has proven to be beneficial. The circumstances required for the existence of these solitons are presented. Keywords: higher order Schrodinger equation; extended rational sinh-cosh; ‘nonlinearity, optical solitons.

Keywords: Higher order Schrodinger equation; extended rational sinh-cosh; ‘nonlinearity, optical solitons

[1] John, F.: Partial Differential Equations. Applied Mathematical Sciences, 4th edn. Springer, Berlin (1991)

[2] Protter, M.H., Weinberger, H.F.: Maximum Principles in Differential Equations, Corrected Reprint of the1967 Original. Springer, New York (1984)

[3] Schatzman, M.: Numerical Analysis—A Mathematical Introduction. Oxford University Press, Oxford (2002)

[4] Smith, G.D.: Numerical Solutions of Partial Differential Equations, Finite Difference Methods. Oxford Uni-versity Press, New York (1985)

CALCULATING AND MAPPING OF TOPOGRAPHIC COMPLEXITY USING ENTROPY EQUATIONS

Özdemir ŞENTÜRK[1], Ahmet MERT[2]

[1] Department of Forestry, Gölhisar Vocational School, Burdur Mehmet Akif Ersoy University, Burdur, Turkey, osenturk@mehmetakif.edu.tr

[2] Department of Wildlife Ecology and Management, Faculty of Forest, Isparta University of Applied Sciences, Isparta, Turkey, ahmetmert@isparta.edu.tr

Abstract

In applied ecology, environmental factors have been frequently used for distribution modelling of biodiversity. However, the complexities or heterogeneities of those factors have been almost completely ignored. The present study draws attention to this issue. We conducted a study to explain and illustrate how to prepare complexity maps of elevation, radiation index, and slope degree using the data from Burdur Lake watershed, Turkey. In the present study, firstly, large grid sizes of 750x750 meters were formed throughout the watershed. Each grid was divided equally into 36 units with the size of 125x125 meters. Next, for each grid, Shannon Entropy, Rao's Quadratic Entropy, Simpson Index, and Gray Level Co-occurrence Matrix (GLCM) algorithms were calculated for each topographic feature. Lastly, the calculated complexity values were visualized. Thus complexity maps of the topographic features were created for the Burdur Lake watershed.

Keywords: Environmental heterogeneity; Habitat heterogeneity; Landscape diversity; Species diversity; Topographical variability.

REFERENCES

1. Hall-Beyer, M., GLCM texture: a tutorial, National Council on Geographic Information and Analysis Remote Sensing Core Curriculum, Vol: 3, No: 1, 75, 2000.
2. Haralick, R. M., Shanmugam, K., Dinstein, I. H., Textural features for image classification, IEEE Transactions on systems, man, and cybernetics, Vol: 6, 610 – 621, 1973.
3. Rao, C. R., Diversity: its measurement, decomposition, apportionment and analysis, Sankhya: The Indian Journal of Statistics, A 44, 1 – 22, 1982.
4. Shannon, C. E., A mathematical theory of communication, Bell System Technical Journal, Vol: 27, No: 3, 379 – 423, 1948.
5. Simpson, E. H., Measurement of diversity, Nature, Vol: 163, No: 4148, 688, 1949.

Dynamic behavior of the Fokas system arising in monomode optical fibers through the unified auxiliary equation method

Karmina K. Ali[1], Sibel Tarla[2], Resat Yilmazer[2]

[1] Department of Mathematics, University of Zakho, Zakho, Iraq

2 Department of Mathematics, University of Firat, Elazig, Turkey

karmina.ali@uoz.edu.krd

sibeltarla@gmail.com

rstyilmazer@gmail.com

Abstract

In this study, we investigate the Fokas system which represents the spread of irregular pulse in monomode optical fibers by using the unified auxiliary equation method, which is the most powerful method for exploring exact solutions for nonlinear models. The wave transform is used in order to convert a partial differential equation into an ordinary differential equation. As a result, several distinct solutions are obtained such as dark, bright, dark–bright, singular, Jacobi elliptic function, and exponential solutions. Furthermore, 3D surfaces and 2D graphics for the obtained solutions are drawn by choosing appropriate values for the involved free parameters. particular values to the parameters involved.

Keywords: Fokas system; the unified auxiliary equation method; Exact solutions.

REFERENCES

1. Tarla, S., Ali, K. K., Sun, T. C., Yilmazer, R., & Osman, M. S. (2022). Nonlinear pulse propagation for novel optical solitons modeled by Fokas system in monomode optical fibers. Results in Physics, 105381.
2. Zayed, E. M. E., & Shohib, R. M. A. (2019). Solitons and Other Solutions for Two Higher-Order Nonlinear Wave Equations of KdV Type Using the Unified Auxiliary Equation Method. Acta Physica Polonica, A., 136(1).
3. Zhang, S., & Xia, T. (2007). A generalized new auxiliary equation method and its applications to nonlinear partial differential equations. Physics Letters A, 363(5-6), 356-360.

*Abstract Submission should be prepared only **1 page**.

A class of solutions for the Boussinesq-like equations including Spatio-temporal dispersion

Dilara Altan Koç [1] Hajar F. Ismael [2,3] Hasan Bulut[3]

[1] Department of Mathematics, Mugla Sitki Kocman University, Mugla, Turkey

[2]Department of Mathematics, University of Zakho, Zakho, Iraq

[3]Department of Mathematics, University of Firat, Elazig, Turkey

dilaraaltan@mu.edu.tr, hajar.ismael@uoz.edu.krd, hbulut@firat.edu.tr,

Abstract

The Boussinesq equation is a very famous nonlinear evolution equation developed for describing the motion of water with small amplitude and long wave. Boussinesq-type equations which also include dispersion, time dependence and weak nonlinearity have become the most important equation in the prediction of wave transformations in coastal areas. In this study, the $\left(m+\frac{1}{G'}\right)$-expansion method is presented. By using this method, we compute the exact solutions of the Boussinesq-like equations including Spatio-temporal dispersion. As a result of calculations travelling wave solutions are obtained. The graphs of these solutions are drawn and analyzed.

Keywords: $m+\frac{1}{G'(\xi)}$ method; Boussinesq equation; Travelling wave solution.

REFERENCES

1. Anjan Biswas, Daniela Milovic, Arjuna Ranasinghe, Solitary waves of Boussinesq equation in a powerlaw media, Communications in Nonlinear Science and Numerical Simulation 14, 3738–3742, 2009.
2. Christov, C.I., Maugin, G.A., Velarde, M.G., Well-posed Boussinesq paradigm with purely spatial higher-order derivatives, Physical Review E 54, 3621–3638, 1996.
3. Seyma Tuluce Demiray, Yusuf Pandir, Hasan Bulut, New solitary wave solutions of Maccari system. Ocean Engineering, 103, 153–159, 2015.
4. Kang-Jia Wang, Guo-Dong Wang, Solitary and periodic wave solutions of the generalized fourth-order Boussinesq equation via He's variational methods, Mathematical Methods in the Applied Sciences, 44, 5617–5625, 2021.
5. Serbay Duran, Extractions of travelling wave solutions of (2 + 1)-dimensional Boiti–Leon–Pempinelli system via (G′/G, 1/G)-expansion method, Optical and Quantum Electronics,) 53:299, 2021.

OPINIONS OF PRE-SERVICE SCIENCE TEACHERS ABOUT VIRTUAL TOURS TO SCIENCE CENTERS

Esma Kurban [1]

[1] Department of Mathematics and Science Education, Ordu University, Ordu, Turkey

esmakurban19@gmail.com,

Fatma Nur Büyükbayraktar [2]

[2] Department of Mathematics and Science Education, Ordu University, Ordu, Turkey

fnbuyukbayraktar@gmail.com,

Abstract

Science centers are defined as places that offer experimental and hands-on activities for individuals of different age levels, and that make science and technology understandable and accessible (1). Due to reasons such as the lack of science centers in every province in Turkey and the cost of visits, science centers can be reached through virtual tours. In this study, it is aimed to examine the views of pre-service science teachers about their virtual tours to science centers. The sample of the research consists of 6 pre-service teachers studying in the 3rd grade of the Science Teaching undergraduate program of the Faculty of Education of Ordu University. This research was carried out the case study method. In the research, semi-structured interview form and two worksheets were used as data collection tools. The findings obtained in this interview were interpreted through descriptive analysis. According to the results of the analysis of the findings obtained from the interview data, pre-service science teachers define science centers as the means that provide scientific information and technology. At the same time, they think of science centers as areas prepared to conduct experiments and reveal new ideas. Pre-service science teachers define virtual tour to science centers as reaching physical areas in virtual environment. In these virtual tours, besides visiting the areas, they expect to make discoveries by interacting with the areas. However, it has been revealed that pre-service science teachers do not have knowledge about science centers in their own countries. In our study, pre-service science teachers were provided to get to know Konya Science Center through virtual tours. During the virtual tour were made applications. In these applications, activities that pre-service science teachers can use in their professional lives are included. It is thought that virtual tours will contribute to the professional development of pre-service science teacher.

Keywords: Science centers; Virtual tour; Pre-service science teacher.

REFERENCES

1. TUBITAK (The Scientific and Technological Research Council of Turkey), 2019, http://www.tubitak.gov.tr/

ON SOLUTION OF THE NON-NEWTONIAN VOLTERRA INTEGRAL EQUATIONS WITH CONVOLUTION KERNEL

Nihan Güngör [1]

[1] Department of Mathematical Engineering, University of Gumushane, Gumushane, Turkey

nihanmath@gmail.com,

Abstract

The non-Newtonian calculus, which has many field of application such as interest rates, theory of elasticity in the economy, the viscosity of the blood, computer science including image processing and artificial intelligence, biology, differential equations, functional analysis and probability theory, was built as an infinite family of calculus consisting of the branches of geometric, bigeometric, harmonic, biharmonic, quadratic and biquadratic calculus by Grossman and Katz [1]. In this study, the Volttera integral equations with convolution kernel are defined in the non-Newtonian calculus with the aid of $*$-integral. The focus of this study is obtained the exact solutions of the convolution type linear non-Newtonian Volterra integral equations of the first kind and also the second kind by using non-Newtonian Laplace transform.

Keywords: Non-Newtonian calculus; Volterra integral equations; Integral transforms.

REFERENCES

1. Grosmann M., Katz R., Non-Newtonian Calculus, Lee Press, Pigeon Cove Massachussets, 1972.
2. Grosmann M., An Introduction to Non-Newtonian Calculus, International Journal of Mathematical Education in Science and Technology, 10(4), 525-528, 1979.
3. Erdoğan M., Duyar C., Non-Newtonian Improper Integrals, Journal of Science and Arts 1(42), 49-74, 2018.
4. Güngör N., A note on linear non-Newtonian Volterra integral equations, Mathematical Sciences, 1-15, https://doi.org/10.1007/s40096-021-00427-z, 2021.

A STUDY ON CONJUGATE CONNECTIONS ON PARA ANTI-HERMITIAN MANIFOLDS

Sibel Turanlı[1] **and Sedanur Uçan**[2]

[1] Department of Mathematics, Faculty of Sciences, University of Erzurum Technical, Erzurum, Turkey

[2]Department of Mathematics, Institute of Science, University of Atatürk, Erzurum, Turkey

sibel.turanli@erzurum.edu.tr

sedanurr15@gmail.com

Abstract

Let ∇ be a linear connection on an almost para anti-Hermitian manifold M_{2n}, equipped with an almost para complex structure φ and a pseudo-Riemannian metric g. Also, associated para anti-Hermitian metric of almost para anti-Hermitian manifold is defined by $G(X,Y) = (g \circ \varphi)(X,Y)$ and called twin para anti-Hermitian metric of g. In this study, we introduce three types of conjugate connections of linear connections relative to g, φ, G and investigate some properties of these conjugate connections.

Keywords: Conjugate Connections; para anti-Hermitian Manifolds.

REFERENCES

1. Salimov, A. A., Iscan, M., Akbulut, K., Notes on Para-Norden-Walker 4-Manifolds, International Journal of Geometric Methods in Modern Physics, 7(8), 2010, 1331-1347.

2. Gezer, A., Cakicioglu, H., Notes Concerning Codazzi Pairs on Almost Anti-Hermitian Manifolds, Inpressed.

3. Fei, T., Zhang, J., Interaction of Codazzi Couplings with (Para-) Kahler Geometry, Results Math, 72, 2017, 2037-2056.

4. Nomizu, K., Simon, U., *Notes on Conjugate Connections*, In Geometry and Topology of Submanifolds IV, eds. F. Dillen and L. Verstraelen, Word Scientific, 1992.

THE EXPLOSION OF THE SOLUTIONS OF STOCHASTIC WAVE EQUATIONS

Beytullah YAĞIZ[1], Yener ALTUN[2], Sinan DENİZ[3]

[1]Institute of science, Department of Statistics, Van YYU, Van, Turkey
[2] Faculty of Economics and Administrative Sciences, Department of Business Administration, Van YYU, Van, Turkey
[3]Institute of science, Department of Statistics, Van YYU, Van, Turkey
[1]beytmat@gmail.com, [2]yeneraltun@yyu.edu.tr, [3]sinan_deniz65@hotmail.com;

Abstract

In this study, a class of nonlinear stochastic wave equations is considered. Using appropriate energy inequalities, some sufficient conditions are established for the local solutions of the stochastic wave equation to blow-up with positive probability. Under these conditions, it is examined whether the solution of the equation in question exists in finite and infinite time. As a result, it has been determined that the solution of the equation approaches infinity as the variable approaches a finite time.

Keywords: Blow-up, Global solution, Local solution, Stochastic wave equation.

REFERENCES

1. Arnold, L., 1992. ***Stochastic Differential Equations: Theory and Applications***. A Wiley - Inter Science Publication., USA.
2. Bo, L., Tang, D., Wang, Y., 2008. Explosive solutions of stochastic wave equations with damping on Rd. ***Journal of Differential Equations,*** **244:** 170–187.
3. Chow, P. L., 2002. Stochastic Wave Equations with Polynomial Nonlinearity, ***Ann. Appl. Probab.*****12 (1):** 361-381.

ESTIMATING SAMPLE COMPLETENESS FOR BIODIVERSITY ASSESSMENT BASED ON TREE AND SHRUB SPECIES IN SARIKAYA WLDA/ANTALYA

Mertcan Gülben[1], Emirhan Berberoğlu[2], Zeynep Öz[1], Murat Görgöz[3], Candan Aykurt[1], Kürşad Özkan[3], Özdemir Şentürk[4]

[1] Akdeniz University, Department of Biology, Antalya, Turkey.

[2] Akdeniz University, Department of Geography, Antalya, Turkey.

[3] Isparta University of Applied Sciences, Department of Forest Engineering, Isparta, Turkey.

[4] Burdur Mehmet Akif Ersoy University, Department of Forestry, Burdur, Turkey.

gulbenmertcan@gmail.com

Abstract

We carried out a study to test sample completeness in Sarıkaya Wildlife Development Area (WLDA), one of the most important biodiversity hotspots in Turkey. The data used consists of 103 tree and shrub species collected from 104 sampling plots. To test sample completeness, we used five methods, Chao 2 $(\hat{S}_{Chao2})$, adjusted Chao 2 $(\hat{S}_{iChao2})$, Jacknife 1 $(\hat{S}_{jk1})$, Jacknife 2 $(\hat{S}_{jk2})$, boostrap $(\hat{S}_{boot})$ and expected species richness based on the number of only once seen species $(\hat{S}_{Q_1})$.

The results indicated that the lowest estimated richness values were provided by $\hat{S}_{Chao2}$, $\hat{S}_{iChao2}$ with 113 and 115 species, respectively. The highest estimated richness corresponds to the values of 123, 122 and 121 obtained by $\hat{S}_{jk2}$, $\hat{S}_{Q_1}$ and $\hat{S}_{jk1}$ while the value of $\hat{S}_{boot}$ is equal to 112. According to average values of the estimators, estimated sample completeness corresponds to a value of 87.5% with the confidential intervals between 85% and 90.2%. This relative value provides sufficient information to examine vegetation-environmental relationships or ecological land classification because woody species have been generally used for such studies as explanatory data. However, for biodiversity studies, not only woody species but also herb species are considered. Hence, we concluded that field work needs to continue to grand sufficient data collection by taking extra sampling plots in the study area.

Keywords: true species richness, sample coverage, negative bias, bio complexity, bias-corrected estimators

REFERENCES

1. Brose, U., Martinez, N.D. & Williams, R.J., Estimating species richness: sensitivity to sample coverage and insensitivity to spatial patterns, Ecology, 84, 2364–2377, 2003.
2. Chao, A. and Chiu, C. H., Species richness: estimation and comparison. Wiley StatsRef: Statistics Reference Online, 1-26, 2016.
3. Chiu, C. H., and Chao, A., Estimating and comparing microbial diversity in the presence of sequencing errors PeerJ, 4, e1634-e1634, 2016.
4. Gotelli, N. J. and Chao, A., Measuring and estimating species richness, species diversity, and biotic similarity from sampling data, in The Encyclopedia of Biodiversity 2nd Edition, (ed S.A. Levin), Elsevier, New York, pp. 195–211, 2013.
5. Özkan, K., Estimating True Species Richness from Braun Blanquet Scale, Kastamonu Univ. Journal of Forestry Faculty, 21(3), 306-314, 2021.

LMI-BASED STABILITY ANALYSIS OF NONLINEAR SYSTEMS WITH TWO ADDITIVE TIME-VARYING DELAY COMPONENTS

Yener ALTUN[1]

[1] Faculty of Economics and Administrative Sciences, Department of Business Administration, Van YYU, Van, Turkey

yeneraltun@yyu.edu.tr

Abstract. This study is concerned with the LMI-based stability analysis of nonlinear systems with two additive time-varying delay components in Lyapunov–Krasovskii framework. By constructing a suitable augmented Lyapunov-Krasovskii functional (LKF) with integral terms, some new stability criteria are established in terms of linear matrix inequalities (LMIs), which are easily tested by various convex optimization techniques. More information of the lower and upper delay bounds of time-varying delays are used to obtain the stability criteria, which can lead less conservative results. In this direction, this study presents a new stability analysis result for a class of nonlinear systems with two additive time delays. Finally, two numerical examples are given to demonstrate the effectiveness and to show the applicability of the proposed method over existing results by using the MATLAB-Simulink.

Keywords: LKF, Stability, Additive time delay, LMI.

REFERENCES

1. Altun, Y. (2020). Improved results on the stability analysis of linear neutral systems with delay decay approach. Math Meth Appl Sci. **43**(3), 1467–1483.
2. Ramakrishnan, K., Ray, G. (2010). Stability criteria with less LMI variables for linear system with additive time-delays. J. Appl. Math. Comput. 215, 3740–3745.
3. Jiao, J. M. (2013). A stability criterion for singular systems with two additive time-varying delay components. Int. J. Autom. Comput. 10(1), 39–45.

The potential distribution of *Teucrium chamaedrys* L. in Turkey

Ahmet ACARER[1], Halil İbrahim ÜRKMEZ[2], Ali ŞENOL[2], Ahmet MERT[1]

[1] Department of Wildlife Ecology and Management, Faculty of Forest, Isparta University of Applied Sciences, Isparta, Turkey

[2] Department of Forest Engineering, Faculty of Forestry, Isparta University of Applied Sciences, Isparta, Turkey

aacarer32@gmail.com

Abstract

Teucrium chamaedrys L. is a plant species evaluated as an evergreen and non-wood forest product. It has been used in the treatment of diseases such as hypertension, gout, diabetes and obesity from past to present. Because of these properties, it has been consumed in herbal tea and capsule form since the 1980s. It is aimed to determine the potential distribution of *Teucrium chamaedrys* L. species depending on environmental factors in Turkey, which contributes to the economy of Turkey due to its medicinal and aromatic plant properties. For this purpose, present data records of *Teucrium chamaedrys* L. species in Turkey were downloaded from the Global Biodiversity Information Facility (www.gbif.org). Then, the 63 present data of the target species were associated with the independent variables using the Maximum entropy method and the potential distribution map of the species in Turkey was obtained.

Keywords: Maximum entropy method; Medicinal and Aromatic plants; Modeling and mapping; Plant species distribution.

REFERENCES

1 Elith, J., Phillips, S. J., Hastie, T., Dudík, M., Chee, Y. E., & Yates, C. J. (2011). A statistical explanation of MaxEnt for ecologists. Diversity and distributions, 17(1), 43-57.

2 GBIF (2018). GBIF Occurrence Data. https://doi.org/ 10.15468/dl.fbftf6. Erişim: 21 Nisan 2022.

3 Piccolella, S., Scognamiglio, M., D'Abrosca, B., Esposito, A., Fiorentino, A., & Pacifico, S. (2021). Chemical Fractionation Joint to In-Mixture NMR Analysis for Avoiding the Hepatotoxicity of *Teucrium chamaedrys L. subsp. chamaedrys*. Biomolecules,11(5), 690. (https://doi.org/10.3390/biom11050690).

Approximation by a new construction of parametric Baskakov operators

Fatih Rıza Çelik[1] , **Gürhan İçöz**[1]

[1] Department of Mathematics, University of Gazi, Ankara, Turkey

fatihrizacelik@gmail.com, gurhanicoz@gazi.edu.tr

Abstract

In the present study, we introduce a non-negative real parametric generalization of Baskakov operators and call them α-Baskakov operators whose construction depends on a real-valued function τ by using two sequences u_n and v_n of functions. We prove that the new operators provide a weighted uniform approximation on [0,∞). In terms of weighted moduli of smoothness, we obtain degrees of approximation associated with the function τ. Also, we prove the Voronovskaya-type theorem.

Keywords: Baskakov, approximation, parametric, sequence

REFERENCES

1. Usta, F. On approximation properties of a new construction of Baskakov operators. Adv Differ Equ 2021, 269 (2021).
2. A. Aral, H. Erbay, Parametric generalization of Baskakov operators, Math. Commun., 24 (2019),119—131
3. Patel, P., Mishra, V.N., Örkçü, M.: Some approximation properties of the generalized Baskakov operators. J. Interdiscip. Math. 21(3), 611--622 (2018)

To Alternate the Routes Encoded by Bloom Filters in Computer Networks Represented by Graphs

Gökçe ÇAYLAK KAYATURAN [1]

[1] Department of Mathematics, Ordu University, Ordu, Turkey

gokcekayaturan@odu.edu.tr,

Abstract

Compressing the data to take up less space can be considered as a solution for the reliability of data exchange between computers in computer network systems, communication speed between computers, the data traffic in the network or the limited storage areas in the computers for the big data . There may be many other solutions to be produced for the problems can be faced during routing in computer networks, as well as the use of the Bloom filter is one of them. The Bloom filter is a construct used to represent a set in compressed form [1]. However, it has potential to produce errors denoted by false positives with a probability. In order to produce mathematical solutions to real-time network problems such as network attacks or physical disruptions during message transportation or communication between computers in networks, a computer network system model can be represented by a mathematical graph and the edges of the graph are represented by Bloom filters. We aim to evaluate the probability of false positives that may be encountered during message transmission in the computer network system and possible physical deterioration situations that may occur on the edges of the graph, and accordingly we present the most appropriate and reliable transmission paths. With some assumptions (such as the graph is known, the risk situations of the roads are known, and whether the paths are the shortest path between two given points), We aim to obtain a model that presents the alternative routes to the users in an undirected network systems.

Keywords: Computer Network; Bloom filter; False positives.

REFERENCES

1. Burton H Bloom, Space/time trade-offs in hash coding with allowable errors, Communications of the ACM, Vol:13, No:7, 422–426, 1970.

Time Optimal Trajectory Generation in Joint Space for 6R Industrial Serial Robots Using Cuckoo Search Algorithm

Oguzhan KARAHAN[1], Hasan KARCI[2], Ali TANGEL[3]

[1,2,3]Department of Electronics and Communications Engineering, Kocaeli University, Kocaeli, Turkey

oguzhan.karahan@kocaeli.edu.tr, hsnkarci06@gmail.com, atangel@kocaeli.edu.tr

Abstract

In this paper, an optimal trajectory planning approach is proposed based on optimal time by utilizing the interpolation spline method. The method including a combination of cubic spline and 7th order polynomial is used for generating the trajectory in joint space for robot manipulators. Cuckoo Search (CS) optimization algorithm is chosen to optimize the joint trajectories based on objective, namely minimizing total travelling time along the whole trajectory. The spline method has been applied on PUMA robot for optimizing the joint trajectories with the CS algorithm based on objective. Moreover, results from the proposed algorithm have been compared with that of the algorithms suggested by earlier studies. With the trajectory planning method, the joint velocities, accelerations and jerks along the whole trajectory optimized by CS meet the requirements of the kinematic constraints in case of objective. Simulation results validated that the used trajectory planning method based on the proposed algorithm are very effective in comparison with the same methods based on the algorithms proposed by earlier authors.

Keywords: Trajectory planning, Interpolation spline methods, Industrial robots, CS.

Call Assistant Application for Call Centers

Mesut Tartuk[1], Furkan Taha Nurdağ[2], Vedat Acar[3], Mehmet Fatih Akay[4], Fatih Abut[5]

[1]*Comdata Teknoloji ve Müşteri Hizmetleri A.Ş., Turkey; mesut.tartuk@comdatagroup.com*

[2]*Comdata Teknoloji ve Müşteri Hizmetleri A.Ş., Turkey; furkan.nurdag@comdatagroup.com*

[3]*Comdata Teknoloji ve Müşteri Hizmetleri A.Ş., Turkey; vedat.acar@comdatagroup.com*

[4]*Çukurova University, Computer Engineering Department, Adana, Turkey; mfakay@cu.edu.tr*

[5]*Çukurova University, Computer Engineering Department, Adana, Turkey; fabut@cu.edu.tr*

Abstract

In this study, we develop a new application to handle incoming calls into call centers when the centers are on holiday, non-working time, or peak call time. Previously, we were using a Voicemail server from Avaya by license. To reduce side effects like overloading and high currency cost, we made a definition between Interactive Voice Response (IVR) and Central. With this definition, the load of the server is reduced significantly. Then, we developed an intermediate application that would forward the incoming calls from the exchange to the server for recording incoming calls on the IVR side. Incoming Voicemail will be directed to the IVR according to the determined conditions, and the customer's speech will be translated into the written text over IVR. Afterward, these texts will be sent by email to the authorized people within the defined projects. This mail will contain the voice file and information of the person who sent this Voicemail. Afterward, the authorized person who receives this email will respond to this request within his own team.

Keywords: Voicemail; Callback Requests; Abandon Calls; Interactive Voice Response

REFERENCES

[1] LeFebvre, L. E., & Haggadone, B. A. (2019). Remembering the voice: Exploring information and sentiment in Voicemail archival practices. Mobile Media & Communication, 7(2), pp. 195-214.

[2] Mehrotra, V., Ozlük, O., & Saltzman, R. (2010). Intelligent procedures for intra-day updating of call center agent schedules. Production and operations management, 19(3), pp. 353-367.

[3] Ringel, M., & Hirschberg, J. (2002). Automated message prioritization: Making Voicemail retrieval more efficient. In Procs. of Extended Abstracts on Human Factors in Computing Systems (pp. 592-593).

AN APPLICATION ON THE EXTENDED TECHNOLOGY ACCEPTANCE MODEL

Sinan DENİZ[1] , Şakir İŞLEYEN[2] , Beytullah YAĞIZ[3]

[1] Institute of science, Department of Statistics, Van YYU, Van, Turkey

[2] Faculty of Economics and Administrative Sciences, Department of Econometrics, Van YYU, Van, Turkey

[3] Institute of science, Department of Statistics, Van YYU, Van, Turkey

[1] sinan_deniz65@hotmail.com; [2] sakirisleyen@yyu.edu.tr, [3] beytmat@gmail.com;

The effective use of technology in education, which affects every aspect of life in the age we live in, has an important place. It is of great importance to determine the factors affecting the use of technology in education and training. In this study, data analyzes were performed using IBM SPSS 22 and AMOS V24 programs. In this process, some results were obtained by using various statistical techniques such as structural equation modeling (SEM), multivariate statistical techniques (Confirmatory Factor Analysis (CFA) and bootstrap) and descriptive statistics. As a result of the analysis, it was determined that perceived ease of use, attitude towards use and perceived usefulness had positive effects on behavioral intention and results similar to the literature were obtained.

Keywords: Technology Acceptance Model, Structural Equation Model, Multivariate Statistical Techniques.

REFERENCES

Ajzen, I. (1991). Theory of Planned Behavior, Organizational Behavior and Human Decision Processes. *50* (2), 179-211.

Çelik, H. E., Arslan, T. and Yilmaz, V. (2012). Examination of satisfaction from family medicine by structural equation modeling: The example of Turkey. *African Journal of Business Management, 6*(9), 3282-3287.

Davis, F. D., Bagozzi, R. P. and Warshaw, P. R. (1989). User Acceptance of Computer Technology: A Compari-son of Two Theoretical Models. Management Science, 35 (8), 982-1003.

NEW OPERATION TYPES ON INTERVAL-VALUED NEUTROSOPHIC SOFT SETS

Yıldıray Çelik

Department of Mathematics, University of Ordu, Ordu, Turkey

ycelik61@gmail.com

Abstract

In this paper, we introduce the notion of interval-valued neutrosophic soft set which is a generalization of neutrosophic soft sets. Then we define some new operations on interval-valued neutrosophic soft sets and investigate basic properties related this operations. We also present numerical examples for better understanding of the subject.

Keywords: Interval-valued set; Neutrosophic set; Interval-valued neutrosophic set.

REFERENCES

1. I. Deli, Interval-valued neutrosophic soft sets and its decision making, International Journal of Machine Learning and Cybernetics, Vol:8, 1-12, 2015.
2. P.K. Maji, Neutrosophic soft set, Annals of Fuzzy Mathematics and Informatics, Vol:5, No:1, 157-168, 2013.
3. D. Molodtsov, Soft set theory-first results, Computers and Mathematics with Applications, Vol:37, 19-31, 1999.
4. F. Smarandache, Neutrosophic set-a generalization of the intuitionistic fuzzy set, Granular Computing, 2006 IEEE International Conference, 38-42, 2006.
5. X. Yang, T.Y. Lin, J. Yang, Y. Li and D. Yu, Combination of interval-valued fuzzy set and soft set, Computers and Mathematics with Applications, Vol:58, No:3, 521-527, 2009.

A NEW VIEW ON INTUITIONISTIC FUZZY SOFT GRAPHS

Yıldıray Çelik [1], **Gül Ergün**[2]

[1]Department of Mathematics, University of Ordu, Ordu, Turkey

ycelik61@gmail.com

[2]Department of Mathematics, University of Ordu, Ordu, Turkey

gul.ergunn06@gmail.com

Abstract

In this paper, we introduce the concept of intuitionistic neutrosophic soft graphs and give some new operations such as parametric $\wedge$-intersection and parametric $\vee$-union on intuitionistic neutrosophic soft graphs. We have also applied the concept of intuitionistic neutrosophic soft graph in a decision-making problem and then give an algorithm fort the selection of optimal object.

Keywords: Intuitionistic neutrosophic soft set; Intuitionistic neutrosophic soft graph; Decision-making problem.

REFERENCES

1. M. Akram, S. Shahzadi, Neutrosophic soft graphs with applications, Journal of Intelligent and Fuzzy Systems, Vol:32, No:1, 841-858, 2017.
2. K. Atanassov, Intuitionistic fuzzy sets, Fuzzy Sets and Systems, Vol:20, 87-96, 1986.
3. P. Bhattacharya, Some Remarks on Fuzzy Graphs, Pattern Recognition Letters, Vol:6, No:5, 1987.
4. S. Broumi, F. Smarandache, Intuitionistic Neutrosophic Soft Set, Journal of Information and Computing Science, Vo:8, No:2, 130-140, 2013.
5. F. Smarandache, Neutrosophic set-a generalization of the intuitionistic fuzzy set, Granular Computing, IEEE International Conference, 38-42, 2006.
6. R. Parvathi, M.G. Karunambigai, Intuitionistic Fuzzy Graphs, Computational intelligence, theory and application, Springer, Berlin, Heidelberg, 139-150, 2006.

A Study on the Solutions of (1+1)-Dimensional Mikhailov-Novikov-Wang Equation

Seyma Tuluce Demiray, Ugur Bayrakci

Department of Mathematics, University of Osmaniye Korkut Ata, Osmaniye, Turkey

seymatuluce@gmail.com, ubayrakci42@gmail.com

Abstract

The main purpose of this study is to obtain various solutions of the (1+1) dimensional Mikhailov-Novikov-Wang integrable equation. For this purpose, the generalized exponential rational function method has been applied to this equation. Thus, some trigonometric function, hyperbolic function and bright soliton solutions of the studied equation were obtained. In addition, 2D and 3D graphics of the obtained solutions were drawn for certain values. The obtained results and the graphic drawings of the results were made using Wolfram Mathematica 12.

Keywords: Generalized exponential rational function method; (1+1)-dimensional Mikhailov-Novikov-Wang integrable equation.

REFERENCES

1. Nauman Raza, Aly R. Seadawy, Saima Arshed, Muhammad H. Rafiq, A variety of soliton solutions for the Mikhailov-Novikov-Wang dynamical equation via three analytical methods, Journal of Geometry and Physics, Vol:176, 104515, 2022.
2. Santanu Saha Ray, Shailendra Singh, New various multisoliton kink-type solutions of the (1+1)-dimensional Mikhailov–Novikov–Wang equation, Mathematical Methods in the Applied Sciences, Vol:44, No:18, 14690-14702, 2021.
3. Hari M. Srivastava, Hatira Günerhan, Behzad Ghanbari. Exact traveling wave solutions for resonance nonlinear Schrödinger equation with intermodal dispersions and the Kerr law nonlinearity. Mathematical Methods in the Applied Sciences, Vol:42, No:18, 7210-7221, 2019.
4. Hajar Farhan Ismael, Hasan Bulut, Haci Mehmet Baskonus, W-shaped surfaces to the nematic liquid crystals with three nonlinearity laws, Soft Computing, Vol:25, 4513-4524, 2021.

ON STATISTICAL CONVERGENCE OF GENERALIZED FUZZY NUMBER SEQUENCES

Büşra Nur ER and Yavuz ALTIN

Department of Mathematics, Firat University Elazig, Turkey

nurb37332@gmail.com, yaltin23@yahoo.com

Abstract Fuzzy set theory was first studied by Zadeh[7]. This theory has been applied to many different fields of science. Fuzzy number sequences play an important role in mathematical analysis and functional analysis. In this study, we analyzed some inclusion theorems by defining a new statistical convergent fuzzy sequence space using Δ_p^m −difference sequence.

Keywords: Statistical Convergence, Fuzzy Sequence.

REFERENCES

1. Fast, H. Sur la convergence statistique, Colloq. Math.. 2 241-244, 1951.

2. Et, M.; Çolak, R. On some generalized difference sequence spaces. Soochow J. Math. 21 (1995), no. 4, 377-386.

3. Karakaş, A.; Altin, Y.; Et, M. Δ_p^m −statistical convergence of order α. Filomat 32 (2018), no. 16, 5565-5572.

4. Kızmaz, H. On certain sequence spaces. Canad. Math. Bull. 24 (1981), no. 2, 169-176

5. Matloka, M. Sequences of fuzzy numbers, BUSEFAL 28, 28-37, 1986.

6.Nuray, F. ; Savaş, E. Statistical convergence of sequences of fuzzy real numbers, Math. Slovaca 45 (3) 269-273, 1995.

7. Zadeh, L. A. Fuzzy sets. Information and Control 8 (1965), 338-353.

Neimark-Sacker bifurcation in a prey-predator model with conformable fractional derivative

Guven KAYA[1]

[1] Department of Mathematics, Bingol University, Bingol,Turkey

gkaya@bingol.edu.tr,

Abstract

Using bifurcation analysis is a very effective method to better understand the dynamic behavior of competitive models. Recently, many researchers have studied the bifurcation theory[1-3]. In this paper, we examine Neimar-Sacker bifurcation of a prey-predator model with conformable fractional derivative using the center manifold theorem and bifurcation theory. In addition, the effect of fractional order derivative on dynamic behavior of the model is investigated. Finally, all theoretical results are supported by numerical simulations.

Keywords: Conformable fractional derivative; Discretization; Stability; Bifurcation.

REFERENCES

1. M. S. Khan, M. A. Khan, M. S. Shabbir and Q. Din, Stability, bifurcation and chaos control in a discrete-time prey-predator model with Holling type-II response. Network Biology, Vol:9, No:3, 58-77, 2019.
2. F. Gurcan, G. Kaya and S. Kartal, Conformable fractional order Lotka-Volterra predator-prey model: discretization, stability and bifurcation. Journal of Computational and Nonlinear Dynamics, Vol:14, No:11, 111007, 2019.
3. Q. Din, M. S. Shabbir, M. A. Khan, M. A. and K. Ahmad, Bifurcation analysis and chaos control for a plant-herbivore model with weak predator functional response, Journal of biological Dynamics, Vol:13, No:1, 481-501, 2019.
4. E. Balcı, I. Ozturk and S. Kartal, Dynamical behaviour of fractional order tumor model with Caputo and conformable fractional derivative. Chaos, Solitons & Fractals, Vol:123, 43-51, 2019.
5. Q. Din, Dynamics and chaos control for a novel model incorporating plant quality index and larch budmoth interaction. Chaos, Solitons & Fractals, Vol:153, 111595, 2021.

APPROXIMATION PROPERTIES OF A NEW TYPE OF GAMMA OPERATOR

Gürhan İçöz[1], Seda Demir[2]

[1,2]Department of Mathematics, University of Gazi, Ankara, Turkey

gurhanicoz@gazi.edu.tr, seda.demir@gazi.edu.tr

Abstract

In this paper, a new type of Gamma operator will be presented. The approximation properties of the presented operator are tested on the Voronovskaya type theorem, weighted approximation. The rate of convergence was found and pointwise estimation was examined. Numerical example is given to show that the last newly defined operator approaches the function faster.

Keywords: Gamma function; voronovskaya type theorem; weighted approximation; rate of convergence; pointwise estimates.

REFERENCES

1. Alexandru Lupaş, M. Müller, Approximationseigenschaften der Gammaoperatoren, Mathematische Zeitschrift, Vol:98, 208–226, 1967.
2. Chrysi G. Kokologiannaki, Valmir Krasniqi, Some properties of the k-Gamma function, Le Matematiche, Vol:68, 13–22, 2013.
3. Asifa Tassaddiq, A new representation of the k-Gamma Functions, Mathematics, Vol:7, No:133, 1–3, 2019.
4. Ömür Betuş, Fuat Usta, Approximations of functions by a new type of Gamma operators, Numerical Methods for Partial Differential Equations, 1–12, 2020.
5. Qing-Bo Cai, Bayram Çekim, Gürhan İçöz, Gamma Generalization Operators Involving Analytic Functions, Mathematics, Vol:9, No:13, 1–8, 2021.
6. Fuat Usta, Ömür Betuş, A new modification of Gamma operators with a better error estimation, Linear and Multilinear Algebra, 1–12, 2020.
7. Ronald A. DeVore, George G. Lorentz, Constructive Approximation, Springer-Verlag,1993.

On metrizability of topological groups

Ali FARAJZADEH[1], Alkan ÖZKAN[2]

[1]Department of mathematics, Faculty of Science, Razi University, Kermanshah, Iran

farajzadehali@gmail.com , A.Farajzadeh@razi.ac.ir

[2]Department of Mathematics, Faculty of Arts and Sciences, Igdır University, Igdır, Turkey

alkan.ozkan@igdir.edu.tr, alkan_mat@hotmail.com

Abstract

In this paper, we are going to discuss about the following facts and when a topological space equipped with an algebraic structure, for instance group structure, how can one reduce the assumptions given in the facts.

Fact 1. If X is a regular, locally separable, meta-Lindelöf, $w\Delta$-space with a G_δ−diagonal, then X is metrizable.

Remember that a topological space X is called G_δ-diagonal if $\{(x,x): x \in X\}$ is a G_δ subset of $X \times X$, that is it is as intersection of a family of open subsets of $X \times X$. Also the topological space X is said to be metacompact (paracompact) if each open cover U of X has an open refinement SS such that each $x \in X$ is an element of only finitely many elements of SS; SS is said to be a point-finite refinement of U. X is said to be para-Lindelöf (meta-Lindelöf) if each open cover U of X has an open refinement SS such that each $x \in X$ has a neighbourhood which intersects only countably many elements of $x \in X$.

Fact 2. A regular metacompact M−space X is metrizable if and only if it has a G_δ .

Fact 3. Regular space X with a uniform base is metrizable if and only if it is an M−space.

Fact 4. If X is a regular, locally countably compact, metacompact space with a G_δ-diagonal , then X is metrizable and locally compact.

Keywords: Regular, locally separable, metacompact, topological group

REFERENCES

1. P. J. Cohen. Ideal approach to convergence in functional spaces. https://arxiv.org/abs/2111.05049, 2021.

A COMBINED FINITE DIFFERENCE-COLLOCATION METHOD FOR FITZHUGH-NAGUMO EQUATION

Ömer Oruç [1]

[1] Department of Mathematics, Dicle University, Diyarbakır, Turkey

omer.oruc@dicle.edu.tr,

Abstract

In this study, we propose a numerical method to solve Fitzhugh–Nagumo equation which is an important partial differential equation in biology. The proposed numerical method uses finite difference method for time discretization of the equation. For space discretization Fibonacci and Lucas polynomials are used. Two problems are solved numerically. Acquired numerical solutions are compared with exact solutions and with methods available in literature. The comparsions indicate efficiency of the proposed method.

Keywords: Lucas polynomials; Partial differential equations; Computational method.

REFERENCES

1. R. Jiwari, R.K. Gupta, Vikas Kumar, Polynomial differential quadrature method for numerical solutions of the generalized Fitzhugh–Nagumo equation with time-dependent coefficients, Ain Shams Engineering Journal, 5, 1343–1350, 2014.
2. Ö. Oruç, A new algorithm based on Lucas polynomials for approximate solution of 1D and 2D nonlinear generalized Benjamin-Bona-Mahony Burgers equation, Computers & Mathematics with Applications, 74 (12), 3042-3057, 2017.
3. J. Nagumo, S. Arimoto, S. Yoshizawa, An active pulse transmission line simulating nerve axon, Proceedings of the IRE, 50, 2061–2070, 1962.
4. R. Fitzhugh, Impulse and physiological states in models of nerve membrane. Biophys Journal,1(6), 445–66, 1961.

Nonlinear differential equations with according to the Darboux frame

Fatma BULUT[1], **Mehmet BEKTAŞ**[2]

[1] Department of Mathematics, University of Bitlis Eren, Bitlis, Turkey.

[2] Department of Mathematics, University of Firat, Elazig, Turkey.

fbulut@beu.edu.tr, mbektas@firat.edu.tr.

Abstract

In the current study, we define k-type slant helices of timelike curves in Euclidean 4-space consisting of Darboux vector. Also, we characterize k-type slant helices in terms of their curvatures. On the other hand, nonlinear differantial equations are obtained by utilizing k-type slant helices.

Keywords: Nonlinear differential equations; Curve; Darboux frame.

REFERENCES

[1] Ali A.T., Önder M., Some characterizations of rectifying spacelike curves in the Minkowski space-time, Global Journal of Science Frontier Research Mathematics, 12(1), 2249-4626, 2012.

[2] Camcı Ç., İlarslan K., Kula L., Hacısalihoğlu H.H., Harmonic curvatures and generalized helices in E^n, Chaos, Solitons and Fractals, 40(5), 2590-2596, 2009.

[3] Düldül M., Vector fields and planes in E^4 which play the role of Darboux vector, Turkish Journal of Mathematics, 44, 274-283, 2020.

[4] Izumiya S., Takeuchi N., New special curves and developable surfaces, Turkish Journal of Mathematics, 28(2), 153-163, 2004.

[5] İlarslan K., Boyacıoğlu Ö., Position vectors of a timelike and a null helix in Minkowski 3-space, Chaos, Solitions and Fractals, 38, 1383-1389, 2008.

[6] İlarslan K., Nesovic E., Some characterizations of null osculating curves in the Minkowski space-time, Proceedings of the Estonian Academy of Sciences, 61(1), 1-8, 2012.

Potential distribution modeling of *Styrax officinalis* L. in Mediterranean ecosystems

Serkan Gülsoy[1], Candan Aykurt[2], Halil Süel[3], Ahmet Mert[1], Zeynep Öz[2], Ali Şenol[1]

[1]Faculty of Forestry, Isparta University of Applied Sciences, Isparta, Turkey

[2]Faculty of Arts and Sciences, Department of Biology, Akdeniz University, Antalya, Turkey

[3]Sütçüler Prof. Dr. Hasan Gürbüz Vocational School, Isparta University of Applied Sciences, Isparta, Turkey

serkangulsoy@isparta.edu.tr

Abstract

For accurate ecosystem-based management plans and strategies today and in the future, it is necessary for modeling and mapping of ecological characteristics of target species. In the present study, data from Kumluca district representing the Mediterranean ecosystem of Turkey were analyzed using two different modeling techniques to determine the potential distribution of *Styrax officinalis* L. According to the results of Logistic Regression Analysis (LRA), which is the first of the modeling techniques, precipitation, elevation, radiation index, heat index, solar radiation index, and topographic humidity index were the significant explanatory variables for the potential distribution of the species. The ROC values of the training and test dataset of the LRA model were found 0.915 and 0.874, respectively. Elevation, precipitation, and hillshade index were the variables that structured the model in the Classification Tree (CT) technique as the second method. The ROC values of the training and test dataset of the CT model were found 0.937 and 0.834, respectively. The common results of the two models revealed that the distribution of the species in Mediterranean ecosystems above 1500 m is generally limited. Below 1500 m, mainly precipitation and partially other model variables were statistically significant descriptors for the potential distribution areas of S. *officinalis* in the district.

Keywords: Classification tree; Ecological factors; Lojistic Regression Analysis; Precipitation

REFERENCES

1. Davis, P. H. (1978). *Styrax* L. in Flora of Turkey and the East Aegean Islands. Vol. 6. p. 144.
2. Jaradat, N. (2020). Phytochemistry, traditional uses and biological effects of the desert plant Styrax officinalis L. *Journal of Arid Environments*, *182*, 104253.
3. Yesilyurt, M. K., & Cesur, C. (2020). Biodiesel synthesis from Styrax officinalis L. seed oil as a novel and potential non-edible feedstock: a parametric optimization study through the Taguchi technique. *Fuel*, *265*, 117025.

On the Numerical Solutions of time Fractional FitzHugh Nagumo equation with Forced Term

Güllü Esra Köse, Yusuf Ucar and Alaattin Esen

Department of Mathematics, University of İnönü, Malatya, Turkey

esra_bubek@hotmail.com

Abstract

Neuronal behavior in terms of membrane potentials are modeled using some partial differential equations such as Hodgkin-Huxley model (1952), FitzHugh-Nagumo model (1969), Morris-Lecar model (1981), Hindmarsh-Rose model (1984). Among these models, FitzHugh-Nagumo model due to its mathematically simple nature and producing a rich dynamical behavior, has been so successful. The aim of this paper is two fold: firstly, to construct a finite element scheme to investigate the underlying fractional numerical behaviour the time fractional FitzHugh model with forced term and Robin boundary conditions. To support these numerical and theoretical observations, numerical example for the model and graphical behavior of the solutions are presented under the proposed fractional framework. Results indicate that the proposed method are very effective mathematical tools for solving reaction--diffusion systems.

Keywords: FitzHugh-Nagumo model, Collocation method, B-splines.

REFERENCES

1. Hodgkin AL, Huxley AF (1952) A quantitative description of membrane current and application to conduction to conduction and excitation in nerve. J Physiol 117: 500-544.
2. FitzHugh R. Impulses and Physiological States in Theoretical Models of Nerve Membrane. Biophysical Journal.1961;1(6):445-466.
3. Fritz, Marvin, Mabel L. Rajendran, and Barbara Wohlmuth. "Time-fractional Cahn--Hilliard equation: Well-posedness, degeneracy, and numerical solutions." Computers & Mathematics with Applications 108 (2022):66-87.
4. B. Jin, R. Lazarov, and Z. Zhou, An analysis of the L1 scheme for the subdiffusion equation with nonsmooth data, IMA J. of Numer. Anal., 36(2016), pp. 197-221
5. Tchier, F., Inc, M., Korpinar, Z. S., Baleanu, D. (2016). Solutions of the time fractional reaction-diffusion equations with residual power series method. Advances in Mechanical Engineering, 8(10),1687814016670867.
6. Kumar, Devendra, Jagdev Singh, and Dumitru Baleanu. "A new numerical algorithm for fractional Fitzhugh-Nagumo equation arising in transmission of nerve impulses." Nonlinear Dynamics 91.1 (2018): 307-317.
7. Srivastava, H. M., and Khaled M. Saad. "New approximate solution of the time-fractional Nagumo equation involving fractional integrals without singular kernel." Appl. Math. Inf. Sci 14.1 (2020): 1-8.

The Köthe Teoplitz Duals of $V_\lambda^0(G)$

Tuba Dinç and Çiğdem Bektaş

Department of Mathematics, University of Firat, Elazig, Turkey

karahan.tuba@gmail.com, cbektas@firat.edu.tr

Abstract

In this paper, we define the sequence spaces $V_\lambda^0(G)$, where $G = (g_k)$ is a sequence of modülüs functions. Also we will show that this space is a linear space and a solid AK-FK-space with an absolutely monotone F-norm. Furthermore we will introduce the The Köthe-Toeplitz Duals of $V_\lambda^0(G)$.

Keywords: Köthe-Toeplitz duals; AK-FK-spaces; Modülüs functions; Summability.

REFERENCES

1. Ahmad, Z. U.; Mursaleen, M. (1987). Köthe-Toeplitz duals of some new sequence spaces and their matrix maps. Publ. Inst. Math. (Beograd) (N.S.) 42(56), 57–61.

2. Altin, Y. (2009). Properties of some sets of sequences defined by a modulus function. Acta Math. Sci. Ser. B Engl. Ed. 29(2), 427–434.

3. Başarir, M. (2010). On the generalized Riesz B-difference sequence spaces. Filomat, 24(4), 35-52.

4. Bektaş, Ç. A. ; Et, M., & Çolak, R. (2004). Generalized difference sequence spaces and their dual spaces. Journal of Mathematical Analysis and Applications, 292(2), 423-432.

5. Bilgin, T. (2001). Lacunary strong A-convergence with respect to a modulus. Mathematica, 46(4), 39-46.

6. Chandra, P. ; Tripathy, B. C. (2002). On generalised Köthe-Toeplitz duals of some sequence spaces. Indian J. Pure Appl. Math. 33, no. 8, 1301–1306.

7. Connor, J. (1989). On strong matrix summability with respect to a modulus and statistical convergence. Canadian Mathematical Bulletin, 32(2), 194-198.

8. Cooke, R. G. (1955). Infinite Matrices and Sequence Spaces, Dover Publ.

9. Et, M. (1993). On some difference sequence spaces, Turk. J. Math. 17, 18-24.

APPLICATIONS OF COMPLEX FUZZY SOFT SETS IN RING THEORY

Gökhan Yüce[1], Yıldıray Çelik[2]

[1]Department of Mathematics, University of Ordu, Ordu, Turkey

gokhanyuce86@hotmail.com

[2]Department of Mathematics, University of Ordu, Ordu, Turkey

ycelik61@gmail.com

Abstract

In this paper, we extend the notion of ring to inside the algebraic structures of complex fuzzy soft sets. Then we introduce the concept of complex fuzzy soft rings (ideals) and study some of their properties and structural characteristics. We also give some theorems of homomorphic image and homomorphic pre-image of complex fuzzy soft rings under a complex fuzzy soft homomorphism.

Keywords: Complex fuzzy soft set; Complex fuzzy soft ring; Complex fuzzy soft ideal.

REFERENCES

1. M. Alsarahead, A. Ahmad, Complex Fuzzy Subring, International Journal of Pure and Applied Mathematics, Vol:117, 563-577, 2017.
2. Y. Çelik, C. Ekiz, S. Yamak, Applications of fuzzy soft sets in ring theory, Annals of Fuzzy Mathematics and Informatics, Vol:5, No:3, 451-462, 2013.
3. A. Nadia, Complex fuzzy soft set, MSc Research Project, Universiti Kebangsaan Malaysia, 2010.
4. D. Ramot, R. Milo, M. Friedman, A. Kandel, Complex fuzzy set, IEEE Transaction on Fuzzy Systems, Vol:10, 171-186, 2002.
5. L.A. Zadeh, Fuzzy sets, Information and Control, Vol:8, 338-353, 1965.

On the complex solutions to the Hirota–Satsuma–Ito equation

Hajar F. Ismael [1,2] **, Arif Özkul** [2] **, Hasan Bulut** [2]

Hasan Bulut [1]

[1] Department of Mathematics, University of Zakho, Zakho, Iraq

[2] Department of Mathematics, University of Firat, Elazig, Turkey

hajar.ismael@uoz.edu.krd,

Abstract

In this work, we study the Hirota–Satsuma–Ito equation that describes the propagation of unidirectional shallow-water waves and interactions of two long waves with different dispersion forms. For this study, we use the sine-Gordon expansion method for the suggested equation. Complex solutions in terms of dark-bright, dark and bright are constructed. Obtained solutions are constructed in three-dimensional figures to understand these physical phenomena.

Keywords: Complex solutions, Hirota–Satsuma–Ito equation, Dark, Bright, Dark-bright.

REFERENCES

1. Zhang, Y., Deng, S.F. and Chen, D.Y., The novel multi-soliton solutions of equation for shallow water waves, Journal of the Physical Society of Japan, 72(3), pp.763-764, 2003.
2. Hirota, R., The direct method in soliton theory, Cambridge University Press, 2004.
3. Lü, X., Tian, B., Sun, K. and Wang, P., Bell-polynomial manipulations on the Bäcklund transformations and Lax pairs for some soliton equations with one Tau-function, Journal of Mathematical Physics, 51(11), p.113506, 2010.
4. Lü, X., Ma, W.X., Chen, S.T. and Khalique, C.M., A note on rational solutions to a Hirota-Satsuma-like equation, Applied Mathematics Letters, 58, pp.13-18, 2016.
5. Wu, Y.Q., Asymptotic Behavior of Periodic Wave Solution to the Hirota—Satsuma Equation, Chinese Physics Letters, 28(6), p.060204, 2011.

A SECOND-ORDER DİFFERENCE SCHEME ON BAKHVALOV MESH FOR THE SİNGULARLY PERTURBED DELAY PSEUDO-PARABOLİC EQUATİONS

Baransel Gunes[1] **, Hakki Duru**[2]

1,2Department of Mathematics,Van Yuzuncu Yil University,Van,Turkey

baranselgunes23@gmail.com, hakkiduru@gmail.com

Abstract

This paper presents the finite difference scheme on Bakhvalov mesh for the singularly perturbed pseudo-parabolic problems with time-delay. Priori bounds of the solution is investigated by using the energy estimates. An error analysis is derived in the discrete maximum norm and the almost second order convergence is obtained. A numerical example is included to confirm the theoretical findings.

Keywords: Delay differential equation, difference scheme, error estimate, pseudo-parabolic problem, singular perturbation

REFERENCES

1. Amiraliyev, G. M., Mamedov, Y. D.: Difference schemes on the uniform mesh for singularly perturbed pseudo-parabolic equations. Turkish Journal of Mathematics. 19, 207-222 (1995).

2. Chinayeh A. B., Duru, H.: On adaptive mesh for the initial boundary value singularly perturbed delay Sobolev problems, Numerical MethodS for Partial Differential Equations (2019) doi:10.1002/num.22417.

3. Mohapatra J., Shakti D.: Numerical treatment for the solution of singularly perturbed pseudo-parabolic problem on an equidistributed grid. Nonlinear Engineering, 9, 169-174 (2020).

A ROBUST NUMERİCAL METHOD FOR THE SİNGULARLY PERTURBED SECOND-ORDER VOLTERRA-FREDHOLM INTEGRO-DİFFERENTİAL EQUATİONS

Baransel Gunes [1] , Musa Cakir[2]

1,2Department of Mathematics,Van Yuzuncu Yil University,Van,Turkey

baranselgunes23@gmail.com, cakirmusa@hotmail.com

Abstract

In this study, second-order Volterra-Fredholm integro-differential equations with layer behavior is investigated. Firstly, preliminary results for the analytical problem are given. Then, a uniform discretization is presented on layer-adapted mesh. The stability and convergence bounds of the proposed discretization are analyzed in the discrete maximum norm. Finally, numerical examples are given to clarify the theory.

Keywords: Error estimate, Finite difference scheme, Layer-adapted mesh, Singular perturbation.

REFERENCES

1. G. M. AMIRALIYEV, M. E. DURMAZ AND M. KUDU, Fitted second order numerical method for a singularly perturbed Fredholm integro-differential equations, Bull. Belg. Math. Soc. Simon Stevin, 27 (2020), pp. 71-88.

2. E. CIMEN AND M. CAKIR, A uniform numerical method for solving singularly perturbed Fredholm integro-differential problem, Computational and Applied Mathematics 40 (2021), https://doi.org/10.1007/s40314-021-01412-x.

3. N. A. MBROH, S. C. O. NOUTCHIE AND R. Y. M. MASSOUKOU, A second order finite difference scheme for singularly perturbed Volterra integro-differential equation, Alexandria Engineering Journal, 59 (2020), pp. 2441-2447.

AN APPLICATION OF INTUITIONISTIC NEUTROSOPHIC SOFT GRAPHS IN A DECISION-MAKING PROBLEM

Onur Zihni[1], **Yıldıray Çelik**[2]

[1]Department of Mathematics, University of Ordu, Ordu, Turkey

kuzeydenizli@gmail.com

[2]Department of Mathematics, University of Ordu, Ordu, Turkey

ycelik61@gmail.com

Abstract

We encounter countless decision-making problems in our daily lives. With the progress of science, many problems have progressed at the same speed. The universal language of science that produces solutions to such problems is mathematics. In order to find the most accurate solution, the way of thinking that leads us to the right decision is mathematical modeling. In this paper, firstly we introduce the concept of intuitionistic neutrosophic soft graphs and give some new operations on this concept. Further, we discuss a real-life application of intuitionistic neutrosophic soft graphs in decision-making. Lastly, for the selection of optimal object, we elaborate general procedure of our application by designing an algorithm.

Keywords: Intuitionistic neutrosophic soft set; Intuitionistic neutrosophic soft graph; Decision making problem.

ACKNOWLEDGMENTS

This work was supported by Ordu University Scientific Research Projects Coordination Unit in the scope of the project number B-1915.

REFERENCES

1. K. Atanassov, Intuitionistic fuzzy sets, Fuzzy Sets and Systems, Vol:20, 87-96, 1986.
2. P. Bhattacharya, Some Remarks on Fuzzy Graphs, Pattern Recognition Letters, Vol:6, No:5, 1987.
3. S. Broumi, F. Smarandache, Intuitionistic Neutrosophic Soft Set, Journal of Information and Computing Science, Vol:8, No:2, 130-140, 2013.
4. R. Parvathi, M.G. Karunambigai, Intuitionistic Fuzzy Graphs, Computational intelligence, theory and application, Springer, Berlin, Heidelberg, 139-150, 2006.

AN ALGORITHM FOR THE VERTICES ON THE PATHS OF MINIMAL LENGTH IN THE SUBORBITAL GRAPHS

Ümmügülsün AKBABA[1], **Ali Hikmet DEĞER** [1], **Tolga BERBER**[1]

[1] Department of Mathematics, University of Karadeniz Technical, Trabzon, Turkey

ummugulsun.akbaba@gmail.com,

Abstract

The suborbital graphs are formed by the imprimitive action of the modular group Γ on the rational projective line $\widetilde{\mathbb{Q}} = \mathbb{Q} \cup \{\infty\}$. Jones, Singerman and Wicks extend the results of Farey graph $\mathbf{F}_{1,1}$ to suborbital graphs $\mathbf{F}_{u,N}$, where $(u, N) = 1$ and $N > 1$. Then, Deger defined the farthest vertices on the paths of minimal length on the suborbital graphs and investigated the related continued fractions.

In this paper, we present an algorithm which gives the values of the vertices on the paths of minimal length in the suborbital graphs $\mathbf{F}_{u,N}$.

Keywords: Suborbital graphs, Imprimitive action, Continued fractions

REFERENCES

1. M. Akbas, On suborbital graphs for the modular group. Bulletin of the London Mathematical Society, 33, 647-152, 2001.
2. G.A. Jones, D. Singerman, K. Wicks, The modular group and generalized Farey graphs, London Math. Soc. Lecture Note Ser., 160, 316-338, 1991.
3. A.H. Deger, Vertices of paths of minimal lengths on suborbital graphs, Filomat, 31, 913-923, 2017.
4. A Cuyt, VB Petersen, B Verdonk, WB Jones, Handbook of Continued Fractions for Special Continued Fractions, New York: Springer, 2008.

Solitary Wave Solutions to NLEEs for Higher Balance Number through the Modified Simple Equation Method

[1]M. Ali Akbar, [2]Md. Ekramul Islam, [3]Fatma Berna Benli and [4]Onur Alp İlhan

[1]Department of Applied Mathematics, University of Rajshahi, Bangladesh

Email: ali_math74@yahoo.com

[2]Department of Mathematics, Pabna University of Science and Technology, Bangladesh

Email: ekramul.math@pust.ac.bd

[3,4]Department of Mathematics, Faculty of Education , Erciyes University, 38039

Melikgazi, Kayseri, Turkey

Email : [3] akpinarb@erciyes.edu.tr ; [4] oailhan@erciyes.edu.tr

Abstract: Although the modified simple equation (MSE) method effectively provides exact traveling wave solutions to nonlinear evolution equations (NLEEs) that emerge in engineering and mathematical physics, it has some drawbacks. When the balance number is greater than one, the approach typically fails to provide solution. We have addressed this shortcoming in this article and figured out a process to implement the MSE approach to investigate NLEEs for balancing number two. Two NLEEs, namely, the regularized long wave and the Jimbo-Miwa equations, have been investigated in order to affirm the approach. Through this scheme, we found further generic wave solutions related to physical parameters, and when the parameters receive particular values, solitons emerge from the exact solutions. Graphs are used to investigate the solitary wave properties of the attained solutions. This shows the validity, usefulness, and compatibility of the process.

Keywords: MSE method; nonlinear evolution equations; solitary wave solutions; exact solutions; regularized long wave equation; Jimbo Miwa equation

References

1. Ablowitz, M.J., Clarkson, P.A., Soliton, nonlinear evolution equations and inverse scattering, Cambridge University Press, New York, 1991.
2. Hirota, R., The direct method in soliton theory, Cambridge University Press, Cambridge, 2004.
3. Xin, X., Xia, Y., Liu, H., Zhang, L., Darboux transformation of the variable coefficient nonlocal equation, J. Math. Anal. Appl., 490 (2020): 124227.

New exact solutions for the Ivancevic option pricing model

Faruk Düşünceli [1]

[1] Department of Economics, Mardin ArtukluUniversity, Mardin, Turkey

farukdusunceli@artuklu.edu.tr,

Abstract

The Ivancevic option pricing model is presented by the application of the improved Bernoulli subequation function method. The equation which is a nonlinear partial differential equation is transformed into nonlinear ordinary differential equation using a wave transformation and then is solved by IBSEFM. Some new solutions are successfully constructed. Under suitable choice of the parameter values, interesting two- and three-dimensional graphs of all the obtained solutions are plotted.

Keywords: The Ivancevic option pricing model; the improved Bernoulli subequation function method; Exact solutions.

REFERENCES

1. Chen, Q., Baskonus, H. M., Gao, W., & Ilhan, E. (2022). Soliton theory and modulation instability analysis: The Ivancevic option pricing model in economy. *Alexandria Engineering Journal*, *61*(10), 7843-7851.
2. Faruk Düşünceli, Hacı Mehmet Baskonus, Alaattin Esen and Hasan Bulut, New mixed-dark soliton solutions to the hyperbolic generalization of the Burgers equation, Journal of Balıkesir University Insiıtute of Science and Technology, 21(2): 503-511, 2019.
3. Faruk Düşünceli New Exact Solutions for the (3 + 1) Dimensional B-type Kadomtsev-Petviashvili Equation. Erzincan Üniversitesi Fen Bilimleri Enstitüsü Dergisi, 12 (1), 463-468,(2019). DOI: 10.18185/erzifbed.493777
4. Pourghanbar, S., Manafian, J., Ranjbar, M., Aliyeva, A., & Gasimov, Y. S. (2020). An efficient alternating direction explicit method for solving a nonlinear partial differential equation. Mathematical Problems in Engineering, Volume 2020, Article ID 9647416, https://doi.org/10.1155/2020/9647416.
5. Srivastava, H.M., Iqbal, J., Arif, M., Khan, A., Gasimov, Y.S., & Chinram, R. (2021). A new application of Gauss quadrature method for solving systems of nonlinear equations. Symmetry, 13(3), 432.

On the partial (p,q)-differential equations

İnci Çakmak [1] **and Aynur Şahin** [2]

[1,2] Department of Mathematics, Sakarya University, 54050, Sakarya, Türkiye

incicakmak13113@gmail.com, ayuce@sakarya.edu.tr

Abstract

The quantum calculus (or recalled q-calculus) appeared as a connection between mathematics and physics. It has many applications in different mathematical areas, such as number theory, combinatorics, orthogonal polynomials, and other sciences: quantum theory, mechanics, and theory of relativity. Further, there is the possibility of extension of the q-calculus to post-quantum calculus denoted by the (p,q)-calculus. When the case $p=1$, the (p,q)-calculus reduces to the q-calculus. Recently, Jafari et al. (Rom. Journ. Phys. 59, 399-407, 2014) presented the reduced q-differential transform method for solving partial q-differential equations. In this study, we define the concept of partial (p,q)-derivative for a multivariable function and generalize the method of Jafari et al. to partial (p,q)-differential equations. Also, we give some examples to discover the effectiveness and performance of the proposed method.

Keywords: q-calculus; (p,q)-calculus; Partial differential equation; Initial value problem.

References

1. Jafari, H., Haghbin, A., Hesam, S., Baleanu, D., Solving partial q-differential equations within reduced q-differential transform method, Rom. Journ. Phys., Vol: 59, No: 5-6, 399-407, 2014.
2. Sadjang, P. N., On the fundamental theorem of (p,q)-calculus and some (p,q)-Taylor formulas, Results Math., 73:39, 2018.
3. Aracı, S., Duran, U., Açıkgöz, M., Srivastava, H. M., A certain (p,q)-derivative operator and associated divided differences, J. Ineq. Appl., 2016:301, 2016.
4. El-Shahed, M., Gaber, M., Two-dimensional q-differential transformation and its application, App. Math. Comput., Vol:217, 9165-9172, 2011.
5. Akça, H., Benbourenane J., Eleuch, H., The q-derivative and differential equation, Journal of Physics: Conference Series, 1411:012002, 2019.

Some special Smarandache Ruled Surfaces according to Frenet Frame in E^3-II

[1]Süleyman ŞENYURT, [2]Davut CANLI, [3]Elif ÇAN
[1,2,3]Faculty of Arts and Sciences, Department of Mathematics,Ordu University, Ordu,

TURKEY

[1]ssenyurt@odu.edu.tr, [2]davut_canli@hotmail.com, [3]eliff.cann@hotmail.com

Abstract.
In this study, firstly, the Smarandache curve was defined using Frenet vectors of a curve. Along this curve, the Gaussian and average curvatures of the ruled surfaces formed by the vectors obtained from the Frenet vectors are calculated, and the conditions of being openable or minimal are given. Finally, examples were given for each surface and drawings were made with the maple 17 program.

Key Words: Smarandache ruled surfaces, mean curvature, Gaussian curvature,vivian cuve

References

[1] Ali, A.T., Special Smarandache curves in the Euclidean space, International Journal of Mathematical Combinatorics, 2(2010),30-36, 2010

[2]Berk, A., A Structural Basis for Surface Discretization of Free Form Structures: Integration of Geometry, Materials and Fabrication, Ph. D Thesis, Michigan University, ABD, 2012

[3]Do-Carmo,P.M., Differential geometry of curvesand surfaces. IMPA, 1976

[4]Fenchel,W., On the Differential Geometry of Closed Space Curves, Bulletin of American Mathematical Society, 57(44-54), 1951

[5]Gray A. Modern differential geometry of curves and surfaces with Mathematica, 1997

[6]Ouarab, S., Smarandache Ruled Surfaces according to Frenet-Serret Frame of a Regular Curve in E3, Hindawi Abstract and Applied Analysis, Vol. 2021, Article ID 5526536, 8 pages

[7]Ouarab, S., Smarandache ruled Surfaces according to Darboux Frame in E^3, Hindawi Journal of Mathematics, Vol. 2021, Article ID 9912624, 10 pages

[8]Ouarab,S., NC-Smarandache ruled surface and NW-Smarandache ruled surface according to alternative moving frame in E^3, HindawiJournal of Mathematics ,Vol. 2021, Article ID 9951434, 6 pages

[9]Pottmann, H., Asperl, A., Hofer, M. ve Killian, A., Architectural Geometry, Bentley, D. (ed.), First Edition, Bentley Institute Press, 2007

[10]Pressley A.,Elementary Differential Geometry, Second Edition, Springer London, 201

PELL AND PELL-LUCAS NUMBERS IN SUBORBITAL GRAPHS

Ümmügülsün AKBABA[1], **Ali Hikmet DEĞER** [1]

[1] Department of Mathematics, University of Karadeniz Technical, Trabzon, Turkey

ummugulsun.akbaba@gmail.com,

Abstract

Continued fractions and their matrix connections have been used in many studies to generate new identities. On the other hand, many examinations have been made in the suborbital graphs under circuit and forest conditions. Special number sequences and special vertex values of minimal length paths in suborbital graphs have been associated in our previous studies. In this study, some special natural number values of $k > 2$ are used for the continued fractions $\boldsymbol{K}(-1/k)$, which is a special periodic continued fraction. The relation between Pell, Pell- Lucas numbers and the suborbital graphs are examined by using this continued fraction. Also, some new identities and some new matrices related to Pell, and Pell-Lucas number sequences are produced.

Keywords: Pell numbers, Pell-Lucas numbers, Continued fractions, Suborbital graphs

REFERENCES

1. M. Akbas, On suborbital graphs for the modular group. Bulletin of the London Mathematical Society, 33, 647-152, 2001.
2. G.A. Jones, D. Singerman, K. Wicks, The modular group and generalized Farey graphs, London Math. Soc. Lecture Note Ser., 160, 316-338, 1991.
3. A.H. Deger, Vertices of paths of minimal lengths on suborbital graphs, Filomat, 31, 913-923, 2017.
4. A Cuyt, VB Petersen, B Verdonk, WB Jones, Handbook of Continued Fractions for Special Continued Fractions, New York: Springer, 2008.
5. T Koshy, Pell, and Pell-Lucas numbers with applications. Springer. New York,2014.

SOME RELATIONS BETWEEN THE SETS OF $\Delta_v^m(f)$-STATİSTİCAL CONVERGENCE OF ORDER α

Emine ÖZÇELİK and Çiğdem A. BEKTAŞ

Department of Mathematics, University of Firat, Elazig, Turkey

eminemozcelik@gmail.com , cbektas@firat.edu.tr

Abstract. In this article we introduce the relationship between $\Delta_v^m(f)$-statistically convergent and $\Delta_v^m(f)$-statistically bounded of order α .

Mathematics subject classification. 40A35 . 40G15 . 40A05 . 46A45

Keywords and phrases. Density . Modulus functıon . Statistical convergence . Statistical boundedness . Difference sequence

REFERENCES

[1] Aizpuru A, Listån-Garcìa MC, Rambla-Barreno F, Density by moduli and statistical convergence, Questiones Mathematicae Quaestiones Mathematicae, Vol:37 No:4, 525-530, 2014, DOI:10.2989/16073606.2014.981683

[2] Bhardwaj VK, Singh N, On some sequence spaces defined by a modulus, Indian J Pure Appl Math Vol:30, No: 8, 809-817, 1999.

[3] Bhardwaj VK, Dhawan S, and Gupta S (2016) Density by Moduli and Statistical Boundedness, Abstract and Applied Analysis Volume 2016, Article ID 2143018, 6 pages, 2016 http://dx.doi.org/10.1155/2016/2143018

[4] Connor JS, The Statistical and Strong p-Cesåro Convergence of Sequences, Analysis Vol:8, 47-63, 1988.

[5] Fast H, Sur la convergence statistique Colloq Math 2, 241-244, 1951.

[6] Fridy J, On statistical convergence, Analysis 5, 301-313, 1985.

[7] Fridy JA and Orhan C, (1997) Statistical limit superior and limit inferior, Proc. Amer. Math. Soc. Vol:125 No:12, 3625-3631, 1997.

DETERMINING THE SUSTAINABLE CONSUMPTION BEHAVIORS OF TEACHER CANDIDATES

Gökce İrem Kızılkaya[1]* – Asist. Prof. Dr. Elif Çil[2] – Assoc. Prof. Dr. Sevda Türkiş[3]

Department of Mathematics and Science Education, University of Ordu, Ordu, Turkey

gokceiremkizilkaya@odu.edu.tr, elifcil@odu.edu.tr, sevdaturkis@odu.edu.tr

Abstract

Consumption is defined in the dictionary as the use and consumption of things produced. Consuming is carried out for different purposes, from fulfilling the vital activities of living things to meeting essential needs. It is known that this action is affected by many factors such as age, gender, lifestyle, occupation, and environmental awareness. Obviously, with the unconscious increase in consumption, there will be problems that are difficult to solve in the world. The concept of sustainable consumption has a critical position in this sense. Sustainable consumption behavior is a form of action that aims to make the present and the future livable by ensuring that the consumer mass is conscious while performing the consumption act. For life to be sustainable, it is essential to give the proper social consumption awareness to manage world resources. In raising conscious consumers, the role of teachers is at least as necessary as that of the family. Therefore, this study it is aimed to determine the sustainable consumption behaviors of science and classroom teacher candidates. The study sample, which was carried out with the descriptive survey model from quantitative research, consists of pre-service teachers who continue their education in science and classroom teaching departments at a state university in North Anatolia. The study consists of 154 persons. The "Sustainable Consumption Behaviors Scale developed by Doğan et al. (2015)" was used as a data collection tool. The data analysis was carried out with the SPSS 21.0 package program.

Keywords: Science teacher candidate, Primary school teacher candidate, Quantitative Analysis

REFERENCES

1. Doğan, O., Bulut, Z. A., & Çımrın, F. K. (2015). A Scale Development Study To Measure Indıvıduals' Sustaınable Consumptıon Behavıor. *Ataturk University Journal Of Economics & Administrative Sciences*, *29*(4).
2. Şahin, B. G., & Akballı, E. E. (2019). Tüketici Davranışlarını Etkileyen Faktörler Ve Yöntem Analizi. *Uluslararası Sosyal Bilimler Ve Eğitim Dergisi*, *1*(1), 43-85.

AN EIGENVALUE PROBLEM WITH A SPECTRAL PARAMETER IN BOUNDARY CONDITIONS

Ayşe Kabataş [1]

[1] Department of Mathematics, Karadeniz Technical University, Trabzon, Turkey

akabatas@ktu.edu.tr,

Abstract

In this paper, we improve asymptotic estimates of eigenfunctions for an eigenvalue problem considered by Kerimov and Mamedov (1999) with the spectral parameter in all boundary conditions when the potential function is continuous, also its differentiation exists and is integrable.

Keywords: Eigenvalue problem; Eigenfunctions; Asymptotics.

REFERENCES

1. Charles T. Fulton, Two-Point Boundary Value Problems with Eigenvalue Parameter Contained in the Boundary Conditions, Proceedings of the Royal Society of Edinburgh Section A: Mathematics, Vol:77, Issue:3-4, 293-308, 1977.
2. Elif Başkaya, Asymptotic of Eigenvalues for Sturm-Liouville Problem with Eigenparameter-Dependent Boundary Conditions, New Trends in Mathematical Sciences, Vol:6, No:2, 247-257, 2018.
3. B. J. Harris, The Form of the Spectral Functions Associated with Sturm-Liouville Problems with Continuous Spectrum, Mathematika, Vol:44, 149-161, 1997.
4. Haskız Coşkun, Ayşe Kabataş, Asymptotic Approximations of Eigenfunctions for Regular Sturm-Liouville Problems with Eigenvalue Parameter in the Boundary Condition for Integrable Potential, Mathematica Scandinavica, Vol:113, No:1, 143-160, 2013.
5. N. B. Kerimov, Kh. R. Mamedov, On One Boundary Value Problem with A Spectral Parameter in the Boundary Conditions, Vol:40, No:2, 281-290, 1999.

A NEW LOOK AT : THE NUMERİCAL SOLUTİON OF THE GENERALİZED BURGERS–HUXLEY EQUATİON

Melike Karta [1]

Department of Mathematics, Ağrı ibrahim Çeçen University, Ağrı, Turkey

mkarta@agri.edu.tr

Abstract

In this study, a new technique for the numerical approximation of the generalized Burgers–Huxley equation is recommended using Strang time splitting technique which divides the problem into two sub-equations suh that each of them transform a system of partial differential equations. After, each partial differential equation is reduced a system of first-order ordinary differential equations using cubic B-spline collocation method for spatial derivatives. Afterwards, the resulting ODEs for time derivatives are solved using the Strang time splitting technique via Runge–Kutta (RK-4) approach. Our aim here is to measure how effective the technique we applied to the generalized Burgers–Huxley equation is. For this, L_∞ error norm has been calculated by choosing small enough to the time step length used by different researchers and the solutions acquired have been compared with the solutions of some researchers in the literature. It can be seen that the technique used is useful and trustworthy and easily applied to other diffusion–reaction equations and said that the obtained scheme is not only simple but also effective and easy in terms of computability.

Keywords: Generalized Burgers–Huxley equation; Cubic B-splines; RK4 approach; Collocation method

REFERENCES

1. Ramesh Chand Mittal, Amit Tripathi, Numerical solutions of generalized Burgers–Fisher and generalized Burgers–Huxley equations using collocation of cubic B-splines, International Jurnal of Computer Mathematics, Vol:92, No:5, 1053-1077, 2015.

2 Mohammad Javidi, Ahmad Golbabai, A new domain decomposition algorithm for generalized Burger's–Huxley equation based on Chebyshev polynomials and preconditioning, Chaos, Solitons and Fractals, Vol:39, No:2, 849–857, 2009.

3. Yali Duan, Linghua Kong, Rui Zhang, A lattice Boltzmann model for the generalized Burgers–Huxley equation. Phys. A., Vol:391, No:3, 625–632, 2012.

4. Kolade M. Owolabi, Numerical Solution of the Generalized Burgers-Huxley Equation by Exponential Time Differencing Scheme, International Journal of Bioinformatics and Biomedical Engineering, Vol:1, No:1, 43-52, 2015.

5. Murat Sari ,Gürhan Gürarslan, Numerical Solutions of the Generalized Burgers-Huxley Equation by a Differential Quadrature Method, Hindawi Publishing Corporation Mathematical Problems in Engineering, Vol: 2009, No:11, Article ID 370765, pages doi:10.1155/2009/370765.

6. Rongpei Zhang, Xijun Yu, Guozhong Zhao, The local discontinuous Galerkin method for Burger's–Huxley and Burger's–Fisher equations, Applied Mathematics and Computation, 218 (2012): 8773–8778. at ht://dx.doi.org/10.1016/j.amc.2012. 02.035.

AN INVERSE RECONSTRUCTION PROBLEM

Mehmet Açil

Department of Mathematics, Van Yüzüncü Yıl University, Van, Türkiye

mehmetacil@yyu.edu.tr,

Abstract

In this paper, a method is proposed in order to calculate the potential function in Sturm-Liouville differential equation in the normal form with Dirichlet boundary conditions by using some sets of partial data. One of these data is defined in this paper as the other is a data that have been studied. To introduce the method, we give some lemmas and theorems. Then, we give numerical results for some test potentials to show the efficiency of the method.

Keywords: Inverse problems involving ordinary differential equations; Boundary eigenvalue problems for ordinary differential equations;

REFERENCES

1. O. H. Hald, J. R. McLaughlin, Solution of inverse nodal problem, Inverse Problems, Vol:5, 307–347, 1989.

2. A. Neamaty, Sh. Akbarpoor, Numerical solution of inverse nodal problem with an eigenvalue in the boundary condition, Inverse Problems in Science and Engineering, Vol: 25, 978-994, 2016

3. J. R. McLaughlin, W. Rundell, A Uniqueness Theorem for an Inverse Sturm-Liouville Problem, J. Math. Phys., Vol: 28, 1471–1472, 1987.

4. K. Chadan, D. Colton, L. Paivarinta, W. Rundell, An Introduction to Inverse Scattering and Inverse Spectral Problem, SIAM Monographs on Mathematical modeling and computation, SIAM, Philadelphia, 1997.

BIENERGIES FOR SYMPLECTIC REGULAR CURVE WITH SYMPLECTIC FRAME

Mehmet BEKTAŞ[1] , Esra ÇİÇEK ÇETİN, Sümeyye GÜR MAZLUM[2]

1 Department of Mathematic, Faculty of Science, University of Firat Elazığ, TURKEY.

2 Department of Computer Technology Gümüşhane University, Gümüşhane, TURKEY.

Abstract: In this work , we research geometrical interpration involved with the bienergy for symplectic regular curve with symplectic frame.

Keywords: Frenet vectors, Symplectic curve, energy functions.

REFERENCES

[1] T. Körpınar, Y.Ünlütürk, A new wersion of bienergies and biangles for curve framed by extended darboux frame, Soft Computing (2022), 26, 45-54.

[2] T.Körpınar and R.C. Demirkol , Curvature and Torsion Dependent energy of Elastica and Nonelastica for a light-like curve in the Minkowski Space, Ukr. Math.J. Vol 72 No:8 2021, 1267-1279.

[3] E.Ç.Çetin, M.Bektaş, The characterizations of Affine symplectic curves in R^4, Mathematics mdpı 2019, 7(1), 110.

[4] E.Ç.Çetin, M.Bektaş, Some new characterizations of symplectic curve in 4- dimensional symplectic space, Communications in advanced mathematical Sciences , 20192(4), 331-334.

[5] E.Ç.Çetin, M.Bektaş, k-type slant helices for symplectic curve in 4-dimensional symplectic space, Facta Universitatis, Series: Mathematics and Informatics, 2020,641-646.

[6] N. Kamran, P. Olver, and K. Tenenblat,KTGĞ*J*UUJPLKMğk ÜĞÜÜÜÜÜŞNŞÇÇİN L Local symplectic invariants for curves, Commun. Contemp. Math., 11(2): (2009)., 165-183.

[7] F. Valiquette, Geometric affine symplectic curve flows in R^4, Differ. Geom. Appl., 30(6), (2012), 631-641.

[8] S.S Chern, H.S. Wang, Differential Geometry in symplectic space, I. Sci.Rep. Nat.Tsing Hua Univ., 4(1947), 453-477.

[9] V. Decochy, Hypersurfaces in symplectic affine geometry, Differ. Appl., 17(1) (2002), 1-13.

SOME NEW ASSOCIATED CURVES OF A NULL AND PSEUDO NULL CURVE IN 4-DIMENSIONAL MINKOWSKI SPACE-TIME

Esra ÇİÇEK ÇETİN[1], Mehmet BEKTAŞ[1]

[1] Department of Mathematics, Faculty of Science, University of Firat, Elazig, Turkey

esracicek@gmail.com, mbektas@firat.edu.tr

Abstract: In this paper on curves, we have designated Principal direction and Binormal direction curves of a granted Null curve and Pseudo Null curve by employing integral curves designated 4-dimensional Minkowski Space-time. Besides, give some characterizations of these curves are granted together with the intercourses between the curvatures of each related curve and the situation of Null and Pseudo Null curves to be helical with respect to each other was examinated.

Keywords: Minkowski Space-time, Principal direction curve, Binormal direction curve, Null curve, Pseudo Null curve, slant helix, B_2- slant helix

References

[1] A., Kazan, B., H., Karadağ, Magnetic Pseudo Null and Magnetic Null curves in Minkowski 3-space, İnternational Mathematical forum, 12:3, 119-132, (2017)

[2] E., Nesovic, U., Öztürk, B.,E, Koç, Öztürk, On k-type Pseudo Null Darboux helices in minkowski 3-space, Journal of mathematical analysis and applications, 439:2,690-700,(2016)

[3] H.,Öztekin, S., Tarla, Rectifying curves in the equiform geometry of the Galilean 4-space, The fourth international conference on computational mathematics and engineering sciences, 653-662, (2019)

[4] K., Ilarslan, E., Nesovic, Some characterizations of null, pseudo null and partially null rectifying curves in Minkowski space-time, Taiwanese journal of mathematics,12:5,1035-1044, August (2008)

[5] L., Kula, N., Ekmekçi, Y., Yaylı, K., İlarslan, Charecterizations of slant helices in Euclidean 3-space, Turkısh journal of mathematics, 34:261-273, (2010)

[6] M., Babaarslan, Y., A., Tandoğan, Y., Yaylı, A note on Bertrand curves and constant slope surfaces according to Darboux Frame, journal of advanced mathematical studies, 5:87-96, (2012)

[7] M., Grbovic, K., Ilarslan, and E., Nesovic, On Null and Pseudo Null Mannheim curves in Minkowski 3-space, J. Geom. 105:177-183, (2014)

[8] N., Macit, M., Akbıyık, S., Yüce, Some new associated curves of an Admissible Frenet curve in 3-dimensional and 4- dimensional Galilean spaces, Romanian Journal of mathematics and computer science, 7(2), 110-122, 2017.

[9] W. B. Bonnor, Curves with null normals in minkowski space-time, a random walk in relativily and consmology, Wiley Easten Limitid, 33-47,(1985)

BİSECTOR CURVES OF COMFORMABLE CURVES IN $\mathbb{R}^2$

Şeyda ÖZEL[1], Mehmet BEKTAŞ[1]

[1] Deparment of Mathematics, Faculty of Science, University of Fırat,Elazığ, Turkey

s_demir2323@outlook.com, mbektas@firat.edu.tr

Abstract: In this study, the bisector curves of two regular comformable curves from C^1-regular parametric category is inspected in $\mathbb{R}^2$. Then, multivariable function which is corresponded to bisector curves of regular comformable curves is calculated. The bisector curves are procured by two different paths. As a result, the equations which are corresponded to bisector curves are obtained in $\mathbb{R}^2$.

Keywords. Bisector curve, Comformable Derivative, Frenet Frame, Bivariate Functions.

References

[1] K., Nishimoto, An Essence of Nishimoto's Fractional Calculus, Integrals and Differentiations of Arbitrary Order, (1991) 1-11

[2] R., Khalil, M., Al Horani, A., Yousef, M., Sababheh, A new definition of fractional derivative, Journal of Computational and Applied Mathematics, 264 (2014), 65–70

[3] A., Atangana, D., Baleanu, A., Alsaedi, New properties of conformable derivative, Open Mathematics, 13 (2015), 889–898

[4] D.R., Anderson, D.J., Ulness, Newly defined conformable derivatives, Advances in dynamical systems and applications, 10(2) (2015), 109–137

[5] H., Aminikhah, A.R., Sheikhani, H., Rezazadeh, Sub-equation method for the fractional regularized long-wave equations with conformable fractional derivatives, Scientia Iranica, 23(3) (2016), 1048–1054

[6]] N.Y., Gözütok, U., Gözütok, Multivariable Conformable Fractional Calculus, Filomat, 32(1) (2018), 45–53

[7] M., Dede, Y., Ünlütürk, C., Ekici, Bisector curves of planar rational curves in Lorantzian plane, International Journal of Geometry, 2(1), 2013

[8] U., Gözütok, H., Çoban, Y., Sağıroğlu, Frenet frame with respect to conformable derivative, Filomat., 33 (6), (2019)

ON THE CHARACTERİZATİONS OF CURVES WİTH MODIFIED ORTHOGONAL FRAME IN EUCLIDEAN 3-SPACE

Şeyda ÖZEL[1], Mehmet BEKTAŞ[1]

[1] Deparment of Mathematics, Faculty of Science, University of Fırat,Elazığ, Turkey

s_demir2323@outlook.com, mbektas@firat.edu.tr

Abstract: In this paper, we study (k,m) −type slant helices according to the modified orthogonal frame in Euclidean 3-space.

Key Words: Helices, Slant Helices, Euclidean 3-space.

References

[1] M.S., Lone, H., Es, M. K., Karacan, B., Bükcü, Mannheim curves with modified orthogonal frame in Euclidean 3-space, arXiv preprint arXiv:1809.10876, (2018)

[2] M.S., Lone, H., Es, M.K., Karacan, B., Bükcü, On some curves with modified orthogonal frame in Euclidean 3-Space, Iranian Journal of Science and Technology, Transactions A: Science, 43(4)2019: 1905-1916

[3] AT., Ali, R., Lopez, Slant helices in Minkowski space E_1^3, J Korean Math Soc 48:159–167, (2019)

[4] T., Sasai, The fundamental theorem of analytic space curves and apparent singularities of Fuchsian diferential equations, Tohoku Math J 36:17–24, (1984)

[5] L., Kula, Y., Yaylı, On slant helix and its spherical indicatrix, Appl Math Comput 169(1):600–607, (2005)

[6] W.K., Schief, On the integrability of Bertrand curves and Razzaboni surfaces, J Geom Phys 45:130–150, (2003)

[7] L., Kula, N., Ekmekçi, Y., Yaylı, K., Ilarslan, Characterizations of slant helices in Euclidean 3-space, Turk J Math 34:261–273, (2010)

[8] A.O., Ogrenmis, M., Ergut, M., Bektas, On the helices in the Galilean space G_3, Iran J Sci Technol Trans A 31(A2):177–181, (2007)

[9] M., Barros, General helices and a theorem of Lancret, Proc Am Math Soc 125(5):1503–1509, (1997)

[10] J., Monterde, Salkowski curves revisted: A family of curves with constant curvature and non-constant torsion, Comput Aided Geom Des 26:271–27, (2009)

Novel Numerical solutions of Time fractional Korteweg-de Vries-Burgers Equation

Berat Karaagac [1], **Nuri Murat Yagmurlu** [2], **Alaattin Esen** [2]

[1] Department of Mathematics Education, University of Adıyaman, Adıyaman, Turkey

[2] Department of Mathematics, University of Inonu, Malatya, Turkey

bkaraagac@adiyaman.edu.tr,

murat.yagmurlu@inonu.edu.tr,

alaattin.esen@inonu.edu.tr.

Abstract

The current paper uses the well-known and efficient numerical technique, known as Finite element method, to carry out numerical solutions of time fractional Korteweg–de Vries–Burgers equation. For the discretization of fractional time derivative, the $L1$ algorithm is applied, and for the spatial discretization Collocation method which is a finite element method to approximate the exact solution with trigonometric Quintic B-spline is used. The numerical results obtained using the method are presented via tables and graphics. The novel results demonstrate the efficiency and reliability of the method.

Keywords: Time Fractional Korteweg–de Vries–Burgers equation, Collocation method; Trigonometric Quintic B-splines.

REFERENCES

1. Santanu Saha Ray, Arun Kumar Gupta, Wavelet methods for solving partial differential equations and fractional differential equations. CRC Press, 2018.
2. Adel R., Hadhoud, Abdulqawi AM Rageh, Taha Radwan, Computational Solution of the Time-Fractional Schrödinger Equation by Using Trigonometric B-Spline Collocation Method, Fractal and Fractional, vol:6, No:127, 1-16, 2022.
3. A. K. Gupta, and S. Saha Ray, On the solution of time-fractional KdV–Burgers equation using Petrov–Galerkin method for propagation of long wave in shallow water, Chaos, Solitons & Fractals, vol: 116 , 376-380, 2018.

ON SOME FIXED POINT THEOREMS IN QUASI-METRIC SPACES AND ITS APPLICATIONS

İrem Eroğlu [1]

[1] Department of Mathematics, University of Ordu, Ordu, Turkey

iremeroglu@odu.edu.tr,

Abstract

In this paper we extend the Nieto fixed point results to the asymmetric context and we give some fixed point results in partially ordered quasi-metric spaces. We show that our results can not be weakened. Finally, we give an application of our results to the complexity analysis of algorithms.

Keywords: quasi-metric, bicomplete, partial order, contraction, monotone map, fixed point..

REFERENCES

1. S.Romaguera, P. Tirado, O. Valero, New results on the mathematical foundations of asymptotic complexity analysis of algorithms via complexity spaces, Int.J.Comput.Math. 89 (2012), 1728-1741.
2. A. Baranga, The contraction principle as a particular case of Kleene's fixed point theorem, Discrete Math. 98 (1991), 75-79.
3. S. Carl, S. Heikkila, Fixed Point Theory in Ordered Sets and Applications: From Diferential and Integral Equations to Game Theory, Springer, 2011.
4. B.A. Davey, H.A. Priestley, Introduction to Lattices and Order, Cambridge University Press, Cambridge, 1990.
5. H. P. A. Kunzi, Nonsymmetric topology, in: Proc. Colloquium on topology, 1993, Szekszard, Hungary, Colloq. Math. Soc. Janos Bolyai Math. Studies, 4 (1995), 303-338.
6. J.J. Nieto, R. Rodriguez-Lopez, Contractive mapping theorems in partially ordered sets and applications to ordinary diferential equations, Order 22 (2005), 223-239.

AN APPLICATION ON EARTHQUAKE DATA UNDER NON-RESPONSE SCHEME IN SAMPLING THEORY

Ceren Unal [1], **Gamze Ozel** [2]

[1,2] Department of Statistics, Hacettepe University, Ankara, Turkey

cerenunal@hacettepe.edu.tr, gamzeozl@hacettepe.edu.tr,

Abstract

While working on the population, the difficult situation of reaching all units is sometimes even impossible, as well as financial, time, manpower, etc. There are also restrictions. For this reason, the sample that can represent the population is used instead of the population. At this point, the samples are determined with the most appropriate sampling method, an estimation is made for the population parameters (mean, variance, ratio, etc.) by using these samples. The aim is to estimate the population parameter of the study with the most efficient estimator among existing estimators. For this reason, the most efficient estimator is tried to be proposed by using or producing many different methods in the literature. In such studies, it is assumed that all information is obtained for the variable of study (y) and auxiliary variable (x) in the estimators. However, this will not always be the case when considering the situations that may be encountered in daily life. Therefore, with a new approach developed by Hansen and Hurwitz (1946), estimators have started to be proposed by taking into account the non-response situation.

For the purpose of the study, an estimator was proposed in case of non-response in both the variable of study and the auxiliary variable. Theoretical inferences about the proposed estimator are supported by a numerical example. At this stage, the magnitude which are measured as 4 and larger that occurred and depth of the earthquakes in the Aegean Region were handled in the form of the variable of study and auxiliary variable, respectively, and analyzes were made, and the results were discussed in detail.

Keywords: Non-Response Scheme; Efficiency; Exponential Estimator.

REFERENCES

1. Hansen, M. H., Hurwitz, W. N. The problem of non-response in sample surveys. Journal of the American Statistical Association, 41(236), 517- 529, 1946.
2. Cochran, W.G. Sampling Techniques, John Wiley and Sons, New-York, 1977.
3. Yadav, S. K., Subramani, J., Misra, S., Singh, L., Mishra, S. S. Improved estimation of population mean in presence of non-response using exponential estimator. Int. J. Agricult. Stat. Sci. Vol, 12(1), 271-276, 2016.

A DIFFERENT APPROACH TO THE ASSESSMENT OF PARTICULATE MATTER (PM10) POLLUTION IN ANKARA

Derya Turfan [1]

[1] Department of Statistics, Hacettepe University, Ankara, Turkey

deryaturfan@hacettepe.edu.tr

Abstract

Air pollution is one of the most critical factors affecting the quality of life. Especially, particulate matter (PM) in the air poses a serious threat to human health. These particulates have different sizes. In this study, dust particles (PM10) smaller than 10 micrometers in diameter, which is one of the most harmful pollutants, are studied. The air quality of metropolitan cities should be measured and evaluated frequently, considering the number of people living. For this purpose, there are different stations where air pollution measurements are made in Ankara, the capital city of Turkey. This study aims for Ankara's mean air pollution rate to be calculated by applying the minimum spanning tree (MST) method, with each node representing a station. This ratio makes it possible for Ankara's overall air pollution assessment in terms of PM10.

Keywords: Minimum spanning tree; Air pollution; Particulate matter.

REFERENCES

1. Nevin Güler, Öznur Güneri İşçi, The Regional Prediction Model of PM10 Concentrations for Turkey, Atmospheric Research Vol: 180, 64-77, 2016.
2. Paulene Govender, Venkataraman Sivakumar, Application of K-Means And Hierarchical Clustering Techniques for Analysis of Air Pollution: A Review (1980–2019), Atmospheric Pollution Research, Vol:11, No:1, 40-56, 2020.
3. Hüseyin Toros, Hakan Erdun, Özkan Çapraz, Betül Özer, Esra Bozyazı Daylan, Ali İhsan Öztürk, Air Pollution and Quality Levels in Metropolitans of Turkey for Sustainable Life, European Journal of Science and Technology, Vol:1, No:1, 12-18, 2013.
4. Weihai Xu, Wen Yan, Gan Zhang, Jun Li, Li Miao, Weixia Huang, Levels and Distribution of Heavy Metals in Atmospheric Particular Matters Over the Northern South China Sea, Environmental Science and Pollution Research, Vol: 21, 8774-8781, 2014.

COMPARISON OF AEGEAN REGION MAXIMUM EARTHQUAKE MAGNITUDES WITH TIME SERIES METHODS

Hatice Oncel Cekim [1], Gamze Ozel Kadilar [1], Tuba Eroğlu Azak [2]

[1] Department of Statistics, Hacettepe University, Ankara, Turkey

[2] Department of Civil Engineering, National Defence University, Istanbul, Turkey

oncelhatice@hacettepe.edu.tr, gamzeozl@hacettepe.edu.tr, tuba.eroglu@gmail.com

Abstract

Earthquakes are one of the natural disasters that cause significant damage to people and nature. To reduce this damage, many studies have been carried out on earthquakes. However, earthquakes have a complicated structure to adapt to a model. The developing technology is used for mathematical-based analysis, data mining techniques, and earthquake predictions with artificial neural networks. The Aegean region constitutes one of the crucial areas in Turkey in terms of active tectonics. It is an important region, geologically and active tectonically, and should be investigated. Considering the AFAD-DDA catalog as the main catalog for this region, the repeated and missing earthquakes in the Tan and KOERI catalogs are arranged according to the Gardner and Knopoff (1974) method. Since the magnitudes of the earthquakes are related to the damages they cause, the maximum earthquake magnitude estimates are obtained for the first time with time-dependent classical and artificial neural network methods and the results are compared.

Keywords: Time series method; artificial neural network methods; earthquake magnitude estimates.

REFERENCES

1. Gardner, J. K., Knopoff, L., Is the sequence of earthquakes in Southern California, with aftershocks removed, Poissonian?, Bulletin of Seismological Society of America, Vol: 64, No: 5, 1363-1367. 1974.
2. Wang, Q., Guo, Y., Yu, L., Li, P., Earthquake prediction based on spatio-temporal data mining: an LSTM network approach", IEEE Transactions on Emerging Topics in Computing, Vol: 8, No:1, 148-158, 2017.
3. Yegulalp, T. M., Kuo, J. T., Statistical prediction of the occurrence of maximum magnitude earthquakes, Bulletin of the Seismological Society of America, Vol: 64, No:2, 393-414, 1974.

q-Difference Operator on $L_q^2(0,+\infty)$

Meltem Sertbaş[1] and Coşkun Saral[2]

[1]Karadeniz Technical University, Faculty of Sciences, Department of Mathematics, Trabzon, Turkey

[2]Karadeniz Technical University, Institute of Natural Sciences, Trabzon, Turkey

m.erolsertbas@gmail.com , csaral5361@gmail.com

Abstract

In this research, the minimal and maximal operators defined by q- difference expression are given in the Hilbert space $L_q^2\ (0,+\infty)$. The existence problem of a q^{-1}-normal extension for the minimal operator is mentioned. In addition, the sets of the minimal operator spectrum and the maximal operator spectrum are examined.

Keywords: q -difference operator; Minimal operator; Maximal operator; q -deformed operatör; q -formally normal operatör; Spectrum sets.

REFERENCES

1. L. Euler, Introductio in Analysin Infinitorum. vol. 1. Lausanne, Switzerland, Bousquet, 1748 (in Latin).
2. C.F. Gauss, Disquisitiones Generales Circa Seriem infinitam, Werke, 124-162, 1813.
3. S. Ota, Some classes of q-deformed operators, J. Operator Theory 48, no. 1, 151–186, 2002.
4. S. Ota, F.H. Szafraniec, Notes on q-deformed operators, Studia Math., 165, no. 3, 295-301, 2004.
5. S. Ota, F.H. Szafraniec, q-Positive definiteness and related topics, J. Math. Anal. Appl., 329, no. 2, 987-997, 2007.
6. J. Cimpric, Yu. Savchuk , Schmüdgen , K. On q-normal operators and quantum complex plane, Trans. Amer. Math. Soc., 366, no.1, 135 -158, 2014.
7. A. Lavagno, A.M. Scarfone, q-deformed structure and generalized thermodynamics, Rep. Math. Phys., 55, v.3, 423-433, 2005.

NEW ANALYTICAL APPROACH TO THE GENERALIZED FISHER EQUATION

Gülnur Yel[1*], Miraç Kayhan[2]

[1]Faculty of Education, Final International University, Kyrenia, Mersin 10, Turkey

[2] Department of Mathematics, University of İnonu, Malatya, Turkey

gulnur.yel@final.edu.tr, mirackayhan@yandex.com,

Abstract

In this paper, we use an effective method which is the rational sine-Gordon expansion method to present new wave simulations of a governing model. We consider the conformable (1+1)-dimensional Fisher equation which is used to describe interactive relation between diffusion and reaction. Various type solutions such as dark, bright and singular soliton solutions have been obtained. Finally, the physical behaviours of the obtained solutions are shown by 3D, 2D and contour surfaces.

Keywords: Rational sine-Gordon expansion method; the conformable (1+1)-dimensional Fisher equation; travelling wave solutions.

REFERENCES

1. Aksan EN, Bulut H. , Kayhan M., Some Wave simulation properties of the (2+ 1) dimensional breaking soliton equation, ITM web of conferances, 13 (1014), 2017.
2. Bulut H. , Aksan EN, Kayhan M. , Sulaiman T., New solitary wave structures to the (3+ 1) dimensional Kadomtsev–Petviashvili and Schrödinger equation, Journal of Ocean Engineering and Science, 373-378, 2019.
3. Qin Zhou, M. Ekici, A. Sonmezoglu, J. Manafian, S. Khaleghizadeh, M. Mirzazadeh, Exact solitary wave solutions to the generalized Fisher equation, Optik, 127 (24), 12085-12092, 2016.
4. A.M. Wazwaz, The extended tanh method for abundant solitary wave solutions of nonlinear wave equations, Appl. Math. Comput. 187 ,1131–1142, 2007.

On Modified Weingarten Parallel Ruled Surface

Gülden Altay Suroğlu [1], **Yusuf Güzel** [2]

[1,2] Department of Mathematics, University of Firat, Elazig, Turkey

galtay@firat.edu.tr, ysfgzl023@gmail.com

Abstract

In this paper, we obtain some new characterization for Weingarten ruled surface according to Modified frame.

Keywords: Modified frame; ruled surface; Weingarten surface.

REFERENCES

1. Aslı Aybüke Akdemir, Eğrilik ile modifiye edilmiş ortogonal çatıda eğriler, yüksek lisans tezi, Kayseri, 2021.
2. Dae Won Yoon, Some Properties of Parallel Surfaces in Euclidean 3- Spaces, Honam Mat. J., Vol:30, No:4, 637-641, 2018.
3. Kemal Eren, Hidayet Huda Kosal, Evolution of space curves and the special ruled surfaces with modified orthogonal frame, AIMS Mathematics, Vol:5, No:3, 2027–2039, 2020.
4. Muhammed T. Sariaydin, Talat Körpinar and Vedat Asil, On Parallel p-equidistant Ruled Surfaces by Using Mofied Orthogonal Frame with Curvature in E 3, Bol. Soc. Paran. Mat, 40, 1-7, 2022.
5. Selçuk Baş, Talat Körpınar, Modified Roller Coaster Surface in Space, Mathematics, Vol:7, No:5, 2019.
6. Umut Ziya Savcı, Ali Görgülü, Cumali Ekici, Parallel surfaces of ruled Weingarten surfaces, Vol:3, No:4, 237-246, 2015.

*Abstract Submission should be prepared only **1 page**.

OPINIONS OF TEACHER CANDIDATES ON WEB 2.0 TOOLS AND THEIR USAGE IN EDUCATION

Ertuğrul AKGÜL*, Assist. Prof. Dr. ELİF ÇİL
Department of MathematicsandScienceEducation, Master of Scienceeducation, Ordu University, Ordu.
O19521200023@odu.edu.tr

Theconcept of Web 2.0 wasfirstusedbyDarcyDiNucci in 1990. Internet technologies, called Web 2.0 tools, provideeffectivecommunicationandfastandeasyinformationsharing in education, as in manyareas of ourlives. Withthefeatures of thesetools, such as activedatadesign, informationstorage, feedback, andvisual/verbal data presentation, teachersfindtheopportunitytodesign/designactivitiesthatappealtoparticipantsfromallagegroups. Web 2.0 tools, frequentlyused in education, aregroupedaccordingtotheirusageareas; they can be groupedunder nine headings. These; includemindmaps, digitalboards, creatingpostersandcartoons, writingstoriesandbooks, takingnotesandcreatingblogs, creatingtestsandpuzzles, preparingeffectivepresentations, andcreatinginformationpostersandvirtualclassroomapplications. Inthisstudy, whichwasdesignedbased on thisinformation, pre-service scienceteachers' views, whowerestudying at the 2nd, third, and 4th-grade levels at a stateuniversity in North Anatolia about web 2.0 wereexamined. Thesample of thestudyconsists of thirty-threeteachercandidates. A holisticsingledesignstudy, which is one of thequalitativeresearchmethods, wasusedtodeterminepre-service teachers' viewsabout web 2.0 toolsandtheirusage in education. Thequestionnaireconsisting of nine questions, developedby Timur et al. in 2020 andconsisting of open-endedquestions, wasdistributedtothepre-service teacherswhovolunteeredtoparticipate in thestudyvia Google documents. Thecollecteddatawastransferredtothecomputer. Thenames of theteacher-teachercandidateswere not used in thestudy, andtheywerenamed Ö1, Ö2, etc. Thetransferredanswerswerecategorizedbythethematiccodingmethod, andtheappropriateanswersweresematchedwiththeappropriatethemes. Thefrequencyvalues of thethemeswerecalculatedwiththe MAXQDA program.

Keywords: digital literacy, teacher candidate, internet tools, qualitative analysis.

References

Uysal, M. Z. (2020). *İlkokul 4. sınıf fen bilimleri dersinde web 2.0 animasyon araçları kullanımının çeşitli değişkenlere etkisi* (Master'sthesis, Niğde Ömer Halisdemir Üniversitesi/Eğitim Bilimleri Enstitüsü).

TİMUR, S., TİMUR, B., ARCAGÖK, S., & ÖZTÜRK, G. (2020). Fen Bilimleri Öğretmenlerinin Web 2.0 Araçlarına Yönelik Görüşleri. *Journal of KirsehirEducationFaculty*, *21*(1).

Köse, Ö. Ö., Bayram, H., & Benzer, E. (2021). WEB 2.0 Destekli Argümantasyon Uygulamalarının Ortaokul Öğrencilerinin Kuvvet ve Enerji Konusundaki Başarılarına, Tartışmacı Tutumlarına ve Teknoloji Tutumlarına Etkisi. *Erciyes Journal of Education*, *5*(2), 179-207.

TİMUR, S., YILMAZ, Ş., & KÜÇÜK, D. Web 2.0 Uygulamalarının Fen Bilgisi Öğretmen Adaylarının Öz-Yeterlik İnançları Üzerindeki Etkisinin İncelenmesi. *İstanbul Aydın Üniversitesi Eğitim Fakültesi Dergisi*, *7*(2), 291-311.

GÜRLEROĞLU, L., & YILDIRIM, M. ORTAOKUL ÖĞRENCİLERİNİN WEB 2.0 DESTEKLİ EĞİTSEL WEB SİTESİ İLE İLGİLİ GÖRÜŞLERİNİN İNCELENMESİ. *Milli Eğitim Dergisi*, *51*(233), 191-217.

SEMI-INVARIANT SUBMANIFOLDS OF GENERALIZED KENMOTSU MANIFOLD ADMITTING A QUARTER SYMMETRIC NON METRIC CONNECTION

Ramazan Sarı [1] **and İnan Ünal** [2]

[1] Gümüşhacıköy Hasan Duman Vocational School, Amasya University, Amasya, Turkey

ramazan.sari@amasya.edu.tr,

[2] Department of Computer Engineering, University of Munzur, Tunceli, Turkey

inanunal@munzur.edu.tr

Abstract

In this paper, semi-invariant submanifolds of a generalized Kenmotsu manifold endowed with a quarter symmetric non-metric connection are studied. Necessary and sufficient conditions are given on a submanifold of a generalized Kenmotsu manifold to be semi-invariant submanifold with quarter symmetric non-metric connection. Morever, we studied integrabilities and parallel conditions of the distribution on semi-invariant submanifolds of generalized Kenmotsu manifold with quarter symmetric non-metric connection.

Keywords: Generalized Kenmotsu manifold; semi-invariant submanifold; quarter symmetric non metric connection.

REFERENCES

1. S. K. Chaubey and R.H. Ojha, On semi-symmetric non-metric and quarter symmetric metric connections, Tensor N. S., 70(2), 202-213, 2008.
2. A. K. Dubey and R.H. Ojha and S.K. Chaubey, Some properties of quarter symmetric non-metric connection in a Kahler manifold, Int. J. Contemp. Math.Sciences, 5(20), 1001-1007, 2010.
3. S. . Golab, On semi- symmetric metric and quarter symmetric linear connections, Tensor 29, 249-254, 1975.
4. R. Sarı, M.A. Akyol and E. Aksoy, Some curvature properties of an S-manifold with a quarter symmetric non metric connection, International Journal of Applied Mathematics and Statistic 56(3), 93-102, 2017
5. A. Turgut Vanli, and R. Sari, Generalized Kenmotsu Manifolds. Communications in Mathematics and Applications, 7(4), 311- 328, 2016.

SEMI-INVARIANT RIEMANNIAN SUBMERSIONS WITH RICCI SOLITON

Ramazan Sarı [1] and İnan Ünal [2]

[1] Gümüşhacıköy Hasan Duman Vocational School, Amasya University, Amasya, Turkey

ramazan.sari@amasya.edu.tr,

[2] Department of Computer Engineering, University of Munzur, Tunceli, Turkey

inanunal@munzur.edu.tr

Abstract

In this paper, we introduce ricci solitons on semi invariant Riemannian submersions from contact manifold. We investigate any foliation of such a submersion is a Ricci soliton. Moreover, we show Einstein conditions.

Keywords: Riemannian submersion; semi-invariant Riemannian submersion; Ricci soliton.

REFERENCES

1. M. A. Akyol, R. Sarı and E. Aksoy, Semi-invariant ξ-Riemannian submersions from almost contact metric manifolds, Int. J.Geom. Methods Mod. Phys., 14(5), 1750074, 2017.
2. Y. Gündüzalp, Almost hermitian submersions whose total manifolds admit a ricci soliton, Honam Math. J., 42(4), 733-745,2020.
3. R. S. Hamilton, The Ricci flow on surfaces, Contemp. Math., 71,237-261, 1988.
4. Ş.E. Meriç and E. Kılıç, Riemannian submersions whose total manifolds admit a ricci soliton, Int.J. Geo. Met. and Mod. Phy., 16(12), 1950196, 2019.
5. Ş.E.Meriç, M. Gülbahar and E. Kılıç, Some inequalities for Riemannian submersions, An. Ştiint. Univ. Al. I. Cuza Ia¸si. Mat. (N.S.), 63(3), 471-482, 2017.

ON A HOMOGENEOUS-HETEROGENEOUS REACTIONS WITH NEWTONIAN HEATING IN FLOW OF A THIRD GRADE FLUID

Canan Unlu
Department of Mathematics, University of Istanbul, Turkey

Abstract: The fundamental goal here is to display and analyze the feature of homogeneous-heterogeneous responses in the MHD flow of third grade liquid over an extending surface. Both magnetic and electric fields are considered. Advanced heat transfer technique (i.e., Newtonian heating) and heat creation/absorption impacts are utilized in the formulation. Homogeneous and heterogeneous responses are considered inside the liquid and at the boundary individually. Approximate convergent solutions are constructed. Impacts of different numbers on the velocity, temperature and concentration fields are examined and scrutinized. Numerical estimations of friction and local Nusselt number are determined. Concentration fields for homogeneous and heterogeneous reaction numbers are found inverse.

Keywords: Newtonian heating; electric and magnetic fields; homogeneous-heterogeneous reactions; third-grade fluid.

References

[1] Abbasbandy, S., Yurusoy, M. and Gulluce, H., "Analytical solutions of non-linear equations of power-law fluids of second grade over an infinite porous plate," Math. Comput. Appl., 19 (2), pp. 124 (2014).
[2] Hayat, T., Shafiq, A. and Alsaedi, A., "MHD axisymmetric flow of third grade fluid by a stretching cylinder," Alexandria Engineering Journal, 54 (2), pp. 205-212 (2015).
[3] Sheikholeslami, M., Ellahi, R., Ashorynejad, H. R., Domairry, G. and Hayat, T., "Effects of heat transfer in flow of nanofluids over a permeable stretching wall in a porous medium," J. Comput. Theore. Nanoscience 11 (2) (2014) 486-496.
[4] Hayat,T., Shafiq, A. and Alsaedi, A., "Effect of Joule heating and thermal radiation in flow of third-grade fluid over radiative surface," Plos One, 9 (1), pp. e83153 (2014).

THE EFFECT OF POLYVINYL ALCOHOL FIBER ON THE MECHANICAL PROPERTIES OF HIGH STRENGTH GEOPOLYMER CONCRETES

Ali Öz[1], Oğuzhan Çelebi[2], Gökhan Kaplan[2], Barış Bayrak[3], Haluk Görkem Alcan[3], Abdulkadir Cüneyt Aydın[2]

[1] Narman Vocational School, University of Ataturk, Erzurum, Turkey

[2]Department of Civil Engineering, University of Ataturk, Erzurum, Turkey

[3]Department of Civil Engineering, University of Kafkas, Kars, Turkey

alioz@atauni.edu.tr

Abstract

Today, it is quite common to use conventional Portland cement-based concrete in buildings. Geopolymer concrete has been proposed as an alternative to Portland cement concrete due to the high greenhouse gas effect and the environmental pollution of energy consumption caused by cement consumption. In this study, metakaolin and ground blast furnace slag (GBFS) were preferred as binder materials to increase the early age strength of concrete and not cause loss of strength due to possible porosity in later ages. In addition, due to the high probability of brittle fracture of geopolymer concrete after curing at high temperatures, it aims to cope with these negative properties by using polyvinyl alcohol fibers (PVA) without reducing the flexural, compressive strengths of concrete. In this study, to examine the effect of PVA on the mechanical properties of high strength geopolymer concrete, three mixtures were designed, 0.3% kg/m^3, 0.6 kg/m^3 fiber and did not contain any fibers which were kept in a curing environment at 80 ºC for 10 hours. GBFS and quartz powder vary, with the same weight percentage of metakaolin as the binder in the mixtures. In addition, the aggregate (quartz) and alkali activator contents of the mixtures are the same. The test results showed that the mixture containing 0.6 kg/m^3 PVA reached 136.2 MPa by increasing the compressive strength by 18.77% compared to the without PVA mixture, and reaching 6.03 MPa by increasing the flexural strength by 63.41%. In addition, it has been determined that the use of PVA as a reinforcing element prevents brittle fracture in geopolymer concrete samples. The results showed that PVA could be used as a reinforcement element to prevent the negative behavior of geopolymer concrete, such as high mechanical properties and less energy consumption and prevent its negative behavior, such as a brittle fracture.

Keywords: Geopolymer Concrete, Polyvinylin Alcohol Fiber, High Strength Concrete

MECHANICAL PROPERTIES OF METAKAOLIN BASED GEOPOLYMERS USING WASTE GEOPOLYMER POWDER AND LIME

Gökhan Kaplan[1], Oğuzhan Çelebi[1], Ali Öz[2], Haluk Görkem Alcan[3], Barış Bayrak[3], Abdulkadir Cüneyt Aydın[1]

[1] Department of Civil Engineering, University of Atatürk, Erzurum, Turkey

[2]Narman Vocational School, University of Atatürk, Erzurum, Turkey

[3]Department of Civil Engineering, University of Atatürk, Erzurum, Turkey

gkaplan@atauni.edu.tr,

Abstract

In this study, geopolymer samples were prepared by using binders such as waste geopolymer powder, lime and metakaolin. Sodium silicate and sodium hydroxide were used for alkali activation. Quartz aggregate was used in the preparation of the samples. Geopolymer samples with waste geopolymer powder, with a weight ratio of 36% in $1m^3$ volume, were prepared at 60 ℃ for 6-8-10 hours of curing. The test results showed that the highest compressive strength of the mixtures was 79 MPa with the increase of the lime ratio in the 8 hour curing period. It was determined that the highest compressive strength was 73 MPa in the samples in which metakaolin was never used and the lime ratio was maximum, the compressive strengths of the samples with 0.05% lime ratio of metakaolin and 0.025% lime ratio of metakaolin and 0.05% lime ratio of metakaolin were 74 and 79 MPa, respectively. In addition, capillary water absorption tests were also carried out on the samples. Capillarity coefficients of geopolymer mixtures are below 0.377 $kg/m^2.min^{0.5.}$ It has been suggested that geopolymer mixtures produced from recycled geopolymer powder can be used in the load-bearing elements of structures that need to remain at the level of use immediately under the effect of dynamic load.

Keywords: Geopolymer powder; Recycle concrete; Metakaolin; Lime

ANALYSIS OF PROBLEMS POSED BY PRESERVICE ELEMENTARY MATHEMATICS TEACHERS FOR THE ADDITION AND SUBTRACTION OF FRACTIONS

Author: Şafak Nur DUMAN **Advisor: Doç. Dr. Meral Cansız AKTAŞ**

Mathematics and Science Education, Ordu University, Ordu, Turkey
safaknurduman@odu.edu.tr,

Abstract

In this study, it was aimed to analyze the problems posed by preservice elemantary mathematics teachers candidates for addition and subtraction with fractions. The case study, one of the qualitative research methods, was adopted in the research. Fifty-nine primary school mathematics teacher candidates participated in the research.
The Problem Posing Test (PPT) consisting of ten items for addition and subtraction with fractions was used as a data collection tool in the research. The data were analyzed by using the error types in the studies of (Işık & Kar, 2014) and (Işık & Kar, 2012). As a result of the analysis, the most common errors are listed as inability to establish a part-whole relationship, assigning a natural number meaning to fraction numbers, not assigning appropriate units to fraction numbers, not assigning meaning to whole parts of integer fractions, and expressing fraction numbers over different wholes.
As a result of the research, it was determined that the deficiencies at the conceptual level regarding the concept of fraction and the difficulties experienced in the verbal language dimension were at the center of the error types detected.

Keywords: problem posing, fractions, addition of fractions, subtraction of fractions

REFERENCES

1. KAR, T., & IŞIK, C. (2014). Ortaokul Yedinci Sınıf Öğrencilerinin Kesirlerle Çıkarma İşlemine Kurdukları Problemlerin Analizi. *Ilkogretim Online*, *13*(4).

1. Işık, C., & Kar, T. (2012). Analyzing problems posed by 7th grade students for addition operation with fractions. Elementary Education Online, 11(4).

INTEGRAL REPRESENTATIONS OF k-SRIVASTAVA HYPERGEOMETRIC FUNCTIONS

Sena Halıcı, Ayşegül Çetinkaya [1]

[1] Department of Mathematics, University of Kırşehir Ahi Evran, Turkey

halıcı.sena@ogr.ahievran.edu.tr, acetinkaya@ahievran.edu.tr

Abstract

In this study, we introduce k-Srivastava hypergeometric functions by using Pochhammer k-symbol. We also present some integral representations for these functions.

Keywords: Srivastava hypergeometric functions; Pochhammer k-symbol; Integral representations.

REFERENCES

1. Choi J., Hasanov A., Srivastava H.M., Turaev M., Integral representations for Srivastava's triple hypergeometric functions, Taiwanese J. Math., 15, 2751-2762, 2011.
2. Diaz R., Pariguan E., On hypergeometric functions and Pochhammer k-symbol, Divulg. Mat., 15, 179–192, 2017.
3. Hasanov A., Srivastava H.M., Turaev M., Decomposition formulas for some triple hypergeometric functions, J. Math. Anal. Appl., 324,955–969, 2006.
4. Kıymaz İ.O., Çetinkaya A., Agarwal P.A., Study on the k-generalizations of some known and fractional operators, Journal of Inequalities and Special Functions, 8 (4), 31-41, 2017.
5. Mubeen S., Habibullah G., An integral representation of some k-hypergeometric functions. Int. Math. Forum 7, 203–207, 2012.
6. Mubeen S., Iqbal S., Rahman G., Contiguous function relations and an integral representation for Appell k-series $F_{1,k}$, Inter. J. Math. Res., 4 (2), 53-63, 2015.
7. Mubeen S., Rahman A., A note on k-gamma function and Pochhammer k-symbol, J. Informatics Math. Sci. 6, 93–107, 2014.
8. Srivastava, H.M, Manocha, H, Treatise on Generating Functions, John Wiley and Sons, New York, USA, 1984.

RECURSION FORMULAS OF k-SRIVASTAVA HYPERGEOMETRIC FUNCTIONS

Sena Halıcı, Ayşegül Çetinkaya [1]

[1] Department of Mathematics, University of Kırşehir Ahi Evran, Turkey

halıcı.sena@ogr.ahievran.edu.tr, acetinkaya@ahievran.edu.tr

Abstract

In this study, various recursion formulas are obtained for k-Srivastava hypergeometric functions defined by k-Pochhammer symbol.

Keywords: Srivastava hypergeometric functions; Pochhammer k-symbol; Recursion formulas.

REFERENCES

1. Diaz R., Pariguan E., On hypergeometric functions and Pochhammer k-symbol, Divulg. Mat., 15, 179–192, 2017.
2. Kıymaz İ.O., Çetinkaya A., Agarwal P.A., Study on the k-generalizations of some known and fractional operators, Journal of Inequalities and Special Functions, 8 (4), 31-41, 2017.
3. Mubeen S., Habibullah G., An integral representation of some k-hypergeometric functions. Int. Math. Forum 7, 203–207, 2012.
4. Mubeen S., Iqbal S., Rahman G., Contiguous function relations and an integral representation for Appell k-series $F_{1,k}$, Inter. J. Math. Res., 4 (2), 53-63, 2015.
5. Mubeen S., Rahman A., A note on k-gamma function and Pochhammer k-symbol, J. Informatics Math. Sci. 6, 93–107, 2014.
6. Srivastava, H.M, Manocha, H, Treatise on Generating Functions, John Wiley and Sons, New York, USA, 1984.
7. Şahin R., Recursion formulas for Srivastava hypergeometric functions. Mathematica Slovaca 65 (6), 1345-1360, 2015.

COMPUTATION OF H_∞-NORM OF A TRANSFER MATRIX VIA BISECTION ALGORİTHM

Hasan GÜNDÜZ[1] **Ercan ÇELİK**[2]

[1] Erzincan Binali Yıldırım University, Faculty of Art and Science, Department of Mathematics, Bingöl-Turkey

[2]Kyrgyz-Turkish Manas University, Department of Applied Mathematics and Informatics, Bishkek-Kyrgyzstan

hgunduz@bingol.edu.tr, ercan.celik@manas.edu.kg

Abstract

In this paper, we compute H_∞-norm of a transfer matrix, via Bisection Algorithm. The algorithm is given and applied some problems. The problems are choosen from diverse areas of control theory such as aircraft models and decentralized interconnected systems,

Keywords: H-infinity control; Bisection method; Hankel singular values, Hamiltonian.

REFERENCES

1. Boyd S., Balakrishnan V., Kabamba P., A Bisection Method for Computing the H_∞-norm of a Transfer Matrix and Related Problems, Math. Control Signals Systems, 2, 207-219, 1989."
2. Leibfritz F., Lipinski W., Description of the benchmark examples in COMPl_eib 1.0, Tech-Report, 2003.
3. Kuster G. E., H-infinity Norm Calculation via a State Space Formulation, Master of Science in Mathematics, Virginia, 2012.
4. Suman S. K., Kumar A., Reduction of Large-Scale Dynamical Systems by Extended Balanced Singular Perturbation Approximation, Internation Journal Mathematical , Engineering and Management Sciences, Vol. 5 No. 5, 939-956, 2020.

SEGMENTATION OF ELECTRICITY SUBSCRIBERS BASED ON THE CONSUMPTION PATTERNS

Hicran Gümüşbaş[1], Ulas Vural[1, 2]

[1] Nar System Technology Inc., Istanbul, Turkey

[2] Kocaeli Health and Technology University, Kocaeli, Turkey

hicran.gumusbas@nar.com.tr, ulas.vural@kocaelisaglik.edu.tr

Abstract

Electricity is a strategic and expensive resource that must be carefully managed. Discovering and accurately modeling the temporal consumption patterns of the subscribers is crucial for offering a reliable energy supply. In addition, energy consumption models are also useful for determining anomalies in the consumption trends and forecasting high-load risks on power distribution nodes.

Feasible modeling of consumption data should be both efficient and accurate. This paper presents a method to cluster customers with similar consumption patterns to increase the overall model accuracy while keeping the number of generated models low. Hourly energy consumption data is collected from real commercial electricity subscribers by using automated meter reading systems. These time-series data are clustered by using the K-means algorithm.

The clustering algorithm is tested on a dataset that contains a 104-day record of hourly energy consumption of 376 commercial subscribers in Turkey. The experimental results are given for two different normalization schemes of daily and weekly usage. The optimal number of clusters for both experiments is determined as 6 by using the elbow heuristic. The inertia metric is calculated as 365.9 for weekly normalized and as 340.2 for the daily normalized data.

Keywords: Customer Segmentation; Electricity Consumption; Clustering; Time-series.

REFERENCES

1. Pilo F, Pisano G, Ruggeri S, Troncia M. Data analytics for profiling low-voltage customers with smart meter readings. Applied Sciences. 2021.
2. Rajabi A, Eskandari M, Ghadi MJ, Li L, Zhang J, Siano P. A comparative study of clustering techniques for electrical load pattern segmentation. Ren. and Sust. Energy Reviews. 2020.
3. Morales F, García-Torres M, Velázquez G, Daumas-Ladouce F, Gardel-Sotomayor PE, Gómez-Vela F, Divina F, Vázquez Noguera JL, Sauer Ayala C, Pinto-Roa DP, Mello-Román JC. Analysis of Electric Energy Consumption Profiles Using a Machine Learning Approach: A Paraguayan Case Study. Electronics. 2022.

Dstabil: A Matlab Package for Stability Analysis and Analytical/Numerical Solution of Linear Delay Differential Equation Systems

Noorolhuda Wyal[1], Erhan Coşkun[2]

Department of Mathematics, Karadeniz Technical University, Turkey

noorwyal@gmail.com

erhan@ktu.edu.tr

Abstract

In this study we consider a linear system of delay differential equations and develop an application called "Dstabil" to carry out stability analysis of equilibrium points of the following system.

$$X' = AX(t - D) + BX, t > 0$$
$$X(t) = \phi(t), t \in [-D, 0]$$

A is a square matrix with dimension $n \times n$ $(n = 1,2,3)$, D is a diagonal delay matrix of the same size. Dstabil also has components to determine analytical solution interactively using method of steps and method of characteristics.

Keywords: Delay differential equations; stability of equilibrium solutions.

REFERENCES

1. Hal, Smith, An Introduction to Delay Differential Equations with Applications to the Life Sciences Springer New York, 2011.
2. Coşkun, E., Course Notes for Numerical Analysis, www.erhancoskun.com.tr, 2022.

Renewable energy in smart grid: Photovoltaic Power Monitoring System based on Machine Learning using an Open-Source IoT Platform

Hakam Youness[1], Gaga Ahmed[1] and Benachir El Hadadi[1]

[1]Research Laboratory of Physics and Engineers Sciences (LRPSI). Department of Physics, Team Embedded Systems Engineering, Automation, Signal, Telecommunications and Intelligent Materials Polydisciplinary Faculty (FPBM), Sultan Moulay Slimane University (USMS), Beni-Mellal, Morocco
hakamyouness2@gmail.com
gaga.ahmad@gmail.com
doyen.fp@gmail.com

Abstract

This paper proposes a PV monitoring system based on the internet of things, it is the best solution to monitoring our system. in this system, we have utilized the Raspberry Pi card as a server communicates with a NodeMcu (ESP32) as a client using the MQTT and HTTP protocols respectively. Our solution is composed of two steps, the first step is consisting of power measurement. A voltage constant and current was measured by the ACS712 sensor, we measured the power and energy of the solar panels every 5 min. a DHT sensor has utilized to measure the temperature of solar panels. these measures will be shown on the Node-red platform and stored as a database in the SQLite programming language, SQLite is introduced to reduce the database complexity. Raspberry pi card recorded this database in real-time using WiFi or cable ethernet, we can make with this database accurate projections about the efficiency of solar panels and command our system. However, in the second step, we based on this database, we involve the command in our system by the best method(algorithm) of prediction for our case. the power delivered by solar panels was predicted with the use of machine learning (a model decision tree), which enabled us to generate the forecasts. In practice, we use an electronic card that can support this type of machine learning algorithm, for this, we used the raspberry pi card. Node-red is the most suitable interface to apply this algorithm also it allows us to monitor all measured by the dashboard in real-time with a laptop (WiFi) and with a smartphone (4G).as the result of this work, we have made a smart system based on machine learning that allows to integrates PV into the smart grid. this approach allows us to manage our system in a more efficient, automated, and intelligent manner.

Keywords: Power Monitoring; photovoltaic (PV); smart grid; Internet of Things (IoT); WIFI; NodeMcu; Raspberry pi; Node-RED; SQLite; Machine Learning.

Overview of Some Interval Valued Sequence Spaces

Gülsen Kılınç[1] **Mehmet Sezai Yıldırım** [2]

[1] Department of Mathematics and Science Education, Faculty of Education,

Adıyaman University, Adıyaman, TURKEY

Department of Mathematics, Faculty of Science and Arts,

Adıyaman University, Adıyaman, TURKEY

gkilinc@adiyaman.edu.tr,

sezai63songul@gmail.com,

Abstract

In this paper, Interval sequence spaces, which are a different example of quasilinear space, will be introduced and information will be given about the algebraic and topological structure of one of these spaces.

Keywords: Interval valued sequences; Quasilinear space; Haussdorff metric .

REFERENCES

1. Aseev, S. M., Quasilinear operators and their application in the theory of multivalued mappings, Proceedings of the Steklov Instıtute of Mathematics,, Vol:2, 23–52, 1986.

2. Yılmaz, Y., Çakan, S., Aytekin, Ş. Topological Quasilinear spaces, Abstract And Applied Analysis, Article ID 951374, Doi: 1155/2012/951374, 10 pages, 2012.

3. Moore, E., Ramon., Kearfott, R. Baker; Cloud, Michael J., Introduction to interval analysis, Society for ındustrial and Applied mathematics, Philadelphia,, 2009.

4. Maddox, I.J., Elements of functional analysis, Cambridge University Press (Second Edition), Cambridge, 1988.

5. Çakan, S. Some new results related to theory of normed quasilinear spaces, PhD Thesis, Inönü University, Turkey, 2016.

BLOW UP OF SOLUTIONS FOR THE WAVE EQUATION VARIABLE COEFFICIENTS

Erhan Pişkin[1] **and Ayşe Fidan**[1]

[1]Department of Mathematics, University of Dicle, Diyarbakır, Turkey

episkin@dicle.edu.tr and afidanmat@gmail.com

Abstract

In this presentation, we obtain the wave equation with variable coefficients. Under suitable conditions on variable coefficients, we prove the blow up of solutions.

Keywords: Blow up; Wave equation; Variable coefficients.

REFERENCES

1. E. Pişkin, Existence, decay and blow up of solutions for the extensible beam equation with nonlinear damping and source terms, Open Mathematics, 13, 408-420, 2015.
2. X. Zheng, Y. Shang, X. Peng, Blow up of Solutions for a Nonlinear Petrovsky Type Equation with Time-dependent Coefficients, Acta Mathematicae Applicatae Sinica, English Series, 36(4) 836-846, 2020.

ADJOINT CURVES FROM THE MODIFIED FRAME OF INVOLUTE CURVE

Süleyman ŞENYURT,Eda ÖZTÜRK

Department of Mathematics Ordu University, ORDU/TURKEY

senyurtsuleyman52@gmail.com , edaozturk6652@gmail.com

Abstract

In this study, firstly, a modified frame was created from the involute curve of a curve. The adjoint curve of the involute curve was obtained by using the modified frame. Finally, the connections of the obtained adjoint curves with the main curve were calculated.

Keywords: involüt curve, adjoint curve, motifiyle frame

REFERENCES

1) Bukcu B, Karacan MK. On the modified orthogonal frame with curvature and torsion in 3-space. Mathematical Sciences and Applications E-Notes 2016; 4
2) Lone, M.S., ES, H., Karacan, M.K.,2019, “On Some Curves with Modified Orthogonal Frame in Euclidean 3-Space”, Iran J Sci Technol Trans Sci 43, 1905–1916
3) Hacısalihoğlu, H. H., 2000, "Diferensiyel Geometri I. Cilt", Ankara Üniversitesi Yayınları, Ankara, 1-167.
4) Sasai,T., “The Fundamental Theorem of Analytic Space Curves And Apparent Singularities of Fuchsian Differential” Equations,Tohoku Math. Journ., 36 (1984),17-24
5) Nurkan, S.K., Güven, İ.A., Karacan, M.K.,2019, “Characterizations of Adjoint Curves in Euclidean 3-Space”, Proc. Natl. Acad. Sci., India, Sect. A Phys. Sci. 89, 155–161
6) Lone MS, Es H, Karacan MK, Bukcu B.,2019, “Mannheim curves with modified orthogonal frame in Euclidean 3-space”, Turkish Journal of Mathematics. 2019; 43(2): 648- 663.

Several Properties of Compact operators on some generalized Fibonacci difference sequence spaces

Murat Candan

Department of Mathematics, University of İnönü, Malatya, Turkey

murat.candan@inonu.edu.tr

Abstract

In this work, we are going to characterize the matrix classes $\left(l_1, l_p(\hat{F}_r^s)\right)$ $(1 \le p < \infty)$, in which $l_p\left(\hat{F}_r^s\right)$ is some Fibonacci difference sequence spaces for $r, s \in IR$. We also obtain estimates for the norns of bounded linear operators L_A defined by these matrix trasformations and calculate conditions to derive the corresponding subclasses of compact matrix operators by using the Hausdorff measure of noncompactness.

Keywords: Sequence spaces, Fibonacci numbers, compact operators, Hausdorff measure of noncompactness.

REFERENCES

1. Alotaibi A, Mursaleen M, AS Alamsi B and Mohiuddine S.A; Compact operators on some Fibonacci difference sequence spaces; JIA; 2015:203, 8 pages.
2. Candan M; A new sequence space isomorphic to the space $l(p)$ and compact operators; J.Math.Comput.Sci.4(2014), No.2, 306-334.

Operators on Some Vector-Valued Orlicz Difference Sequence Spaces

Murat Candan

Department of Mathematics, University of İnönü, Malatya, Turkey

murat.candan@inonu.edu.tr

In this work, we are going to give some sequences of operators which have the same function with a basis for some vector-valued Orlicz difference sequence spaces. At the same time, we characterize the space $B(h_M(\Delta X), Y)$ of continuous operators from $h_M(\Delta X)$ is the space of all Δx – valued sequences $x = (x_k)$ such that

$$\sum_{k=1}^{\infty} M\left(\frac{\|\Delta x_k\|}{\rho}\right) < \infty \text{ for all } \rho > 0.$$

Exactly, we derive that each $T \in B(h_M(\Delta X), Y)$ is equivalent, under certain conditions, to arbitrary sequence $A = (A_k)_{k=1}^{\infty}$ of operators $A_k \in B(\Delta X, Y)$.

Keywords: Operator spaces, Orlich difference sequence spaces, representations of functionals.

REFERENCES

1. Yılmaz Y, Özdemir M.K, Solak İ, Candan M; Operators on some vector-valued Orlicz sequence spaces; F.Ü.Fen ve Mühendislik bilimleri dergisi, 17(1), 59-71, 2005.

Some results on convergence of soft sequences in soft topological spaces

Murat CANDAN

Department of Mathematics, University of İnönü, Malatya, Turkey

murat.candan@inonu.edu.tr

Abstract

The notion of soft set was given first by Molodtsov in 1999 [1]. Soft set theory, which brought a new approach to the solution of uncertainties, was later studied in many areas of mathematics from different aspects. Shabir and Naz gave notion of soft topological space in 2011 [2]. In 2013, Das and Samanta obtained some results by describing the convergence of sequences in soft metric spaces [3].

In this study, we obtain some characterizations about the convergence of sequences in soft topological spaces.

Keywords: Soft set, soft point, soft topological space, soft sequence,

REFERENCES

1. Molodtsov, D.; Soft set theory-First results, Comput Mat Appl 37 (4/5): 19-31 (1999).
2. Shabir, M. and Naz, M.; On soft topological spaces, Comput Mat Appl 61 (7), 1786-1799 (2011)
3. Reddy, B.S., Yogesh, K., Jalil, S.; Some results on soft sequences, International Journal of Mathematics and its Applications, 3 (4), 37-44 (2015)

THE CHARACTERIZATIONS OF EQUIDISTANT RULED SURFACES GENERATED BY BINORMAL VECTORS

Yanlin Li[1], Süleyman Şenyurt[2,*], Ahmet Özduran[2], Davut Canlı[2]

[1]School of Mathematics, Hangzhou Normal University, Hangzhou, China

[2]Department of Mathematics, Ordu University, Ordu, Turkey

*senyurtsuleyman52@gmail.com,

Abstract

In the paper, for given two differentiable curves, the characteristics of equidistant ruled surfaces along the striction curves of binormal ruled surfaces are examined. In addition, if the ruled surface is closed, then the pitch, the angle of the pitch, and the drall of these are calculated. An example is provided at the end by providing the corresponding asymptotic plane equations to indicate that the distances between the proper points on these planes are always constant.

Keywords: Ruled surfaces; Equidistant ruled surfaces; Integral invariants.

Acknowledgement: This work was supported by Scientific Research Projects Coordination Unit of Ordu University. Project number: B-1828.

REFERENCES

1. Hacısalioğlu, H. H., Diferensiyel Geometri. II.Cilt, Ankara Üniversitesi Fen Fakültesi Yayınları, Ankara, 340, 1994.
2. Masal, M., & Kuruoğlu, N., Some Characteristic Properties of The Parallel P-Equidistant Ruled Surfaces In The Euclidean Space. Pure Applied Mathematic Sciences, India, L1: 35-42, 1999.
3. Masal, M., & Kuruoğlu, N., Some Characteristic Properties of the Shape Operators of Parallel p-Equidistant Ruled Surfaces. Bulletin of Pure and Applied Sciences, 19E (2): 361-364, 2000.
4. Masal, M., & Kuruoğlu, N., Generalized paralel pi-equidistant ruled surfaces. Acta et Commentationes Universitatis Tartuensis de Mathematica, Vol.171, 18 pages, 2013.
5. Şenyurt, S., & As, E., Some Characteristic Properties Of Parallel z-Equidistant Ruled Surfaces. Hindawi Publishing Corporation, Mathematical Problems in Engineering, Article ID 587289, 7 pages, 2013.
6. Valenontis, I., Parallel P-Äquidistante Regelflachen. Manuscripta Math, 54, 391-404, 1986.

The Sine-Gordon Expansion Method and the Modified Expansion Function Method for New Hyperbolic Wave Solutions of Wu-Zhang System Models

Tuğba YAZGAN[1], Esin İLHAN[2], Ercan ÇELİK[3] , Hasan BULUT[4]

[1] *Ataturk University, Faculty of Science, Department of Mathematics, Erzurum-TURKEY*

[2] *Kırşehir Ahi Evran University, Faculty of Engineering and Architecture*, Kırsehir- *TURKEY*

[3] *Kyrgyz-Turkish Manas University, Department of Applied Mathematics and Informatics, Bishkek-KYRGYZSTAN*

[4] *Fırat University, Faculty of Science, Department of Mathematics, Elazığ-TURKEY*

tugba.yazgan@atauni.edu.tr, eilhan@ahievran.edu.tr, ercan.celik@manas.edu.kg, hbulut@firat.edu.tr

Abstract

In this paper, we investigate some solitary wave solutions of the Wu-Zhang system by using the sine-Gordon expansion method and the modified expansion function method. We successfully obtain some new hyperbolic wave solutions of this model. After then, two and three dimensional graphs of the obtained soliton solutions of the models are drawn by choosing the appropriate parameters. All the calculations in this study are done with Wolfram Mathematica software.

Keywords: The Sine-Gordon expansion method, The modified expansion function method, Wu-Zhang system

REFERENCES

1. S. T. Mohyud-Din, S. Bibi, "Exact solutions for nonlinear fractional differential equations using exponential rational function method", Opt. Quant. Electron. 49, 64 (2017)
2. M. Kaplan, A. Bekir, A. Akbulut, E. Aksoy, "Exact solutions of nonlinear fractional differential equations by modified simple equation method", Rom. J. Phys. 60, 1374–1383 (2015)
3. N. Xiong, Y. Q. Li, J. C. Chen, Y. Chen, "One-dimensional optimal system and similarity reductions of Wu-Zhang equation", Commun. Theor. Phys., 66 (1) (2016), p. 1
4. W. X. Ma, T. Huang, and Y. Zhang, "A multiple exp-function method for non-linear differential equations and its application", Physica Scripta, 82 (2010), 065003.

A WOUND/SLOUGH MODEL ACCOUNTING FOR CELLULAR DIFFUSION AND DEBRIDEMENT

Rıdvan Yaprak[1], Erhan Coşkun[2]

[1,2] Department of Mathematics, Karadeniz Technical University, Trabzon, Turkey.

[1]ridvanyaprak@ktu.edu.tr

[2]erhan@ktu.edu.tr

Abstract

In this study, we consider the nondimensionalized wound/slough interaction model [1] and develop it to have impulsive effects which can be stated as

$$U_t = \Delta U + N_1(U,V), \boldsymbol{x} \in [0,1]^n, t > 0$$
$$V_t = D\Delta V + N_2(U,V) + I_i\big(\mathrm{V}(t_i, \boldsymbol{x})\big), i = 0,1,2, \ldots$$

where Δ is the usual Laplace operator for $n = 1,2,3$; D is a nondimensional parameter accounting for cellular diffusion; N_1 and N_2 are the reaction terms; U, V are the cellular density of wound and slough respectively.

The impulsive effects are used to simulate medical treatment technique known as debridement [2] which is implemented at $t = t_1, t_2, \ldots$ to speed up the healing process, the $I_i(.)$'s are control functions to be determined to achieve optimal healing.

Keywords: Wound healing; Impulsive differential equations; Debridement.

REFERENCES

1. Yaprak, R., Coşkun, E., "A Slough/Wound Interaction Model Accounting for Cellular Diffusion.", International Conference on Mathematics and Mathematics Education (ICMME-2021), Gazi University-Ankara, 71-72, 2021.
2. Bradley, M., Nicky, C., Sheldon, T., "The debridement of chronic wounds: a systematic review.", Health Technol Assess, 3(1999).

A STUDY ON THE STATUS OF RESEARCH ON SCİENCE TEACHERS' OPİNİONS ON STEM PRACTİCES

Gözde SEVEN[1], Assist. Prof. Dr. Elif ÇİL[2]

Department of science, University of Ordu, Ordu, Turkey

gozdeseven@odu.edu.tr , elıfcıl@odu.edu.tr

Abstract

21st century (century) skills in education are related to many fields, and these connections enable us to use different disciplines together. The areas that provide these connections in science are mathematics, engineering, and technology. Another goal of 21st-century skills in education is to raise scientifically literate individuals, and these individuals use science, mathematics, engineering, and technology (STEM) skills together. Various studies are carried out to provide individuals with these skills through training programs. In this context, teachers and students are supported with different educational training.

In the literature, it is seen that reports on STEM applications are mostly related to students. On the contrary, studies about the opinion of the teachers are limited. So that the new studies about STEM can be planned, it is essential to make a situation analysis first. The studies examining the opinion of science teachers about STEM practices in graduate studies were compiled, and suggestions were presented. The study method was carried out with document analysis, a qualitative data collection method. The date range, approach, aim, and results of the studies on science teachers' opinions about STEM applications were analyzed. The journal articles and master thesis, which are reached from the national academic database (ULAKBIM) and Higher Education Institution National Thesis Centre (YÖK), were selected for analysis between 2016 and 2022. The scope of the research was limited by the "the science course" keyword. Seventeen reports were obtained within the range of the study. Six master's theses and eleven journal articles were examined in the research. Topics are presented by relating self-efficacy and different variables. The most common problems in the studies have been determined that teachers have a shortage of materials in STEM education and teachers have difficulties integrating knowledge from different disciplines with science education. Following these, science teachers' perception of themselves as inadequate in STEM, crowded classrooms, and problems in evaluating students in STEM applications are other main problems.

According to science teachers, to eliminate these problems in STEM practices, STEM applications should be integrated into the science teaching undergraduate program with compulsory courses. And also, the content of in-service, both online and face-to-face training given to teachers, should be expanded with activities and assessment, and evaluation stages should be added to this training. It is recommended that non-thesis master's programs be opened in universities.

Keywords: STEM; STEM education; FeTeMM education; Content analysis; science course.

REFERENCES

1. Bakırcı, H., ve Kutlu, E. (2018). Fen Bilimleri Öğretmenlerinin Fetemm Yaklaşımı Hakkındaki Görüşlerinin Belirlenmesi. *Turkish Journal of Computer and Mathematics Education (TURCOMAT)*, *9*(2), 367-389.

2. Bölükbaşı, G., ve Ari, A. G. (2019). Fen bilimleri öğretmenlerinin FeTeMM eğitimi ve etkinliklerine yönelik görüşleri. *Academic Perspective Procedia*, *2*(1), 47-56.

3. Çiftçi, M., ve Çınar, S. (2017). Fen Bilgisi Öğretmenlerinin STEM Eğitiminin Fen Bilimleri Dersine Entegrasyonu Hakkındaki Görüşleri. *Sözel bildiri, VII. Uluslararası Eğitimde Araştırmalar Kongresi, Çanakkale*.

4. Deniz, A. (2019). *Fen Bilimleri Öğretmenlerinin Fetemm Ve Sosyobilimsel Konular İle İlgili Görüşlerinin İncelenmesi,* Yüksek Lisans Tezi. Van: Yüzüncü Yıl Üniversitesi Eğitim Bilimleri Enstitüsü.

5. Doğan, E., ve SARAÇOĞLU, S. (2019). Fen Bilimleri Öğretmenlerinin STEM Temelli Fen Eğitimi Hakkındaki Görüşleri. *Journal of Hasan Ali Yücel Faculty of Education/Hasan Ali Yücel Egitim Fakültesi Dergisi (HAYEF)*, *16*(2).

6. Gökçen, S. (2021). *Fen ve Matematik Eğitiminde STEM Uygulamalarına İlişkin Lisansüstü Tezlerin İçerik Analizi*, Yüksek Lisans Tezi. Bartın: Bartın Üniversitesi Lisansüstü Eğitim Enstitüsü.

7. Özmansur, N. (2019). *Fen Bilimleri Öğretmenlerinin Öğretmenlik Mesleğine Yönelik Öz-Yeterlik İnanç Düzeylerine Göre Fen Eğitiminde Kullanılan Stem Etkinlikleri Hakkındaki Görüşlerinin Değerlendirilmesi* (Master's thesis, Sosyal Bilimler Enstitüsü).

8. Yaşar, E.B. (2021). *Fen Bilimleri Öğretmenlerinin 21. Yüzyıl Becerileri Öz Yeterlilik Algıları ve STEM Tutumlarının İncelenmesi,* Yüksek Lisans Tezi. Kırıkkale : Kırıkkale üniversitesi fen bilimleri enstitüsü.

Sustainable Fisheries with Optimal Control Theory

Mahir Demir [1]

[1]Department of Mathematics, Giresun University, Giresun, Turkey

mahir.demir@giresun.edu.tr

Abstract

Fish populations are an important part of marine ecosystems, and any violation in the harvesting of fish populations is directly affecting marine ecosystems and their resilience. Since fish populations are a vital source of food for the human population worldwide, their harvesting strategies are very important for both sustainable fisheries and resilient ecosystems. There have been three common sustainable harvest techniques: seasonal fisheries, no-take marine reserves (marine protected areas), and ecosystem-based fishery management. Having these three most known techniques in mind for sustainable fisheries, in this study, I will discuss a more complete and general technique for sustainable fisheries. This technique consists of the implementation of optimal control theory together with food chain models governed by differential equations or partial differential equations. This technique not only offers optimal harvesting strategies but also offers the most profitable harvesting strategies for fish stocks and helps to investigate the effect of fisheries on their corresponding ecosystems. In many cases, no-take marine reserve areas and seasonal fisheries are natural results of the optimal control application in fishery management. Besides these, this technique provides more accurate and realistic harvesting strategies as compared to traditional harvesting strategies.

Keywords: Sustainable fishery, optimal control, harvesting, harvesting strategies, food chain modeling.

THE ELECTRONIC AND OPTICAL PROPERTIES OF GRAPHENE QUANTUM DOTS PYRENE FAMILY

Hajar Ahessab[1] **Khalid Rahmani**[1] **Mohamed Al-Hattab**[2]

[1] Department of Physics, University of Sultan Moulay Slimane, Ben-mellal, Morocco

hajarahessab2@firat.com,

Abstract

This work is devoted to the study of the electronic and optical properties of graphene-based quantum dots investigated using perturbation theorie GGA of the Perdew-Burke-Ernzerhof (PBE). The ground proprties is calculated using materials studio. Formation energy, hardness, and electrophilicity show that all structures, from pyrene to silicene QD passing through 15 CSi QD configurations, are energetically and chemically stable. It is also found that they are reactive which implies their favorite character for the possible electronic transport and conductivity. The electronic and optical properties such as absorption, Refractive index, reflectivity, and conductivity all these propreties are very sensitive to the number and position of the substituted silicon-atoms as well as the directions of the light polarization. Moreover, quantum confinement effects make the exciton binding energy of CSi quantum dots larger than their higher dimensional allotropes such as silicene, graphene, SiC sheet, and nanotube. It is also higher compared to other shapes of quantum dots like hexagonal graphene QDs and can be tailored from the ultraviolet region to the visible one. The values of the singlet-triplet splitting determined for the X- and Y-light polarized indicate that all configurations have a high fluorescence quantum yield compared to typical semiconductors, which makes them very promising for various applications such as the light-emitting diode material photovoltaic and nanomedicine.

Keywords: Graphene ; Quantum dots; GGA; PBE.

WEB 3.0 (SEMANTICS) APPLICATIONS IN EDUCATION

Süleyman Dölek[1]* – Asist. Prof. Dr. Elif Çil[2] – Assoc. Prof. Dr. Sevda Türkiş[3]
Department of Mathematics and Science Education, University of Ordu, Ordu, Turkey
suleymandolek@odu.edu.tr, elifcil@odu.edu.tr, sevdaturkis@odu.edu.tr

Abstrat

The number of web pages, which was 110 thousand in 1994, reached 63 billion in a short period of fifteen years. Today, this number is many times higher. The use of technology in educationalenvironments continues to increase day by day. As stated by many academics and researchers, the use of technology in educational areas has proven to be a mandatory requirement with the recent Covid-19 pandemic. Thanks to the new technologies, access to information, communication, and high-level calculations have become more accessible. The development ofinternet technology is the main reason for these conveniences. Thanks to the developed infrastructures and lightning-fast internet access, the creation of collaborative learning platforms in educational environments are increasing rapidly. The development of the Internet brings the web to us in different versions. Web 2.0 in the social web category came to the fore in 2004. Although artificial intelligence-based web 3.0, which is called semantic web, has not completed its development yet, it excites people in terms of the future of the web. The rapid increase of data on the Internet and the need to make it meaningful is one of the reasons for theformation of the semantic web.

Web 3.0 (Semantic Web) technology has emerged with the idea of making sense of countless data entered in the internet environment in machine language and accumulating this data on a platform. The data made sense by Web 3.0 technology makes it easier to use the internet environment and access information by returning results according to the history, usage habits, and search filters of the users. There are already technologies in metadata to use Semantic Web technology, which is encoded. Web 3.0 is the next generation of internet technology based on machine learning, artificial intelligence (AI), and blockchain technology. This concept was created by Gavin Wood, founder of Polkadot and co-founder of Ethereum. While Web 2.0 will focus on user-created content located on central websites, Web 3.0 will give users more control over their online data. Web 3.0 aims to provide personalized and relevant information more quickly using artificial intelligence and advanced machine learning techniques. Developing more intelligent algorithms and Big Data analytics means machines can intuitively understandand suggest content. This study aimed to review the studies related to the use of Web 3.0 technologies in education in a systematic way.

Keywords: Semantic Web, literature review, distance education, future education

References

Horban, Y., Humenchuk, A., Karakoz, O., Koshelieva, O., & Shtefan, I. (2021). Application of web 3.0 technologies in distance education (by levels of higher education). *Laplage em Revista*, *7*(Extra-B), 575-586.

FIRAT, E. A., & FIRAT, S. (2020). Web 3.0 in learning environments: A systematic review. *Turkish Online Journal of Distance Education*, *22*(1), 148-169.

Some New Sequence Spaces of Generalized Fractional Order

[1] Muhammed Çınar ; [2]Mikail ET and [3]Koray İ. ATABEY

[1] Department of Mathematics Education, Mus Alparslan University, Mus-TURKEY

[2]Department of Mathematics, Firat University, Elazig-TURKEY

[3] Muş Nizamülmülk Kız Anadolu İmam Hatip High School, Mus-TURKEY

E-mail: [1]muhammedcinar23@gmail.com

[2]mikailet68@gmail.com ; [3]korayatabey7@gmail.com

Abstract

In this study, by using the generalized difference operator, we introduce the new fractional difference sequence spaces $c(B^{\widetilde{\alpha}})$ and $c_0(B^{\widetilde{\alpha}})$. We also investigate their topological properties and compute α−, β− , γ− duals and matrix transform.

Keywords: Difference Operator, Dual Space, Matrix Transform.

REFERENCES

1. Baliarsingh, P. 2013. Some new difference sequence spaces of fractional order and their dual spaces, Appl. Math. Comput. , 219(18), 9737–9742.
2. Başarir, M. and Kayıkçı, M. 2009. On the generalized B^m-Riesz difference sequence space and β-property. Journal of Inequalities and Applications.
3. Başar, F. 2012. Summability theory and its applications. Bentham Science Publishers.
4. Et, M. and Çolak, R. 1995. On some generalized difference sequence spaces. Soochow journal of mathematics, 21(4), 377-386.
5. Et, M. and Esi, A. 2000. On Köthe-Toeplitz duals of generalized difference sequence spaces. Bull. Malays. Math. Sci. Soc., 23(1), 25-32.

MARSHALL OLKİN GARİMA DISTRUBUTİON AND STATİSTİCAL PROPERTIES

Ayşe Metin Karakaş[a], Sinan ÇALIK [b], Aslıhan DEMİR[c]

[a]*Department of Statistics, Bitlis Eren University, Bitlis, Turkeyb*

[b]*Department of Statistics, Firat University, Elazığ, Turkey*

[c]*Department of Statistics, Firat University, Elazığ, Turkey*

aysekarakas5767@gmail.com

Abstract

A two-parameter Marshall-Olkin-Garima distribution is introduced and its structural, statistical properties are investigated. These include the compounding representation of the distribution, the shapes of the density and hazard rate functions, the moments and quantiles as well as the limiting distributions of extreme order statistics. Afterwards, the superiority of Marshall Olkin Garima distribution over the other known distributions on the real data set was shown.

.

Keywords:. Marshall-Olkin Distrubution, Garima distribution.

REFERENCES

1. Muliere, P., Scarsini, M.. Characterization of a Marshall-Olkin type class of distributions. Annals of the Institute of Statistical Mathematics, 39(2), 429-441, 1987.

A NUMERICAL APPROACH FOR SAWADA-KOTERA EQUATION WITH TRIGONOMETRIC QUINTIC B-SPLINE COLLOCATION METHOD

Hatice Karabenli [1], **Alaattin Esen** [1], **N. Murat Yağmurlu** [1]
[1] Department of Mathematics, University of Inonu, Malatya, Turkey

haticekarabenli@gmail.com, alaattin.esen@inonu.edu.tr, murat.yagmurlu@inonu.edu.tr

Abstract

In this paper, we deal with the numerical solution of Sawada-Kotera (SK) equation classified as the type of fifth order Korteweg-de Veries (gfKdV) equations. In the first step of our work consisting of several steps, nonlinear model problem is degraded by using $w_{xxx} = v$ and obtained the system with the coupled new equations. In the second step, Rubin-Graves type linearization are used to get rid of nonlinearity problem. After these applications, approximate solutions are obtained by using the trigonometric quintic B-Spline collocation method. The effiency and accuracy of the present method is demostrated with the tables and graphs. As seen as the tables given with L_2 and L_∞ error norms for different time and space steps, the present method is more accurate for the larger element numbers and smaller time steps.

Keywords: Sawada-Kotera Equation, Collocation Finite Element Method, Trigonometric Quintic B-Spline Basis, Rubin-Graves Type Linearization.

REFERENCES

1. K. Sawada and T. Kotera, A method for finding N-soliton solutions of the KdV equation and KdV-like equation, Progress of Theoretical Physics, No: 51, 1355–1367,1974.

2. H. O. Bakodah, Modified Adomian Decomposition Method for the Generalized Fifth Order KdV Equations, American J. of Comput. Mathematics, No:3, 53-58, 2013.

3. K. Djidjeli, W. G. Price, E. H. Twizell and Y. Wang, "Numerical Methods for the Soltution of the Third and Fifth-Order Dispersive Korteweg-De Vries Equations," Journal of Computational and Applied Mathematics, Vol: 58, No: 3, 307-336, 1995.

4. P. M. Prenter, Splines and Variational Methods, New York, John Wiley, 1975.

5. I. J. Schoenberg, On trigonometric spline interpolation, J. Math. Mech,Vol: 13, 795, 1964.

6. S.G. Rubin and R.A. Graves, A Cubic Spline Approximation for Problems in Fluid Mechanics, NASA, Washington, D.C., October, 1975.

7. N. M. Yagmurlu, O. Tasbozan, Y. Ucar and A. Esen, Numerical solutions of the Combined KdV-mKdV Equation by a Quintic B-spline Collocation Method, Appl. Math. Inf. Sci., V: 4, No:1, 19-24, 2016.

A COLLOCATION FINITE ELEMENT SOLUTION FOR THE FRACTIONAL FOKKER-PLANCK PROBLEM

Hatice Karabenli[1], **Alaattin Esen**[1], **Yusuf Uçar**[1]
[1] Department of Mathematics, University of Inonu, Malatya, Turkey

haticekarabenli@gmail.com, alaattin.esen@inonu.edu.tr, yusuf.ucar@inonu.edu.tr

Abstract

In this study, the numerical solutions of the fractional Fokker-Planck equations have been investigated. Firstly, the general finite element schemes have been obtained by using collocation finite element method suited with the trigonometric quintic B-spline basis functions. Then, the present method has been tested on two fundamental problems having different boundary conditions. The newly numerical results contained error norms L_2 and L_∞ are compared with the exact ones and other solutions for the different time steps and space steps. It is seen that from the tables and graphics, the numerical solutions are compatible with the exact solutions. Also, it can be said that more accurate solutions are avaliable for the larger element numbers and smaller time steps.

Keywords: Fokker-Planck Equation, Collocation Finite Element Method, Trigonometric Quintic B-Spline Basis Functions.

REFERENCES

1.W.Wyss, The fractional diffusion equation, Journal of Mathematical Physics, Vol: 27, No: 11, 2782-2785, 1986.

2. H. Risken, The Fokker-Plank Equation: Method of Solution and Applications, Springer Verlag, Berlin, 1989.

3. F.Neville, X. Jingyu and Y. Yubin, A finite element method for time fractional partial differential equations. Fractional Calculus and Applied Analysis, Vol: 14, No: 3, 454-474, 2011.

4. L.Dongfang, L.Hong-lin, S.Weiwei, W.Jilu and Z. Jiwei, Analysis of L1-Galerkin FEMs for Time-Fractional Nonlinear Parabolic Problems, Communications in Computational Physics, V:24, No:1, 86-103, 2018.

5. N. M. Yagmurlu, O. Tasbozan, Y. Ucar and A. Esen, Numerical solutions of the Combined KdV-mKdV Equation by a Quintic B-spline Collocation Method, Appl. Math. Inf. Sci., V: 4, No:1, 19-24, 2016.

6. I. J. Schoenberg, On trigonometric spline interpolation, J. Math. Mech., Vol. 13, 795, 1964.

THE RELATIONSHIP OF BILINGUALITY OF STUDENTS WITH THE ACHIEVEMENT OF SCIENCE COURSE

Kader Öğmen[1]

[1] Faculty of Education Department of Mathematics and Science Education, University of Ordu, Ordu, Turkey

kkaderogmen@gmail.com,

Abstract

In this study, 67 middle school 5th grade students whose mother tongue is different from the language they are taught in Çermik district of Diyarbakır in the 2019-2020 academic year and are educated with their mother tongue in the town of Camas, Ordu. In the study, the success of the Science course and reading comprehension of the students studying in their mother tongue and those studying in a language other than their mother tongue were examined in the study. It was revealed that gender does not have an effect on the reading comprehension and science achievement of monolingual and bilingual students. It was concluded that there is a positive and hing level relationship between the success of the science courses of students with different native languages and their reading comprehension.

Keywords: Mother Tongue, Monolingualism, Bilingualism, Science Education

REFERENCES

1. Agourram, A. G. (2013). Phenolic content, antioxidant potential and antimicrobial activeites of fruit and vegetable by-produckt extracts. *International Journal of Food Properties, 16*(5), 1092-1104.
2. Akbalık, S., & Yılayaz, Ö. (2020). Ortaokul 8. sınıf öğrencilerinin okuduğunu anlama ve fen başarılarının bazı değişkenler açısından incelenmesi. *Turkish Journal of Educational Studies*, *7*(3), 119-132.
3. Alkaya, Y., & Köysüren, D. (2017). *Gelişim Psikolojisi.* Ankara: Lider Yayınları.
4. Altınok, M. A., Tunç, T., & Özcan, H. (2020). Fen öğretim programlarının fen-teknoloji- toplum ve çevre kazanımları bağlamında 1926'dan günümüze karşılaştırmalı incelemesi. *Amasya Üniversitesi Eğitim Fakültesi Dergisi*, *9*(2), 230-257.
5. Altunkaya, H. (2015). Yabancı dil olarak Türkçe öğrenenlerin okuma kaygıları ile okuduğunu anlama becerileri arasındaki ilişki. Doktora Tezi, İnönü Üniversitesi, Eğitim Bilimleri Enstitüsü, Türkçe Eğitimi Ana Bilim Dalı, Malatya.
6. Arıcı, A. F., & Taşkın, Y. (2019). Okuma becerisinin diğer dil becerileri ile ilişkisi. *International Journal of Field Educations*, *5*(2), 185-194.

ON THE MODIFIED ORTHOGONAL FRAMES OF THE NON-UNIT SPEED CURVES IN EUCLIDEAN SPACE E^3

Sümeyye Gür Mazlum [1] , **Mehmet Bektaş** [2]

[1] Department of Computer Technology, University of Gümüşhane, Gümüşhane, Turkey
sumeyyegur@gumushane.edu.tr

[2] Department of Mathematics, University of Firat, Elazig, Turkey
m.bektas@firat.edu.tr

Abstract

In this study, the modified frames with both the non-zero curvature and the torsion of the non-unit speed curves in Euclidean space E^3 are examined. The relationships between the derivative vectors of the modified frame and the Frenet vectors or the vectors of the modified frame of the curve are given. Besides, the Darboux vectors obtained from the modified orthogonal frames with both the curvature and torsion of the curve and the unit vectors in the direction of these Darboux vectors are investigated. Finally, all these results are shown on the sample curves.

Keywords: Modified orthogonal frame, Frenet frame, Darboux vector

REFERENCES

1. Arikan, M., Nurkan, S. K., Adjoint Curve According to Modified Orthogonal Frame with Torsion in 3-Space, Usak University Journal of Science and Natural Sciences, 2 (2020), 54-64.
2. Azak, A. Z., Involute-Evolute Curves According to Modified Orthogonal Frame, Journal of Science and Arts, 21(2) (2021), 385-394.
3. Bükcü, B., Karacan M. K., On the modified orthogonal frame with curvature and torsion in 3-space, Mathematical Sciences and Applications E-Notes, 4(1) (2016), 184-188.
4. Clelland, J. N., From Frenet to Cartan: the method of moving frames, American Mathematical Soc, 178 (2017).
5. Eren, K., Kosal, H. H., Evolution of space curves and the special ruled surfaces with modified orthogonal frame, AIMS Mathematics, 5(3) (2020), 2027{2039.
6. Gray, A., Modern Differential Geometry of Curves and Surfaces with Mathematica, CRC, New York, (1998).
7. Lone, M. S., Es, H., Karacan, M. K., Bukcu, B., On some curves with modified orthogonal frame in Euclidean 3-Space, Iranian Journal of Science and Technology, Transactions A: Science, 43(4) (2019), 1905-1916.
8. Lone, M. S., Es, H., Karacan, M. K., Bukcu, B., Mannheim curves with modified orthogonal frame in Euclidean 3-space, arXiv e-print, 1809 (2018).
9. Lone, M. S., Karacan, M. K., Tuncer Y., Es, H., Spherical indicatrices with the modified orthogonal frame, arXiv preprint, arXiv: 1810.01082 (2018).
10. Özdemir, M., Diferansiyel Geometri, Altin Nokta Yayinevi, (2020).
11. Sasai, T., The fundamental theorem of analytic space curves and apparent singularities of Fuchsian differential equations, Tohoku Math Journal, 36 (1984), 17-24.
12. Yayli, Y., Gök, I, Hacisalihoglu, H. H., Extended rectifying curves as new kind of modified Darboux vectors, TWMS J Pure Appl Math, 9 (2018), 18-31.

SOFT TOPOLOGICAL HYPERVECTOR SPACES

Gülay Oğuz[1]

[1] Department of Mathematics, University of Harran, Şanlıurfa, Turkey

gulay.oguz@harran.edu.tr

Abstract

The aim of this paper is to introduce soft topological hypervector spaces, a new soft topological structure, by presenting a topological approach to the theory of hyperstructures, which is a generalization of classical theory, and soft set theory, which is one of the powerful mathematical theories that models uncertainty. In this perspective, some important results are obtained by studying its different properties. In addition, the relationships between hypervector spaces, soft hypervector spaces, topological hypervector spaces and soft topological hypervector spaces are examined in detail.

Keywords: Soft set; hypervector space; topological vector space; topological hypervector space; soft topological hypervector space.

REFERENCES

1. F. Marty, Sur une generalization de la notion de groupe, 8th congress des Mathematiciens Scandinaves, Stockholm 45-49 (1934).
2. D. Molodtsov. Soft set theory-first results. Computers Mathematics with Applications, 37(4-5), 19-31, (1999).
3. G. Oguz, B. Davvaz. Soft Topological Hyperstructure. Journal of Intelligent & Fuzzy Systems, 40(5), 8755-8764 (2021).
4. G. Oguz. On Soft Topological Hypergroups. Journal of Hyperstructures, 9(2), 81-95 (2020).
5. E. Ranjbar-Yanehsari, M. Asghari-Larimi, R. Ameri. Soft Hypervector Space and Fuzzy Soft Hypervector Space. *European Journal of Pure and Applied Mathematics*, *12*(1), 118-134 (2019).

NEW NUMERICAL SCHEME FOR SOLVING OF SINGULARLY PERTURBED PROBLEM WITH NONLOCAL BOUNDARY CONDITIONS

Derya Arslan

Department of Mathematics, University of Bitlis Eren, Bitlis, Turkey

[2]ayredlanu@gmail.com.,

Abstract

We consider the singular perturbed problem with nonlocal boundary value on Bakhvalov type mesh. We show that the scheme converges in the first order at the discrete maximum norm independent of the perturbation parameter. The effectiveness of the proposed method is supported by the results of applying the method to a singularly perturbed problem. As a result, it is seen that the numerical operation supports the theoretical results.

REFERENCES

1. Samarskii, A. A., Theory of Difference Schemes, M. Nauka, Moscow, Russia, 1971.
2. Nayfeh, A. H., Introduction to Perturbation Techniques, John Wiley & Sons, New York, NY, USA, 1993.
3. O'Malley, R. E., Singular Perturbation Methods for Ordinary Differential Equations, vol. 89 of Applied Mathematical Sciences, Springer, New York, NY, USA, 1991.

CONTENT ANALYSIS OF THE THESES ABOUT MATHEMATICAL PATTERN IN TURKISH CONTEXT

Fatma Erdoğan [1], **Sude Ay**[2]

[1] Department of Mathematics Education, University of Firat, Elazig, Turkey

[2]Ministry of National Education, Elazig, Turkey

f.erdogan@firat.edu.tr , suudeay35@gmail.com

Abstract

It is a fact that mathematical knowledge is based on patterns and structures (Mulligan and Mitchelmore, 2009). In this study, it was aimed to make an examination from a thematic aspect on the theses which were prepared on mathematical pattren in Turkey. For this purpose, 24 postgraduate theses about mathematical pattern in mathematics education were examined. Theses were examined through document analysis in terms of variables such as type, year, sample type, sample size, method, data collection tools, research topic. In the study, the data were analyzed by content analysis method. Descriptive statistics such as frequency and percentage were used. As a result of the research, it was found out that the number of master's theses was more. It was seen that 2017 was the year when the highest number of thesis were prepared. Qualitative methods were predominantly used in the theses studied. It was determined that a lot of studies were carried out with middle school students. Students' ability to generalize patterns and the effect of different learning methods on students' pattern success are among the most frequently researched topics. In line with the results of the research, some suggestions were made to the researchers considering to work in this field.

Keywords: Mathematical pattern, Postgraduate theses, Mathematics education

REFERENCES

1. J. T. Mulligan and M. C., Awareness of Pattern and Structure in Early Mathematical Development, Mathematics Education Research Journal, Vol:21 No:2, 33–49, 2009.

A NEW TASK FOR MATHEMATICALLY GIFTED STUDENTS: CONSECUTIVE NUMBERS TRIANGLE

Fatma Erdoğan [1], **Neslihan Gül**[2]

[1] Department of Mathematics Education, University of Firat, Elazig, Turkey

[2]Ministry of National Education, Elazig, Turkey

f.erdogan@firat.edu.tr, gulneslihan85@gmail.com

Abstract

One of the characteristic features of mathematically gifted students is their mental effort on mathematical and complex tasks (Leikin, Leikin, Paz-Baruch, Waisman, and Lev, 2017). In addition, gifted students prefer challenging tasks in learning environments (Wu, Jen, and Gentry, 2018). In this context, in this study, it is aimed to design a new number model consisting of counting numbers based on Pascal's Triangle, on which mathematical gifted students can work, and to examine the mathematical structure of this model. In the study, counting numbers were arranged sequentially in a triangular structure in the Excel Program. Later, the special triangular structure was called the "Consecutive Numbers Triangle". When the number groups were examined in the resulting model, it was discovered that each of the numbers was formed according to a rule. In addition, some special numbers (multiples of 2, 3, 4, 5, 10, square and prime numbers) were determined and the patterns formed by coloring only these numbers in the model were determined. In order to determine the patterns exactly, the numbers were written up to 1087 in the model. It was seen that a magnificent pattern was formed for each special issue. Since groups of numbers other than prime numbers are formed with a certain rule, the images of these numbers are formed with a regularity.

Keywords: Pascal's Triangle, Consecutive numbers, Mathematical giftedness, Mathematics education

REFERENCES

1. R. Leikin, M. Leikin, N. Paz-Baruch, I. Waisman, and M. Lev, On the Four Types of Characteristics of Super Mathematically Gifted Students, High Ability Studies, Vol:28, No:1, 107-125, 2017.
2. J. Wu, E. Jen, and M. Gentry, Validating a Classroom Perception Instrument for Gifted Students in a Universitybased Residential Program, Journal of Advanced Academics, Vol:29, No:3, 195–215, 2018.

Analysis of The Peer Bullying in Schools Considering by Fractional Order Metapopulation Modelling

Pelin Yaprakdal-Uzun[1], Kıvanç Uzun[2], İlknur Koca[3]

[1]Burdur Mehmet Akif Ersoy University, Burdur, Turkey, pelinyaprakdal@gmail.com

[2]Burdur Mehmet Akif Ersoy University, Burdur, Turkey, kuzun@mehmetakif.edu.tr

[3]Mugla Sitki Kocman University, Mugla, Turkey, ilknurkoca@mu.edu.tr

Abstract

Peer bullying, which has been up-to-date from past to present and continues to spread rapidly in schools, is a worrying problem on students' well-being and functionality. This problem in social sciences has been addressed in this study with the mathematical model considered by the metapopulation model. Mathematical models can be helpful in explaining a system, examining the effects of different components, and predicting behavior. Through the mathematical model considered in this context, it is aimed to determine the effects of the actors playing a role in peer bullying in schools and their relationship with each other, to determine the level of peer bullying and to obtain important findings about how to prevent bullying. While doing literature reviewing we realized that a comprehensive mathematical model regarding peer bullying, which is an extremely important concept for social sciences and students, has not been considered yet. Based on this deficiency in the literature, a realistic mathematical model of peer bullying in schools has been considered in our study. This mathematical model is handled with the help of the fractional operator, and first the local and global stability analysis of the equilibrium points of the model, then the positivity, existence and uniqueness of the solutions will be shown. According to us, this study is very important in terms of its original and multidisciplinary approach to a subject in the field of social sciences.

Keywords: Peer bullying; Fractional operator; Local stability; Global stability; Existence and uniqueness; Lyapunov function.

REFERENCES

1. Olweus, D. (1997). Bully/victim problems in school: Facts and intervention. *European Journal of Psychology of Education, 12*(4), 495-510.
2. Podlubny, I. (1998). *Fractional differential equations: An introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications.* Elsevier.
3. Toufik, M., & Atangana, A. (2017). New numerical approximation of fractional derivative with non-local and non-singular kernel: Application to chaotic models. *The European Physical Journal Plus*, *132*(10), 1-16.

Mathematical Modelling and Speed Control Optimization of PMSM Machine for Electric Vehicle Application

Megrini Meriem[1], Gaga Ahmed[1], Mehdaoui Youness[1]

[1]Research Laboratory of Physics and Engineers Sciences (LRPSI), Research Team in Embedded Systems, Engineering, Automation, Signal, Telecommunications and Intelligent Materials, Polydisciplinary Faculty (FPBM), Sultan Moulay Slimane University (USMS), Beni Mellal, Morocco

megrinimeriem@gmail.com

gaga.ahmad@gmail.com

youness.mehdaoui@gmail.com

Abstract

Electric vehicles are not something new in this world, however the manufacturers are trying to develop their properties. Since the battery and the electric motors are the essential parts of it, this paper is going to deal with the mathematical modelling and the control of electric motors, especially the permanent magnet synchronous machine (PMSM). The work has begun by using a mathematical model with this machine and analysing it. The first finding revealed that the most fundamental parameters of PMSM, torque and flux, are coupled with each other. Furthermore, we made use of a widely used control, which is FOC, in order to decouple the torque and the flux. We decoupled them by controlling the current as well, since the equation of all the elements is proportional. In the second result, we found an equation that combines speed and torque, in which the speed can also be controlled by the current or the inverse. The results of the two tests using this mathematical model and field oriented control algorithm (FOC) to control the speed of PMSM reveal that the second one is more optimised than the first. Differently, in the first one, we use three PID regulator, but we use only one in the second. These systems are simulated in MATLAB/SIMULINK environment. To conclude, the two tests rendered the same results: fast response and good reference speed tracking, unless the second is the most optimised and appropriate.

Keywords **–** Mathematical Modeling, PMSM, FOC, Speed control, PID regulator

REFERENCES

1. Advan, B.; Kuppuswamy, A.; Jayabaskaran G.; and Jagannathan, V. (2011). Field oriented control of permanent magnet synchronous motor (PMSM) using fuzzy logic controller. IEEE Recent Advances in Intelligent Computational System, 587-592.
2. P. Pillay and R. Krishnan, Control characteristics and speed controller design of a high performance PMSM, in Record IEEE Ind. Appl. Soc.Annu. Meeting., 1985, pp. 627–633.
3. C. Wang, Y. Ji, H. Luan, and Z. Zhang, “Simulation of PMSM sensorless vector control system based on Matlab/Simulink”, Journal of Jilin University (Information Science Edition), vol. 27, no. 1, pp. 17-22, 2009

Analysis of The Vectored Plant Disease Model Considering by Fractional Operator

Pelin Yaprakdal-Uzun [1]

[1] Burdur Mehmet Akif Ersoy University, Burdur, Turkey

pelinyaprakdal@gmail.com

Abstract

Mathematical models are used in social sciences (economics, psychology, sociology, political science, etc.) as well as in natural sciences (physics, biology, earth science, meteorology, etc.) and engineering disciplines (computer science, artificial intelligence, etc.). Plant models of plant-virus and plant-vector-models are very useful for general idea of disease incidence class and general plant virus. This product is similar to a crop from field-harvested cuttings, the second plant-v-vector pattern elevators. This mathematical model is handled with the help of the fractional operator, and first the local and global stability analysis of the equilibrium points of the model, then the positivity, existence and uniqueness of the solutions will be shown.

Keywords: Vectored plant disease; Fractional operator; Local stability; Global stability; Existence and uniqueness; Lyapunov function.

REFERENCES

1. Podlubny, I. (1998). Fractional differential equations: an introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications. Elsevier.
2. Bokil, V. A., Allen, L. J. S., Jeger, M. J., & Lenhart, S. (2019). Optimal control of a vectored plant disease model for a crop with continuous replanting. Journal of Biological Dynamics, 13(sup1), 325-353.

On the Topologies in Dual Space and the Analyticity of Dual Functions

Olgun Durmaz[1], **Buşra Aktaş**[2], **Halit Gündoğan**[3]

[1]Department of Mathematics, University of Atatürk, Erzurum, Turkey

[2,3]Department of Mathematics, University of Kırıkkale, Kırıkkale, Turkey

olgun.durmaz@atauni.edu.tr

baktas6638@gmail.com

hagundogan@hotmail.com

Abstract

In this paper, the properties of the order relation on dual numbers are examined in detail. Using the order relation $<_D$, we obtain the topologies on D^n denoted by $\bar{\tau}_{\bar{d}}$ and $\bar{\tau}$ such that the spaces $(D^n, \bar{\tau}_{\bar{d}})$ and $(D^n, \bar{\tau})$ are Hausdorff spaces. After then, how the analyticity conditions of a dual function which is often expressed in other studies are obtained is given clearly. Making use of the topology $\bar{\tau}_{\bar{d}}$, the dual analytic areas of dual analytic functions are determined.

Keywords: Dual numbers; Dual Inequalities; Topology; Dual Analytic Functions.

REFERENCES

1. E. Halmaghi, P. Liljedahl, Inequalities in the History of Mathematics, From Peculiarities to a Hard Discipline, GeoGebra International Journal of Romania, Vol:4, No:2, 43-56, 2015.

2. G.H. Hardy, J.E. Littlewood, G. Polya, Inequalities, Cambridge Univ. Press., 1934.

3. F.M. Dimentberg, The Screw Calculus and its Applications in Mechanics, Foreign Technology Division, Wright-Patterson Air Force Base, 1965.

4. W.B.V. Kandasamy, F. Smarandache, Dual Numbers, ZIP Publishing, Ohio, 2012.

5. J.R. Munkres, Topology, The Second Edition, ISBN 0-13-181629-2, 2000.

6. G.R. Veldkamp, On the Use of Dual Numbers, Vectors and Matrices in Instantaneous Spatial Kinematics, Mechanism and Machine Theory Vol:11, 141-156, 1976.

7. H. H. Hacısalihoğlu, Hareket Geometrisi ve Kuaterniyonlar Teorisi, Gazi ¨Universitesi Fen Fakültesi Yayınları, Ankara, 1983.

8. I. Fischer, Dual-Number Methods in Kinematics, Statics and Dynamics, CRC Press, 1998.

EXAMINING SECONDARY MATHEMATICS TEACHERS' OPINIONS ABOUT THE EMERGED PROBLEMS DURING THE TEACHING OF MATHEMATICS

Serbay DURAN [1]

[1] Department of Mathematics and Science Education, Adıyaman University, Adıyaman, Turkey

sduran@adiyaman.edu.tr,

Himmet KORKMAZ [2]

[2] Department of Mathematics and Science Education, Ordu University, Ordu, Turkey

himmetkorkmaz@odu.edu.tr,

Abstract

The aim of this study is to examine secondary mathematics teachers' thinking about the problems that arise in the process of teaching mathematics. It is important that the in-service mathematics teachers need to maintain their professional development to keep up with the new era from several perspectives [1]. In this respect, the Ministry of National Education carries out several professional development programs to support teachers' professional development. However, teachers may still experience some troubles in the in-class teaching process stemming from students or themselves. This study, it is aimed to identify the problems experienced by secondary mathematics teachers in the in-class mathematics teaching process and to provide recommendations to cope with them. The current study is a case study that is one of the qualitative research methods. The participants consisted of 12 secondary mathematics teachers working in high schools in a province in the Southeastern Anatolia Region of Turkey who are selected by the purposeful sampling method. The questionnaire created by the researchers was applied to the teachers separately by semi-structured interview technique. The descriptive analysis method was used in the analysis of the obtained raw data. When analyzing the data, codes obtained from the literature and the findings of the research formed themes. The findings showed that teachers expressed frequently the problems such as a low readiness level of students, lack of knowledge in course sources, frequent use of traditional teaching methods rather than new approaches, and typically use of traditional questions during the in-class teaching. Moreover, teachers made suggestions to overcome these troubles.

Keywords: Secondary mathematics teachers; Problems; Mathematics teaching.

REFERENCES

1. Arbaugh, Fran. Study Groups as A Form of Professional Development for Secondary Mathematics Teachers. Journal of Mathematics Teacher Education Vol: 6, No: 2, 139-163, 2003.

FRACTIONAL OPTIMAL CONTROL OF AN EPIDEMIC MODELED WITH TWO TYPES OF NONLINEAR INCIDENCE RATES

Derya AVCI[1,*] **& Fatma SOYTÜRK**[2]

[1] Department of Mathematics, Balıkesir University, Balıkesir, Turkey

[*] dkaradeniz@balikesir.edu.tr,

[2] Department of Mathematics, Balıkesir University, Balıkesir, Turkey

fatmastrk@gmail.com,

Abstract

This study aims to propose the optimal treatment method for an epidemic under the effect of different nonlinear incidence rates [1]. This rate represents the relationship between infected and susceptible individuals, and also makes it possible to predict the progression of disease starting to spread in a population. Depending on the nature of the disease, it can be described by various functional relations. The model under discussion is in terms of the Caputo fractional derivative [2]. The main purpose is to compare the effects of two nonlinear incidence rates on the controlled system. Numerical solutions are simulated with MATLAB software. Comparisons of optimal solutions according to the problem parameters are given in the graphics.

Keywords: Epidemic model; Fractional optimal control; Caputo fractional derivative; Incidence rate

REFERENCES

1. Amira Rachah, Analysis, simulation and optimal control of a SEIR model for Ebola virus with demographic effects, Communications Faculty of Sciences University of Ankara Series A1 Mathematics and Statistics, Vol:67, No:1, 179-197, 2018.
2. Balram Dubey, Preeti Dubey, Uma S Dubey, Dynamics of an SIR model with nonlinear incidence and treatment rate, Applications and Applied Mathematics: An International Journal (AAM), Vol:10, No:2, 718-737, 2015.

FURTHER ON k-FRACTIONAL OPERATORS INVOLVING MITTAG-LEFFLER FUNCTIONS FOR STRONGLY $(\alpha, h-m)$-CONVEX FUNCTIONS

Ayşe Kübra DEMİREL [1]

[1] Central Research Laboratory, University of Ordu, Ordu, Turkey

aysekubrademirel@gmail.com,

Abstract

Integral operators, which form a significant part of fractional calculus, are resources in numerous areas like theory of inequality, statistics, mathematical biology, modelling and engineering which take profit from the fractional analysis. Many inequalities have led to the creation of generalized and new approaches through fractional integral operators.

The purpose of this article is to improve the bounds of the k-fractional integral operators defined for strongly $(\alpha, h-m)$-convexity. To obtain the required results, extended generalized Mittag-Leffler functions are used. The results introduced in this article can be considered an important advancement of previously published conclusions.

Keywords: Mittag-Leffler function; Strongly $(\alpha, h-m)$-convex function; k-fractional integrals.

REFERENCES

1. Yanliang Dong, Maryam Saddiqa, Saleem Ullah, Ghulam Farid, Study of fractional integral operators containing Mittag-Leffler functions via strongly (α, m) -convex functions, Mathematical Problems in Engineering, Vol:2021, Article ID 6693914, 2021.
2. Havva Kavurmacı Önalan, Ahmet Ocak Akdemir, Merve Avcı Ardıç, Dumitru Baleanu, On new general versions of Hermite-Hadamard type integral inequalities via fractional integral operators with Mittag-Leffler kernel, Journal of Inequalities and Applications, Vol:2021, No:186, 1–16, 2021.
3. Tao Yan, Ghulam Farid, Sidra Bibi, Kamsing Nonlaopon, On Caputo fractional derivative inequalities by using strongly $(\alpha, h-m)$-convexity, AIMS Mathematics, Vol:7, No:6, 10165–10179, 2022.
4. Saad Ihsan Butt, Saba Yousaf, Ahmet Ocak Akdemir, Mustafa Ali Dokuyucu, New Hadamard-type integral inequalities via a general form of fractional integral operators, Chaos, Solutions & Fractals, Vol:2021, No:148, 111025, 2021.

EXAMINATION OF 8TH GRADE STUDENTS' OPINIONS ON NEW GENERATION QUESTIONS

Yavuz TURAN[1]

yvztrn@hotmail.com,

Hayal YAVUZ MUMCU[2]

hayalym52@gmail.com,

[1,2]Ordu University, Ordu, Turkey

Abstract

The aim of this study is to examine the opinions of 8th grade students about new generation questions. In the study, the phenomenology method, which examines the subjective experiences of people with the event or situation they live, was used. The sample of the study consist of 8th grade students in Ordu province. The convenient sampling technique (Yıldırım & Şimşek, 2018) which not only adds speed and practicality to the research, but also allows the researcher to choose a situation that is close and easily accessible was used in the study, since the students in the study group consist of the students of the teacher conducting the research. In the study, the Interview Form developed by the researchers was used as data collection tool. In this form, there are open-ended questions to reveal students' views on new generation questions. As a result of the study, it was seen that the students defined the new generation questions as long, challenging, intertwined problems, and boredom. In addition, most of the students stated that they did not encounter these question types until they passed the 8th grade, and many students stated that they had difficulty in solving these question types. Most of the students stated that such questions are unnecessary and that instead of these questions, questions about acquisitions in the curriculum should be asked in the exams. Students think that new generation questions should be simplified. According to the results obtained from the study, it was seen that most of the students did not know why such questions were included in the curriculum. Besides, they said that they were alienated from the mathematics because of these questions, they gave up studying mathematics and they did not want to choose a profession related to mathematics in the future. In the light of the results obtained from the study, suggestions were made for different studies and learning environments that can be done on the subject.

Keywords: New generation questions; 8th grade students, Phenomenology study

REFERENCES

1. Yıldırım, A. ve Şimşek, H. (2008). Sosyal Bilimlerde Nitel Araştırma Yöntemleri. Ankara: Seçkin Yayıncılık.

An optimal control strategy for an H1N1 model

Beyza Billur İskender Eroğlu [1] **and Dilara Yapışkan** [2]

[1,2] Department of Mathematics, Balikesir University, Balikesir, Turkey

[1] biskender@balikesir.edu.tr,

[2] dilara.yapskn@baun.edu.tr,

Abstract

In this work, control strategies to avoid the spread of the H1N1 virus with the three-compartment SIR model are discussed. For this, two control terms, meaning precaution (mask, social distance, hygiene, etc.) and vaccination, were adapted to a without control model [1]. The purpose of these controls is to decline the number of susceptible and infected individuals. First, necessary optimality conditions are obtained for optimal control of disease spreading [2]. Then, numerical solutions of the optimal system are found by applying the forward-backward fourth-order Runge-Kutta method. The dynamics of the controlled and uncontrolled models are simulated using the MATLAB program and the controls decreased the number of susceptible and infected individuals with the desired target.

Keywords: Optimal control, H1N1, SIR epidemiological model, Fourth-order Runge-Kutta Method.

REFERENCES

1. Fathalla A Rihan, Mohamed Naim Anwar, Mohamud Sheek-Hussein, and Srdjan Denic, SIR Model of Swine Influenza Epidemic in Abu Dhabi, Journal of Public Health Frontier, Vol:1, No:4, 85-89, 2012.
2. Donald E. Kirk, Optimal control theory: an introduction, Courier Corporation, 2004.

STOCHASTIC DIFFERENTIAL EQUATION MODELING FOR GOLD DATA WITH YUIMAGUI

Sevda Özdemir[1], **Fevzi Erdoğan**[2]

[1]Ozalp Vocational School, University of Van Yuzuncu Yil, Van, Turkey
[2]Department of Econometrics, University of Van Yuzuncu Yil University, Van, Turkey

sevdaozdemir@yyu.edu.tr, ferdogan@yyu.edu.tr

Abstract

In this study, GOLD data versus USD is investigated by Stochastic Differential Equation Modeling (SDEM). First of all, Square Root Process, Geometric Brownian Motion (GBM), Vasicek Model (VAS), and Cox-Ingersoll-Ross (CIR) models have selected from the most preferred models in finance. Then, the parameters of the mentioned SDE models have estimated using the quasi maximum likelihood method. Secondly, It has been given the p-value for the Kolmogorov-Smirnov test that checks if the empirical and theoretical distribution are the same to see the goodness of fit of the model. Accordingly, all selected models fit well with the given data. After that, model selection has been made among these 4 compatible models according to AIC and BIC criteria. Therefore, for the given data, the VAS model is the most appropriate model according to both criteria. Finally, by using the VAS model, which has been chosen as the most appropriate model, and applying Euler-Maruyama Approximation Method future simulation trajectories of 100 steps between 25.04.2022 and 25.04.2023 have been obtained. Gold opening data with the symbol GC=F between 02.01.2020 and 25.04.2022 have obtained with the help of https://finance.yahoo.com/ link, with the YUIMAGUI interface of the RSTUDIO program and all results have also made by using YUIMAGUI. These results are also corroborated by graphical representation.

Keywords: stochastic differential equation , quasi maximum likelihood, Euler-Maruyama aroximation method,square root process, CIR, GBM, VAS, YUIMAGUI.

REFERENCES

1. Guidotti, Emanuele. YUIMAGUI. 27 04 2022 <https://guidotti.shinyapps.io/yuimaGUI/>.
2. Iacus, Stefano M. ve Nakahiro Yoshida. Simulation and Inference for Stochastic Processes with YUIMA. Springer, 2018.
3. Mao, Xuerong. Stochastic Differential Equatiion and Applications. United Kingdom: Woodhead Publishing, 2007.
4. Sauer, T. Numerical solution of stochastic differential equations in finance. In Handbook of Computational Finance. Berlin, Heidelberg: Springer, 2012.

P-SASAKIAN MANIFOLDS AND THEIR CURVATURE CLASSIFICATION

PAKIZE UYGUN*, SÜLEYMAN DIRIK AND MEHMET ATÇEKEN

Abstract. The object of this paper is to study the curvature tensors of P-Sasakian manifold satisfying the conditions $W_1^* \cdot S = 0$, $W_1^* \cdot P = 0$, $W_1^* \cdot W_9 = 0$ and $W_1^* \cdot W_1^* = 0$. According these cases, P-Sasakian manifolds have been characterized such as η-Einstein and Einstein. In addition, we research W_1^*-flat and W_9-flat a P-Sasakian manifold. The results obtained are interesting and give an idea about the geometry of P-Sasakian manifold.

References

[1] T. Adati and T. Miyazawa, Some properties of P-Sasakian manifolds, TRU Math. 13 (1977), 25-32.
[2] T. Adati and T. Miyazawa, On P-Sasakian manifolds satisfying certain conditions, Tensor (N.S.), 33 (1979), 173-178.
[3] U.C. De and D. Tarafdar, On a type of P-Sasakian manifold, Math. Blkanica (N.S), 7 (1993), 211-215.
[4] D. Kowalezyk., On some subclass of semi-symmetric manifolds, Soochow J. Math., 27 (2001), 445-461.
[5] T. Mert, Characterization of some special curvature tensor on Almost $C(a)$-manifold, Asian Journal of Math. and Computer Research, 29 (1) (2022), 27-41.
[6] T. Mert and M. Atçeken, Almost $C(a)$-manifold on $W_0^\star-$curvature tensor, Applied Mathematical Sciences, 15 (15) (2021), 693-703.
[7] B. O. Neill, Semi-Riemannian Geometry with Applications to Relativity, Academic Press, New York, 1983.
[8] C. Özgür and M. M. Tripathi, On P-Sasakian manifolds satisfying certain conditions on the concircular curvature tensor, Turk J. Math., 31 (2007), 171-179.
[9] G. P. Pokhariyal, Relativistic significance of curvature tensors, Internat. J. Math. Sci., 5 (1) (1982), 133-139.
[10] G.P. Pokhariyal and R.S.Mishra, Curvature tensors and their relativistic significance II, Yokohama Math. J., 19 (2) (1971), 97-103.
[11] B. Prasad, A pseudo projective curvature tensor on a Riemannian manifold, Bull. Calcutta Math. Soc., 94 (3) (2002), 163-166.
[12] I. Sato., On a structure similar to the almost contact structure, 30 (1976), 219-224.
[13] I. Sato and K. Matsumoto., On P-Sasakian manifolds satisfying certain conditions, Tensor (N.S), 33 (1979), 173-178.
[14] Z.I. Szabo, Structure theorems on Riemannian spaces satisfying $R(X,Y) \cdot R = 0$, I: The local version, J. Differential Geom., 17 (4) (1982), 531-582.
[15] M.M. Tripathi and P. Gupta, $T-$curvature tensor on a semi-Riemannian manifold, 4 (1) (2011), 117-129.
[16] P. Uygun and M. Atçeken, On (k,μ)-paracontact metric spaces satisfying some conditions on the $W_0^\star-$curvature tensor, NTMSCI, 9 (2) (2021) 26-37.
[17] K. Yano and S. Bochner, Curvature and Betti numbers, Annals of Mathematics Studies, Princeton University Press, 32 (1953).
[18] K. Yano and M. Kon, Structures on manifolds, Series in Pure Math., World Sci., 3 (1984).

1991 *Mathematics Subject Classification.* 53C15, 53C25.
Key words and phrases. P-Sasakian Manifold, η-Einstein manifold, W_1^* curvature tensor.

ON P-SASAKIAN MANIFOLDS SATISFYING CERTAIN CURVATURE CONDITIONS

PAKIZE UYGUN

Abstract. The aim of this paper is to study the curvature tensors of P-Sasakian manifold satisfying the conditions $W_1 \cdot W_5 = 0$, $W_1 \cdot W_6 = 0$, $W_1 \cdot W_7 = 0$ and $W_1 \cdot W_8 = 0$. Also, we study W_1-pseudo symmetry for a P-Sasakian manifold. We think that some interesting results on a P-Sasakian manifold are obtained.

References

[1] A. Atçeken and P. Uygun, Characterizations for totally geodesic submanifolds of (k, μ)-paracontact metric manifolds, Korean J. Math. 28(2020), 555-571.
[2] T. Adati and K. Matsumoto, On conformally recurrent and conformally symmetric P-Sasakian manifolds, TRU Math., 13 (1997), 25-32.
[3] T. Adati and T. Miyazawa, Some properties of P-Sasakian manifolds, TRU Math. 13 (1977), 25-32.
[4] T. Adati and T. Miyazawa, On P-Sasakian manifolds satisfying certain conditions, Tensor (N.S.), 33 (1979), 173-178.
[5] A. Yıldız, M. Turan and B. E. Acet, On para-Sasakian manifolds, Dumlupınar Uni. Fen bilimleri Enstitüsü Dergisi, 24 (2011).
[6] A. Barman, On para-Sasakian manifolds admitting semi-symmetric metric connection, Publication De linstıtut mathematique, 95 (2014), 239-47.
[7] U.C. De and D. Tarafdar, On a type of P-Sasakian manifold, Math. Blkanica (N.S), 7 (1993), 211-215.
[8] D. Kowalezyk., On some subclass of semi-symmetric manifolds, Soochow J. Math., 27 (2001), 445-461.
[9] T. Mert, Characterization of some special curvature tensor on Almost $C(a)$-manifold, Asian Journal of Math. and Computer Research, 29 (1) (2022), 27-41.
[10] T. Mert and M. Atçeken, Almost $C(a)$-manifold on $W_0^\star-$curvature tensor, Applied Mathematical Sciences, 15 (15) (2021), 693-703.
[11] B. O. Neill, Semi-Riemannian Geometry with Applications to Relativity, Academic Press, New York, 1983.
[12] C. Özgür and M. M. Tripathi, On P-Sasakian manifolds satisfying certain conditions on the concircular curvature tensor, Turk J. Math., 31 (2007), 171-179.
[13] G. P. Pokhariyal, Relativistic significance of curvature tensors, Internat. J. Math. Sci., 5 (1) (1982), 133-139.
[14] G.P. Pokhariyal and R.S.Mishra, Curvature tensors and their relativistic significance II, Yokohama Math. J., 19 (2) (1971), 97-103.
[15] B. Prasad, A pseudo projective curvature tensor on a Riemannian manifold, Bull. Calcutta Math. Soc., 94 (3) (2002), 163-166.
[16] I. Sato., On a structure similar to the almost contact structure, 30 (1976), 219-224.
[17] I. Sato and K. Matsumoto., On P-Sasakian manifolds satisfying certain conditions, Tensor (N.S), 33 (1979), 173-178.
[18] Z.I. Szabo, Structure theorems on Riemannian spaces satisfying $R(X, Y) \cdot R = 0$, I: The local version, J. Differential Geom., 17 (4) (1982), 531-582.
[19] M.M. Tripathi and P. Gupta, $T-$curvature tensor on a semi-Riemannian manifold, 4 (1) (2011), 117-129.

1991 *Mathematics Subject Classification.* 53C15, 53C25.
Key words and phrases. P-Sasakian Manifold, η-Einstein manifold, W_1-curvature tensor.

[20] P. Uygun and M. Atçeken, On (k,μ)-paracontact metric spaces satisfying some conditions on the $W_0^\star-$curvature tensor, NTMSCI, 9 (2) (2021) 26-37.
[21] P. Uygun, M. Atçeken and S. Dirik, Some curvature results on Kenmotsu metric spaces, Bull. Int. Math. Virtual Inst, 12 (2) (2022), 243-251.
[22] P. Uygun, M. Atçeken and S. Dirik, On Kenmotsu metric spaces satisfying some conditions on the W_1-curvature tensor, Filomat, (Kabul edildi).
[23] K. Yano and S. Bochner, Curvature and Betti numbers, Annals of Mathematics Studies, Princeton University Press, 32 (1953).
[24] K. Yano and M. Kon, Structures on manifolds, Series in Pure Math., World Sci., 3 (1984).

*Department of Mathematics, Faculty of Arts and Sciences, Tokat University, 60100, Tokat, Turkey
E-mail address: pakizeuygun@hotmail.com

OMAR KHAYYAM'S PLACE AND IMPORTANCE IN THE HISTORY OF MATHEMATICS

Hüseyin SAMANCI[1]

[1] Department of Arabic Teaching, University of Adiyaman, Adiyaman, Turkey

hsamanci@adiyaman.edu.tr,

Serbay DURAN[2]

[2]Department of Mathematics and Science Education, Adıyaman University, Adıyaman, Turkey

sduran@adiyaman.edu.tr,

Abstract

The aim of this study is to introduce Omar Khayyam and his works in terms of history of mathematics and mathematics education. Omar Khayyam is an Iranian Muslim scholar and is one of the most important mathematicians who contributed to this field by playing a major role in the advancement of mathematics in his time. Khayyam was interested in algebra, geometry, astronomy, physics and medicine, was engaged in music, and also wrote his rubais, poems that immortalized his name. Khayyam had a great influence on the development of mathematics in general and analytical geometry in particular. He showed that a cubic equation can have more than one solution, and how intersections of conic sections, such as parabolas and circles, can be used to obtain geometric solutions of cubic equations. He also proved for the first time that irrational numbers can be used like rational numbers in mathematical operations. In addition to these, Khayyam argued that algebraic facts emerged as geometric facts. Thus, he took an important step towards closing the gap between numerical and geometric algebra long before Descartes. With all these works, he contributed to the science of mathematics and guided the studies in the field of mathematics today.

Keywords: Omar Khayyam; Mathematics education.

REFERENCES

1. Viktor Katz, A History of Mathematics: An İntroductions, HarperCollins College Publishers, New York, 879, 1998.
2. Hazhir Teimourian, Omar Khayyam: Poet, Rebel, Astronomer, The History Press, Sutton, 569, 2008.
3. Yavuz Unat, Ömer Hayyam, İslam Ansiklopedisi, TDV, İstanbul, Vol: 34, 66-68, 2009.

A DOCUMENT ANALYSIS ON THE EFFECT OF FINLAND AND TURKISH EDUCATION SYSTEMS ON PISA SUCCESS

Zeynep Kübra Kazan, Assoc. Prof. Dr. Sevda Türkiş , Assist. Prof. Dr. Elif Çil

Department of Mathematics and Science Education, Faculty of Education, Ordu University, Ordu Turkey

kazanzeynepkubra@gmail.com , sevdaturkis@odu.edu.tr, elifcil@odu.edu.tr

The aim of the study examines the differences between the Finnish and Turkish education models and their impact on PISA success. By considering the factors behind the success in PISA exams, the similarities and differences between the two education models were determined with the document review method. It was aimed to make suggestions for increasing the success of PISA. A holistic approach is aimed at the Finnish curriculum, while a constructivist approach is aimed at the Turkish education model. Since 2018, it has been supported by the spiral approach in the Turkish curriculum. With the spiral approach, repetitive acquisitions and explanations at the different subject and grade levels are included, and learning outcomes that are holistic aim to be gained at once. It is planned to evaluate the effect of the Turkish education model, the constructivist approach model, and the Finnish education model on the science literacy of the Pisa exams.

Keywords: Finnish education system, Turkish education system, science literacy

REFERENCES

1. Benzer, A. (2019). Türkçe ders kitaplarının PISA okuma yeterlik düzeyleri ile imtihanı. *Okuma Yazma Eğitimi Araştırmaları*, *7*(2), 96-109.
2. Gürlen, E., Demirkaya, A. S., & Doğan, N. (2019). Uzmanların Pisa ve Timms sınavlarının eğitim politika ve programlarına etkisine ilişkin görüşleri. *Mehmet Akif Ersoy Üniversitesi Eğitim Fakültesi Dergisi*, (52), 287-319.
3. Özdemir, A. (2017). Bütün öğrencilerin okulu Finlandiya okulları. *İnformal Ortamlarda Araştırmalar Dergisi*, *2*(1), 59-91.
4. Yonca, Z. D. (2018). Finlandiya'nin Pisa Başarisina Etki Eden Faktörler Ve Türkiye Açisindan Karşilaştirilmasi. *Uluslararası Eğitim Bilimleri Dergisi*, (14), 136-146.
5. Yüksel, M. (2019). *PISA 2015 Türkiye ve FİnlandIya verilerine göre okul özellikleri ile öğrencilerin okuma becerileri arasındaki ilişkinin incelenmesi* (Master's thesis, Trakya Üniversitesi, Sosyal Bilimler Enstitüsü).

Some Asymptotic Features of Hyper-Airy Equation

Fatih Say

Department of Mathematics, Faculty of Arts and Sciences, Ordu University, Ordu, Turkey

fatihsay@odu.edu.tr

Abstract

Special functions, such as the gamma function, hypergeometric function, and Airy function, play a central role in the advancement of mathematical asymptotics [1,2,3,4] which has been an active research area in recent years. This talk examines the hyper-Airy differential equation [3], which is a fourth-order version of the Airy equation, in some limits. It addresses some of its asymptotic features that are not explicitly visible to classical asymptotic analysis.

Keywords: Asymptotic analysis; Divergent series; Small exponentials.

REFERENCES

1. Frank W. J. Olver, Asymptotics and Special Functions, Wellesley, MA: A K Peters, Ltd.; 1997.
2. Richard E. Meyer, A simple explanation of the Stokes phenomenon, SIAM Review, 31(3), 435–445, 1989.
3. Carl M. Bender, Steven A. Orszag, Advanced Mathematical Methods for Scientists and Engineers I, Springer-Verlag, New York, 1999.
4. George G. Stokes, On the discontinuity of arbitrary constants which appear in divergent developments. In Mathematical and Physical Papers, Cambridge University Press, 1864.

Total Dominator Chromatic Number of Lexicographic Product of a Cycle to a Graph

Canan Çiftçi[1], Adel P. Kazemi[2]

[1]Department of Mathematics, Faculty of Arts and Sciences, Ordu University, Ordu, Turkey

[2]Department of Mathematics, University of Mohaghegh Ardabili, Ardabil, Iran

cananciftci@odu.edu.tr, adelpkazemi@yahoo.com

Abstract

Coloring and domination are well-studied topics in graph theory. Motivated by the relation between coloring and total domination, the notion of total dominator coloring is studied by many authors [1,2,3]. A total dominator coloring of a graph G with no isolated vertex is a proper coloring of G in which each vertex of the graph is adjacent to every vertex of some (other) color class. The total dominator chromatic number of G is the minimum number of color classes in a total dominator coloring of it. This talk is about the total dominator coloring and examines the total dominator chromatic number of lexicographic product of a cycle to a graph.

Keywords: Total domination; Total dominator coloring; Lexicographic product.

Acknowledgement: This work is supported by TÜBİTAK 3501-Career Development Program under the project number 121F228.

REFERENCES

1. A. Vijayalekshmi, Total dominator colorings in graphs, International Journal of Advancements in Research and Technology, 4, 1-6, 2012.

2. A. P. Kazemi, Total dominator coloring in product graphs, Utilitas Mathematica, 94, 329-345, 2014.

3. M. A. Henning, Total dominator colorings and total domination in graphs, Graphs and Combinatorics, 31(4), 953-974, 2015.

BEZIER CURVES AND PATCHES WITH MODIFIED BERNSTEIN BASIS

Davut Canlı[1], **Süleyman Şenyurt**[1]

[1] Department of Mathematics, Ordu University, Ordu, Turkey

davutcanli@odu.edu.tr, senyurtsuleyman52@gmail.com

Abstract

In this paper, a new generalization of Bezier curves and surfaces are introduced by means of some modified Bernstein polynomial basis. The continuity conditions for the corresponding piecewise Bezier surfaces are discussed. An example is provided to illustrate the comparisons among the new and the classical Bernstein based Bezier objects.

Keywords: Bezier curves; Bezier Surfaces; G1, C1 continuity, Bernstein polynomials.

REFERENCES

1. Chen, X., Tan, J., Liu, Z.,Xie, J., Approximation of functions by a new family of generalized Bernstein operators, Journal of Mathematical Analysis and Applications, 450, 244–261, 2017.
2. Du, W. H., ve Schmitt, F. J., On the G1 continuity of piecewise Bézier surfaces: a review with new results. Computer-Aided Design, 22(9), 556-573, 1990.
3. FarinG, Curves and Surfaces for Comp. Aided Geom. Design, Academic Press, 1996.
4. Han, X. A., Ma, Y., ve Huang, X., A novel generalization of Bézier curve and surface. Journal of Computational and Applied Mathematics, 217(1), 180-193, 2008.
5. Hu, G., Bo, C., ve Qin, X., Continuity conditions for tensor product Q-Bézier surfaces of degree (m, n). Computational and Applied Mathematics, 37(4), 4237-4258, 2008.
6. Izgı, A., Approximation by a class of new type Bernstein polynomials of one and two variables, Global Journal of Pure and Applied Mathematics, 8, 55–71, 2012.
7. Phillips, G.M., Bernstein polynomials based on the q-integers, Annals of Numerical Mathematics, 4, 511–518, 1997.
8. Schurer, F., Linear Positive Operators in Approximation Theory, Math. Inst., Techn. Univ. Delf Report, Delft, 1962.
9. Stancu, D.D., Approximation of functions by a new class of linear polynomial operators, Rev. Roumaine Math. Pures Appl, 13, 1173–1194, 1968.

Revisiting Chatterjea Type Nonunique Fixed Point Theorems via Interpolation in QP_b - Space

Swapnil Verma[1]

[1] Department of Mathematics, University of Delhi, India

swapniliitdelhi@gmail.com

Abstract

In the early years of the 20th century, the renowned mathematician Banach [1] commenced the concept of the Banach Contraction Principle. Due to its consequences and feasible implementations, the idea has been enlarged and generalized in various directions [2]-[6]. Recently, Karapinar [7] adopted an interpolative approach to establish fixed point results in the setting of complete metric space. In 2016, Gupta and Gautam [8]-[9] defined a new metric space known as quasi- partial b -metric space and proved fixed point theorems on this space. Inspired by these results, we have redefined the Chatterjea type contraction in the framework of quasi-partial b-metric space and proved the corresponding common fixed point theorem by adopting the notion of interpolation. The results are further validated with the application based on them. The applications of interpolative contraction in sensitivity analysis of experimental signals and synthesis of scientific data where approximation of natural curves and surfaces is needed, are illustrated here. Examples are given which are based on the new approach.

Keywords: Chatterjea type contraction; quasi-partial b-metric space; interpolative contraction; fixed point

1.Banach, S.: Sur les operations dans les ensembles abstraits et leur application aux equations integrales. Fundam. Math. 3, 133–181 (1922).

2. Bakhtin, I.A.: The contraction mapping principle in almost metric spaces. Funct. Anal. 30, 26–37 (1989).

3. Czerwik, S.: Contraction mappings in *b*-metric spaces. Acta Math. Inform. Univ. Ostrav. 1, 5–11 (1993).

4. Kannan, R.: Some results on fixed points. Bull. Calcutta Math. Soc. 60, 71–76 (1968).

5. Rus, I.A.: Picard operators and applications. Sci. Math. Jpn. 58(1), 191–219 (2003).

6.Matthews, S.G.: Partial metric topology. Ann. N. Y. Acad. Sci. 1994, 728, 183–197.

7. Karapinar, E.: Revisiting the Kannan type contractions via interpolation. Adv. Theory Nonlinear Anal. Appl. 2018, 2, 85–87.

8.Gupta, A.; Gautam, P.: Topological structure of quasi-partial b-metric spaces. Int. J. Pure Math. Sci. 2016, 17, 8–18.

A NOVEL NUMERICAL APPROXIMATION FOR SINGULARLY PERTURBED VOLTERRA INTEGRO-DIFFERENTIAL EQUATION WITH INTEGRAL BOUNDARY CONDITION

Zelal TEMEL[1] and **Musa CAKIR**

[1] Department of Mathematics, University of Van Yuzuncu Yil, Van, Turkey

Department of Mathematics, University of Van Yuzuncu Yil, Van, Turkey

zelaltemel65@gmail.com,

cakirmusa@hotmail.com.

Abstract

In this paper, the initial value problem for the singularly perturbed Volterra integro-differential equation with integral boundary condition was considered. Our purpose is to construct and analyze a numerical approach with uniform convergence according to ε. The numerical solution of the considered problem is discretized on a uniform mesh using implicit difference rules for the differential part and the composite right-side rectangle rule for the integral part. It is shown that the method displays first-order uniform convergence in the perturbation parameter. Also, numerical experiments are studied to show the performance and viability of the proposed approach.

Keywords: Singularly perturbed problem; Numerical solution; Boundary layer; Uniform convergence.

REFERENCES

1. Gabil M.Amiraliyev, Ömer Yapman, Mustafa Kudu, A fitted approximate method for a Volterra delay-integro-differential equation with initial layer. Hacet. J. Math. Stat., 48, 5 1417-29, 2019.
2. Ömer Yapman, Gabil M.Amiraliyev, Convergence analysis of the homogeneous second order difference method for a singularly perturbed Volterra delay-integro-differential equation, Chaos, Solitons and Fractal, 150, 111100, 2021.
3. Musa Cakir, A Numerical Study on the Difference Solution of Singularly Perturbed Semilinear Problem with Integral Boundary Condition, Mathematical Modelling and Analysis, 21, 5, 644-658, 2016.

CONVERGENCE ANALYSIS OF FITTED NUMERICAL METHOD FOR A SINGULARLY PERTURBED NONLINEAR VOLTERRA INTEGRO-DIFFERENTIAL EQUATION WITH INTEGRAL BOUNDARY CONDITION

Zelal TEMEL[1] and **Musa CAKIR**

[1] Department of Mathematics, University of Van Yuzuncu Yil, Van, Turkey

Department of Mathematics, University of Van Yuzuncu Yil, Van, Turkey

zelaltemel65@gmail.com,

cakirmusa@hotmail.com.

Abstract

This work is concerned with a singularly perturbed nonlinear Volterra integro-differential equation with integral boundary condition. Firstly, bounds on the solution and its derivative of the solution of this problem are derived. To solve it numerically, we use a fitted difference scheme on a piecewise mesh (Shishkin type mesh), which is discretized using quadrature rules for the differential part and the composite numerical quadrature rules for both integral part in the equation and integral boundary condition. Then, the stability and convergence analysis of the numerical approach are discussed and analyzed. Furthermore, numerical results supporting the theoretical results are presented.

Keywords: Volterra integro-differential equations; Singularly perturbation; Error estimate; Shishkin mesh.

REFERENCES

1. Ömer Yapman, Gabil M.Amiraliyev, Ilhame Amirali, Convergence analysis of fitted numerical method for a singularly perturbed nonlinear Volterra integro-differential equation with delay, 355, 301-309, 2019.
2. Musa Cakir, Gabil M.Amiraliyev, A finite difference method for the singularly perturbed problem with nonlocal boundary condition, Appl. Math. and Comput, 160, 2, 539-549, 2005.
3. Musa Cakir, A Numerical Study on the Difference Solution of Singularly Perturbed Semilinear Problem with Integral Boundary Condition, Mathematical Modelling and Analysis, 21(5), 644-658, 2016.

EVALUATION OF TEACHER CANDIDATES' BIOTECHNOLOGY KNOWLEDGE LEVELS AND ATTITUDES TOWARDS BIOTECHNOLOGY

Berat GEVŞEK, Assit. Prof. Dr. Elif ÇİL
beratgevsek@odu.edu.tr, elifcil@odu.edu.tr

Today, science and technology are developing rapidly, and the importance of science is increasing in almost every part of society. With these technological developments, the relationship between science and society gains importance. Within the general objectives of science education, besides teaching the scientific concepts underlying the developments in science and technology, individuals follow these developments. It is necessary to ensure that they are scientifically literate. When the literature is examined, it is seen that pre-service science teachers and science teachers need more information in areas such as genetic engineering, cloning, and genetically modified organisms.

This study aims to determine the knowledge level of pre-service science teachers about biotechnology and evaluate their attitudes towards biotechnological developments. The study was modeled according to the descriptive survey model. In the research, the biotechnology knowledge scale with 22 questions developed by Yüce in 2011 and the biotechnology attitude scale with 46 questions developed by Yazıcı in 2009 was used to determine the knowledge level of pre-service science teachers on biotechnology subjects. These scales were applied to 60 science teacher candidates studying at a state university in North Anatolia in the 2021-2022 academic year. The knowledge of pre-service science teachers about biotechnology was investigated according to different variables (gender, class of education, whether or not they had taken a Biotechnology course). The obtained data were analyzed using the SPSS 26 package program.

When the research results are evaluated, it is seen that the biotechnology knowledge levels of the science teacher candidates; It has been determined that there is a statistically significant difference according to the level of education and the status of taking the course or not. This statistical difference favored the 2nd year students who took the course. In addition, a significant ($p < 0.05$) difference was found in the biotechnology attitude levels of science teacher candidates according to their gender. This difference was in favor of female teacher candidates. However, no statistically significant difference was found between knowledge levels and gender. A statistically significant difference was not determined according to the biotechnology attitude levels of teacher candidates, their grade level, and whether they took the course or not.

Keywords: quantitative analysis, pre-service science teacher, attitude

REFERENCES

Yüce, Z. (2011) *"Fen Bilgisi Öğretmenliği Öğrencilerinin Biyoteknoloji Konusundaki Bilgileri ve Biyoteknoloji Uygulamalarına Yönelik Biyoetik Yaklaşımları: Tutum, Görüş ve Değer Yargıları"*, Gazi Üniversitesi Eğitim Bilimleri Enstitüsü, Doktora Tezi, Ankara.

Yazıcı, N. N. (2009). "*Bilim Kurgu ile Biyoetik Grup Tartışmalarının Biyoteknolojiye Yönelik Tutumlar ve Akademik Başarı Üzerine Etkileri"*, Muğla Üniversitesi Fen Bilimleri Enstitüsü, Yüksek Lisans Tezi, Muğla.

Arslankara, V. (2019). *Fen bilgisi eğitiminde biyoteknolojinin önemine yönelik fen bilgisi öğretmen adaylarının görüşleri: Konya örneği* (Doctoral dissertation, Necmettin Erbakan University (Turkey)).Soğukpınar, R., & Karışan, D. (2020). Fen Bilgisi Öğretmen Adaylarının Genetik Okuryazarlık Konularına Ve Biyoteknolojiye Yönelik Tutumlarının İncelenmesi. *Uludağ Üniversitesi Eğitim Fakültesi Dergisi*, *33*(2), 410-445.

A neutrosophic correlation coefficient and its application in recognition of patterns

Rıdvan Şahin [1]

[1] Department of Mathematics Engineering, University of Gumushane, Gumushane, Turkey

mat.ridone@gmail.com,

Abstract

In probability and statistical theory, the correlation coefficient indicates the strength of the linear correlation between two random variables. In this study, a new correlation coefficient is defined between neutrosophic sets (NS). This new approach is defined in the range [-1, 1], similar to classical statistics, and it shows the strength of the relationship between the considered NSs. Finally, to show the usefulness of the proposed new measure, a pattern recognition problem is resolved with the help of the defined measure.

Keywords: Neutrosophic set; Correlation coefficient; Decision making; Pattern recognition

REFERENCES

1. F. Smarandache, Neutrosophic set: A generalization of the intuitionistic fuzzy set, International Journal of Pure and Applied Mathematics, Vol: 24, 287–297, 2005.
2. JS Chai, et al., New similarity measures for single-valued neutrosophic sets with applications in pattern recognition and medical diagnosis problems, Complex & Intelligent Systems Vol:7, No:2, 703-723, 2021.
3. H. Garg, Some new biparametric distance measures on single-valued neutrosophic sets with applications to pattern recognition and medical diagnosis, Information, Vol: 8, No:4, 162, 2017.
4. J. Ye, Multicriteria decision-making method using the correlation coefficient under single-valued neutrosophic environment, International Journal of General Systems, Vol:42, No:4, 386-394, 2013.
5. R. Şahin, M. Karabacak, A novel similarity measure for single-valued neutrosophic sets and their applications in medical diagnosis, taxonomy, and clustering analysis, In Optimization theory based on neutrosophic and plithogenic sets (pp. 315-341). Academic Press. 2020.

New distance measure for neutrosophic sets and its application in medical diagnosis

Rıdvan Şahin [1]

[1] Department of Mathematics Engineering, University of Gumushane, Gumushane, Turkey

mat.ridone@gmail.com,

Abstract

As a new extended form of intuitionistic fuzzy sets, neutrosophic sets are powerful tools for describing uncertainty and uncertainty in complex problems. In this study, we describe a new measure of distance for neutrosophic sets based on triangular divergence. Next, we prove that the proposed distance measurement methods meet the axiomatic requirements of the distance function. We also use proposed distance measure method to solve a medical diagnostic problem in the neutrosophic universe.

Keywords: Neutrosophic set; Distance measure; Medical diagnostic, Decision making

REFERENCES

1. F. Smarandache, Neutrosophic set: A generalization of the intuitionistic fuzzy set, International Journal of Pure and Applied Mathematics, Vol: 24, 287–297, 2005.
2. J. Ye, Single-valued neutrosophic minimum spanning tree and its clustering method, Journal of intelligent Systems, Vol:23, No: 3, 311-324, 2014.
3. R. Şahin, M. Karabacak, A novel similarity measure for single-valued neutrosophic sets and their applications in medical diagnosis, taxonomy, and clustering analysis, In Optimization theory based on neutrosophic and plithogenic sets (pp. 315-341). Academic Press. 2020.
4. H. L. Huang, New distance measure of single-valued neutrosophic sets and its application, International Journal of Intelligent Systems, Vol:31 No:10, 1021-1032, 2016.
5. J. Ye, Improved cosine similarity measures of simplified neutrosophic sets for medical diagnoses, Artificial intelligence in medicine, Vol: 63, No:3, 171-179, 2015.

A STUDY ON SOME SEQUENCE SETS BASED ON LUCAS BAND MATRIX AND MODULUS FUNCTIONS

Mustafa I. Hatim, Çiğdem A. Bektaş

Department of Mathematics, University of Firat, Elazig, Turkey

mustafa.ih88@gmail.com cbektas@firat.edu.tr

Abstract

In this paper, we introduce the sequence sets $\ell_p\left(F^k, u, \hat{E}(r,s)\right)$ and $\ell_\infty\left(F^k, u, \hat{E}(r,s)\right)$ which are depended on Lucas band matrix and modulus functions. After that we extend our study and generalize the sequence sets $\ell_p\left(F^k, u, N, \hat{E}(r,s)\right)$ and $\ell_\infty\left(F^k, u, N, \hat{E}(r,s)\right)$ over a normed linear space (S, N). Furthermore, we study on some properties of these sequence sets and obtain some inclusion relations related to them.

Keywords: Lucas numbers, Modulus function, Sequence space, BK-space.

REFERENCES

1. Gülcan Atici, Çiğdem Bektaş, On some new generalized difference sequence spaces defined by a sequence of moduli, Mathematica Slovaca 61.5, 789-798, 2011.
2. Sayd Abdul Mohiuddine, Kuldip Raj, Anu Choudhary, Difference sequence spaces based on Lucas band matrix and modulus function, São Paulo Journal of Mathematical Sciences, 1-12, 2021.
3. M. Karakas, A. M. Karakas, A study on Lucas difference sequence spaces $\ell_p(\hat{G}(r,s))$ and $\ell_\infty(\hat{G}(r,s))$, Maejo Int. J. Sci. Technol. 12, 70–78, 2018.
4. Serpil Pehlivan, Brian Fisher, On some sequence spaces, Indian Journal of Pure and Applied Mathematics 25, 1067-1072, 1994.
5. Hidegorô Nakano, Modular sequence spaces, Proc. Japan Acad 27, 508-512, 1953.
6. Ivor J. Maddox, Sequence spaces defined by a modulus, Math. Proc. Camb. Phil. Soc. 100, 161–166, 1986.
7. Yavuz Altın, Hifsi Altinok, Rifat Çolak, On some semi normed sequence spaces defined by a modulus function, Kragujevac J. Math. 29, 121-132, 2006.
8. Mohammad Mursaleen, Abdullah Kaid Noman, On some new sequence spaces of non-absolute type related to the spaces ℓ_p and ℓ_∞ II, Math. Commun. 16, 383-398, 2011.

THE EFFECT OF MIND AND INTELLIGENCE GAMES COURSE ON STUDENTS' MATHEMATICS ACHIEVEMENT: AN EXPERIMENTAL STUDY

Elif ŞENKAL [1]

elifcvcn@gmail.com,

Hayal YAVUZ MUMCU[2]

hayalym52@gmail.com,

[1,2]Ordu University, Ordu, Turkey

Abstract

In this study, it is aimed to examine the effect of mind and intelligence games on the academic achievement of students in mathematics. Sudoku from reasoning and operation games, reversi, go and mangala games from strategy games were determined. Experimental study method, one of the quantitative research methods, was used in the study. The research was carried out in a public school in Fatsa district of ORDU province in the 2021-2022 academic year. The study group consists of 30 students studying in the 7th grade. Students were selected by convenient sampling method, which is not a random sampling method. The students in the experimental group received training on mind and intelligence games for two hours a week during the 8-week period. "Mathematics course academic achievement test" prepared by the researcher was used as a data collection tool. For the validity of the test, the opinions of two mathematics educators who are experts in the field were taken, and item and reliability analyzes were made for its reliability. In this study, the games were first explained and then the students were allowed to play. Accordingly, the teaching process lasted 8 weeks. As a result of the research, it was seen that the academic success of the students who received mind and intelligence games training increased.

Keywords: Mind and intelligence games course; 7th grade students, Experimental study

A New Reproducing Kernel Approach for Nonlinear Variable Order Fractional Volterra Integro-Differential Equations

Mehmet Giyas SAKAR [1]

[1] Department of Mathematics, Faculty of Science, Van Yüzüncü Yıl University, Van, Turkey

giyassakar@hotmail.com

Abstract

In this article, a new reproducing kernel approach is developed for nonlinear variable order fractional Volterra integro-differential equations. The fractional derivatives are taken in the Caputo sense. This approach is based on reproducing kernel which is constructed by shifted Legendre polynomials. In order to shows the robustness of the proposed method, some examples are solved and numerical results are given in tabulated forms.

Keywords: Reproducing kernel method; Legendre polynomials; Caputo derivative; Variable order; Volterra integro-differential equations.

REFERENCES

1. J. Yang, H. Yao, B. Wu, An efficient numerical method for variable order fractional functional differential equation, Applied Mathematics Letters, 76, 221–226 (2018).

2. M. G. Sakar, Iterative reproducing kernel Hilbert spaces method for Riccati differential equations, Journal of Computational and Applied Mathematics, 309, 163–174 (2017).

3. X. Li, H. Li, B. Wu, A new numerical method for variable order fractional functional differential equations, Applied Mathematics Letters, 68, 80–86 (2017).

4. M. Gui, Y. Lin. *Nonlinear Numercial Analysis in the Reproducing Kernel Space*. Nova Science Publisher: New York, 2008.

EXAMINATION OF 8th GRADE STUDENTS' OPINIONS ON MODEL ELICITING ACTIVITIES

Büşra UYGUN, Meral CANSIZ AKTAŞ

Department of Mathematics Education, University of Ordu, Turkey

busrauygun96@gmail.com, cansizmeral@hotmail.com

Abstract

Many students see mathematics as a discipline separated from everyday life. In order to break this prejudice toward mathematics in students, it is very important to expose students to daily life problems that can be solved by using mathematical skills (Lesh & Zawojewsky, 2007). For this purpose, we suggested that using model eliciting activities may increase students' mathematical skills and encourage them to use mathematics in daily life.

Model eliciting activities are non-routine problem situations that can be encountered in everyday life. These problems may have different possible solutions; hence, the goal is not to conclude with a single correct answer. The purpose of these activities is to interpret a problem by using mathematics and to offer a perspective for people who may encounter the same situations (Lesh & Zawojewsky, 2007).

The aim of this study is to examine the opinions of 8th-grade students about model eliciting activities. In the scope of the research, we have studied with 10 students studying in a public school in Van, Turkey. This is a qualitative study that examines students' opinions. In the study, educational modeling activities created by Purdue University have used as a data collection tool. The students have divided into two groups and, the activities have carried out as teamwork. At the end of the activities, semi-structured interview questions have used to get individual opinions of the students and the collected data have analyzed with the descriptive analysis method. We observed that students have found model eliciting activities difficult and complex as they got used to routine problems. However, trying to create solutions via mathematics for daily life problems helped students to understand the importance of mathematics. We concluded that the model eliciting activities are important and helpful for easier learning of mathematical concepts and it can distract the lessons from monotony.

Keywords: Model eliciting activities; Teamwork

REFERENCES

1. Bukova Güzel, E., Tekin Dede, A., Hıdıroğlu, Ç.N., Kula Ünver, S. ve Özaltun Çelik, A. (2019).Matematiksel modellemeye giriş. In E. Bukova Güzel (Eds.), *Matematik eğitiminde matematiksel modelleme* (pp.1-160).Ankara: Anı Yayıncılık.,
2. Hıdıroğlu, Ç.N. ve Özkan Hıdıroğlu, Y. (2017). Altıncı sınıf öğrencilerinin matematiksel modellemede oluşturdukları gerçek yaşam problem durumu modelleri. İlköğretim Online, 16(4), 1702-1731.
3. Lesh, R.A., & Zawojewski, J. (2007). Problem solving and modeling. In F. Lester (Ed.), *Second handbook of research on mathematics teaching and learning: A Project of the national council of teachera of mathematics teacging and learning: A Project of the national council of teachers of mathematics. Charlotte, NC: Information Age Publishing.*

Numerical solution of fuzzy differential equations by twostep modified Simpson rule Interpolation of Fuzzy Number

Ekhtiar Khodadadi[1], Mesut Karabacak[2], Ercan Çelik[3]

[1] Department of Mathematics, Malekan Branch, Islamic Azad University, Malekan, Iran.

[2] Department of Mathematics, Faculty of Science, Atatürk University, Erzurum, Turkey.
[3] Kyrgyz-Turkish Manas University, Department of Applied Mathematics and Informatics, Bishkek-Kyrgyzstan

khodadadi@atauni.edu.tr , mkarabacak@atauni.edu.tr, ercan.celik@manas.edu.kg

Abstract

In this paper, a numerical explicit two-step modified Simpson rule for fuzzy first-order initial value problem is present, and convergence and stability of the mentioned rule are proved, and their applicability is illustrated with some examples.

Keywords: Fuzzy differential equations, Fuzzy Cauchy problem, Two-step methods, Midpoint rule, Trapezoidal rule, Modified Simpson rule

REFERENCES

[1] A. Khastan and K. Ivaz, "Numerical solution of fuzzy differential equations by Nyström method," *Chaos, Solitons & Fractals*, vol. 41, no. 2, pp. 859–868, 2009.

[2] K. Ivaz, A. Khastan, and Juan J. Nieto, A Numerical Method for Fuzzy Differential Equations and Hybrid Fuzzy Differential Equations, Hindawi Publishing Corporation *Abstract and Applied Analysis*, Volume 2013, Article ID 735128.

[3] J. J. Nieto, A. Khastan, and K. Ivaz, "Numerical solution of fuzzy differential equations under generalized differentiability," *Nonlinear Analysis: Hybrid Systems*, vol. 3, no. 4, pp. 700–707, 2009.

[4] M. Friedman, M. Ma, and A. Kandel, "Numerical solutions of fuzzy differential and integral equations," *Fuzzy Sets and Systems*, vol. 106, no. 1, pp. 35–48, 1999,

[5] T. Allahviranloo, N. Ahmady, and E. Ahmady, "Numerical solution of fuzzy differential equations by predictor-corrector method," *Information Sciences*, vol. 177, no. 7, pp. 1633–1647, 2007.

A NUMERICAL APPROACH FOR SINGULAR BOUNDARY VALUE PROBLEM

Onur Saldır,[1] **Mehmet Giyas Sakar**[2]

[1,2] Department of Mathematics, Faculty of Science, Van Yüzüncü Yıl University, Van, Turkey

onursaldir@gmail.com

Abstract

In this study, a numerical method is proposed to obtain an approximate solution for singular boundary value problems. The presented method is based on reproducing kernel method and Legendre polynomials. Comparison of numerical results with other methods is given by tables and graphics.

Keywords: Reproducing kernel method; Sindular boundary value problem; Legendre polynomials.

REFERENCES

1. Geng F, Cui M. A novel method for nonlinear two-point boundary value problems: combination of ADM and RKM. Applied Mathematics and Computation. 2011;217(9):4676–4681.

2. Zhou F, Xu X. Numerical solutions for the linear and nonlinear singular boundary value problems using Laguerre wavelets. Advances in Difference Equations. 2011;17(2016):1–15.

3. Rashidinia J, Nabati M, Ahanj S. Sinc-Galerkin method for solving a class of nonlinear two-point boundary value problems. Bulletin of the Iranian Mathematical Society. 2015;41(2):333–352.

.

A NEW APPROACH FOR NUMERICAL SOLUTION OF FOURTH ORDER EMDEN-FOWLER EQUATION

Onur Saldır[1]

[1] Department of Mathematics, Faculty of Science, Van Yüzüncü Yıl University, Van, Turkey

onursaldir@gmail.com

Abstract

In this study, a new numerical approach will be proposed for fourth order Emden-Fowler equation. This approach is based on the reproducing kernel method with Legendre polynomials. To observe the effect of the approach numerical results will be given tables and graphics.

Keywords: Reproducing kernel method; Legendre polynomials; Fourth order Emden-Fowler equation.

REFERENCES

1. R. H. Fowler, The form near infinity of real, continuous solutions of a certain differential equation of the second order, Applied Mathematics and Computation, Quart. J. Math., 1914.
2. S. A. Khuri, An alternative solution algorithm for the nonlinear generalized Emden-Fowler equation, International Journal of Nonlinear Sciences and Numerical Simulation. 2, 299-302, 2001.
3. S. A. Khuri, A. Sayfy, Numerical solution for the nonlinear Emden-Fowler type equations by a fourth-order adaptive method, Int. J. Comput. Methods. 11, 1-21, 2014.

A FINITE DIFFERENCE METHOD FOR FREDHOLM INTEGRO-DIFFERENTIAL EQUATIONS

Fırat Çakır [1], Hayriye Güçkır Çakır[2], Musa Çakır[3]

[1] Department of Mathematics, University of Batman, Batman, Turkey

firat.cakir@batman.edu.tr,

[2]Department of Mathematics, Adiyaman University, Adiyaman, Turkey

hcakir@adiyaman.edu.tr

[3]Department of Mathematics, Van Yuzuncu Yil University,Van,Turker

cakirmusa@hotmail.com

Abstract

In this work, we study the numerical solution of a second-order linear Fredholm integro-differential equation (FIDE) by a finite difference method. The discretization of the problem is obtained by a finite difference method on a uniform mesh. We construct the method using the integral identity method with basis functions and dealing with the integral terms by interpolating quadrature rules with remainder terms. We further employ the factorization method to establish the algorithm. We demonstrate the error estimates and the convergence of the method. The numerical results are demonstrated to verify the order of accuracy.

Keywords: FIDE; Difference Scheme; Error estimates.

REFERENCES

1. E. Cimen and M. Cakir. A uniform numerical method for solving singularly perturbed fredholm integro-differential problem. Comp. Appl. Math., 40:42, 2021.
2. M. Cakir, B. Gunes, and H. Duru. A novel computational method for solving nonlinear volterra integro-differential equation. Kuwait J. Sci, 48(1):1–9, 2021.
3. O. Yapman, M. Amiraliyev G., and I. Amirali. Convergence analysis of fitted numerical method for a singularly perturbed nonlinear volterra integro-differential equation with delay. J. Comput. Appl. Math., 355:301–309, 2019.

Spectra of a linear pencil of pentadiagonal matrices

Hasen Mekki ÖZTÜRK

Department of Mathematics, Ordu University, Ordu, Turkey

hasenozturk@odu.edu.tr

Abstract

Let $c \in \mathbb{R}$, and define the pentadiagonal matrix A and the diagonal matrix B by

$$A = \begin{pmatrix} c & 1 & 1 & & & \\ 1 & c & 1 & \ddots & & \\ 1 & 1 & c & \ddots & \ddots & \\ & \ddots & \ddots & \ddots & \ddots & 1 \\ & & \ddots & \ddots & \ddots & 1 \\ & & & 1 & 1 & c \end{pmatrix}, \qquad B = \begin{pmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & 1 & & & \\ & & & -1 & & \\ & & & & \ddots & \\ & & & & & -1 \end{pmatrix},$$

respectively, where the size of both matrices is $(2n)\times(2n)$, $n \in \mathbb{N}$, and the diagonal matrix B has n plus ones and n minus ones. In this research, we conduct numerical experiments to investigate the behaviour of the eigenvalues of a sign-indefinite linear pencil $P(\lambda) = A - \lambda B$. Motivated by our numerics, we pose some conjectures concerning the localization of the spectrum of $P(\lambda)$.

Keywords: Linear matrix pencils; Non-self-adjoint matrices; Complex eigenvalues; Spectral theory.

REFERENCES

1. E. B. Davies and M. Levitin, Spectra of a class of non-self-adjoint matrices, Linear Algebra and its Applications, Vol:448, 55–84, 2014.
2. A. S. Markus, Introduction to the Spectral Theory of Polynomial Operator Pencils, Amer. Math. Soc., Providence, RI, 1988.

EXAMİNİNG OF MİDDLE SCHOOL 7TH GRADE STUDENTS' PROBLEM POSİNG STATUS ABOUT 'EQUALİTY AND EQUATİON'.

Gizem Nur Battal[1]

Assis. Prof Meral Cansız Aktaş[2]

[1] Master Student, University of Ordu, Ordu, Turkey, gizemnurbattal@gmail.com

[2] Faculty of Education, University of Ordu, Ordu, Turkey, meralcansizaktas@odu.edu.tr

The aim of the research is to examine the problem posing status of middle school 7th grade students about 'equality and equation'. The participants of the research consist of 24 students, 17 girls and 7 boys. The Problem Posing test was used as a data collection tool. The problem posing test was developed by researchers and includes 10 problem posing activities. Problem posing activities were arranged according to Stoyanova and Ellerton's (1996) problem posing classification. The analysis of the data obtained was classified according to the problem posing classification scheme of Türnüklü et al. (2017). According to the results obtained from the research, some suggestions were presented.

Keywords: Problem posing, equality and equation

REFERENCES

Stoyanova, E., & Ellerton, N. F. (1996). A framework for research into students' problem posing in school mathematics. In P. Clarkson (Ed.). Technology in Mathematics Education (pp. 518–525). Mathematics Education Research Group of Australasia.

Türnüklü, E., Ergin, A. S. ve Aydoğdu, M.Z. (2017). 8. sınıf öğrencilerinin üçgenler konusunda problem kurma çalışmalarının incelenmesi. Bayburt Eğitim Fakültesi Dergisi, 12(24), 467-486

Predictor-Corrector $(m+1)$ −step Method for Solving Fuzzy Ordinary Differential Equations by Using the Interpolation of Fuzzy Number

Ekhtiar Khodadadi[1], Mesut Karabacak[2], Ercan Çelik[3]

[1] Department of Mathematics, Malekan Branch, Islamic Azad University, Malekan, Iran.

[2] Department of Mathematics, Faculty of Science, Atatürk University, Erzurum, Turkey.

[3] Kyrgyz-Turkish Manas University, Department of Applied Mathematics and Informatics, Bishkek-Kyrgyzstan

khodadadi@atauni.edu.tr , mkarabacak@atauni.edu.tr, ercan.celik@manas.edu.kg

Abstract

In this paper, an Adams Predictor-Corrector method $(m+1)$ −step is used to solve first-order fuzzy linear ordinary differential equations. The Predictor-Corrector method is generated by combining an explicit $(m+1)$ −step method. The convergence of these methods is proven in detail. Finally, these methods are illustrated using examples of fuzzy initial value problems.

Keywords: Predictor-Corrector method, Fuzzy number, Fuzzy differential equation, Adams-Bashforth method, Hukuhara differentiable, Adams-Moulton method

REFERENCES

[1] El Naschie, M.S., A review of E-infinite theory and the mass spectrum of high-energy particle physics. Chaos, Solitons and Fractals 19 (2004), 209-236.

[2] Zhang, H., Liao, X. and Yu, J., Fuzzy modelling and synchronization of hyperchaotic systems.Chaos, Solitons and Fractals 26 (2005), 835-843.

[3] Feng, G. and Chen, G., Adaptive control of discrete-time chaotic systems: a fuzzy control approach. Chaos, Solitons and Fractals 23 (2005), 459-467.

[4] Tanaka,Y., Mizuna Y. and Kado, T., Chaotic dynamics in the Friedman equation. Chaos, Solitons and Fractals 24 (2005), 407-422. 24

[5] Abbad, M.F., Von Keyserlingk, D.G., Linkens, D.A. and Mahfouf, M., Survey of utilisation of fuzzy technology in medicine and healthcare. Fuzzy Sets and Systems 120 (2001), 331-349.

[6] D.O'Regan, V. Lakshmikantham, J.J. Nieto, Initial and boundary value problems for fuzzy differential equations. Nonlinear Analysis 54 (2003), 405-415

On the stability and stabilization for the implicit differential control system in Banach spaces

Mohamed Hariri [1] **, Zohra Bouteffal** [2] **,**
Nourelhouda Beghersa [3] **and Mehdi Benabdallah** [4]

[1]Department of Mathematics, University of Belhad Bouchaib , Ain Temouchent, Algeria

[2] Department of Mathematics, University of Mustapha Stambouli, Mascara, Algeria.
[3] Department of Mathematics, University of Science and Technology,Oran, Algeria.

[4] Department of Mathematics, University of Science and Technology, Oran, Algeria.

mohamed.hariri@univ-temouchent.edu.dz

Abstract

The aim of this research is to generalize the famous General Theorem of Liapounov of the stability for the explicit systems to the form:

$$Ax'(t) = Bx(t), \quad t \geq 0,$$

using the spectral theory of the corresponding pencil of operators $\lambda A - B$. The achieved results can be applied to stabilizing the implicit control system

$$Ax'(t) = Bx(t) + Cu(t), \ t \geq 0, \ x \in X.$$

Where A, B and C are linear closed operators in X and the control space U are Banach spaces, and the function u is square integrable in the sense of Bochner. The operator A is not necessarily invertible.

Keywords: Spectral theory, stability and stabilization theory, pencil of operators, implicit systems.

REFERENCES

1. J.L. Daleckiĭ and M.G. Krein, Stability of Solutions of Differential Equations in BanachSpace, Amer. Math. Soci. Providence, RI, 1975.
2. M. Benabdallah, A.G. Rutkas and A.A. Soloviev, On the stability of degenerate difference systems in Banach spaces, J. Sov. Math. , No 57 3435-3439, 1991.

3.S.L. Gefter, A.L. Piven, Implicit linear non-homogeneous difference equation in Banach and locally convex spaces, J.Math physics, analysis, geometry, vol 15, No 3, 336-353, 2019.

4. A.G. Rutkas, Spectral methods for studying degenerate differential-operator equations, J.Math. Sci. 144 ,4246–4263, 2007.

SOME NEW EXISTENCE RESULTS FOR FRACTIONAL DIFFERENTIAL INCLUSIONS WITH ANTI-PERIODIC BOUNDARY CONDITIONS

Münevver Tuz[1]

[1] Department of Mathematics, University of Firat, Elazig, Turkey

mtuz@firat.edu.tr,

Abstract

In this paper, the existence of anti-periodic solution for competitive neural networks with delays neural networks under the same framework. Firstly, the existence of anti-periodic solution is discussed by using the Lyapunov functional method.Then some sufficient conditions are obtained to guarantee of anti-periodic solution for such neural networks. The obtained results are new and improve some earlier publications. Finally, numerical example are given to illustrate the effectiveness of the theoretical results.

Keywords: Neural networks; Existence; Linear differential equation; Boundary conditions , Anti-periodic.

REFERENCES

1. Kaslik, E., Sivasundaram, S., Dynamics of Fractional-Order Neural Networks, Proceedings, International Joint Conference on Neural Network (IJCNN -11), San Jose, Cal., USA, 611-618, 2011.

2.Kaslik, E., Sivasundaram, S., Non-Linear Dynamics and Chaos in Fractional-Order Neural Networks, Neural Networks, 32, 245-256, 2012.

3.Wu, R. C., et al., Finite-Time Stability of Fractional-Order Neural Networks with Delay, Communications in Theoretical Physics, 60, 2, 189-193, 2013.

4.Diethelm, K., The Analysis of Fractional Differential Equations, Springer, Berlin, Germany, 2010.

5.Kilbas, A. A., et al., Theory and Applications of Fractional Differential Equations, North-Holland Mathematics Studies, Elsevier Science, Amsterdam, The Netherlands, 2006.

INVESTIGATION OF SOLITARY WAVE SOLUTIONS FOR THE NONLINEAR (2 + 1) -DIMENSIONAL TYPICAL REFRACTION EQUATION

Münevver Tuz[1], **Asıf Yokuş**[2]

[1,2] Department of Mathematics, University of Firat, Elazig, Turkey

[1]mtuz@firat.edu.tr,

[2]ayokus@firat.edu.tr,

Abstract

In this paper, the nonlinear (2 + 1) -dimensional Typical Refraction equation is discussed and new complex hyperbolic propagating wave solutions are obtained for this equation by using the $\left(\frac{G'}{G},\frac{1}{G}\right)$ extension method. In the solutions obtained, graphs were drawn by giving special values to the parameters. The method used is ideal for achieving the goals set for this study, and the solutions obtained are different from the others and shed light on new studies..

Keywords: Extension method; Typical Refraction equation; Complex hyperbolic; wave solutions.

REFERENCES

1. Liu, S., Fu, Z., Liu, S., and Zhao, Q., Jacobi elliptic function expansion method and periodic wave solutions of nonlinear wave equations, Physics Letters A, 289 (1-2), 69-74, 2001.

2. Liu, W., and Chen, K., The functional variable method for finding exact solutions of some nonlinear time-fractional differential equations, Pramana, 81 (3), 377-384, 2013.

3. Yokuş, A., Comparison of caputo and conformable derivatives for time-fractional Korteweg–de Vries equation via the finite difference method, International Journal of Modern Physics B, 32 (29), 1850365, 2018.

4. Yokus, A., and Kaya, D., Numerical and exact solutions for time fractional Burgers'equation, Journal of Nonlinear Sciences and Applications, 3419-3428, 10, 2017.

5. Yokuş, A., An expansion method for finding traveling wave solutions to nonlinear pdes, İstanbul Ticaret Üniversitesi, 2015.

Evolutionary Computation for Multi-Constrained Path Selection Algorithm on the Internet

Habiba Akter[1]

Department of Engineering and Design,

School of Engineering and Informatics,

University of Sussex,

United Kingdom

h.akter@sussex.ac.uk

Abstract

This work investigates the implementation of an Evolutionary Algorithm (EA) that helps in building a path computation tool while benefiting the end-users through Loose-Source Routing. This is done in a scenario where internet tunnels are present in a part of the network topology. The paths calculated may or may not have tunnels present in them. The objective functions for optimisation will be designed based on the cost values associated with the end-to-end path. The approach avoids the need for inter-operator cooperation, although such cooperation would provide a means of extending tunnels across AS peers.

Keywords: Evolutionary Algorithm; Path Calculation; Internet Topology.

REFERENCES

1. Habia Akter and Chris Phillips. 2019. Tunnelling the Internet. Journal of Telecommunications and the Digital Economy 7, 1 (2019).
2. Zhendong Liu, Yawei Kong, and Bin Su. 2016. An improved genetic algorithm based on the shortest path problem. In 2016 IEEE international conference on information and automation (ICIA). IEEE, 328–332.
3. Germán A Montoya, Horacio Riveros Ardila, Sebastián Rivera Díaz, and Yezid Donoso. 2011. Multi-objective evolutive algorithm for efficient coverage problem of WSN using a SPEA approach. In Proceedings of the 2011 international conference on applied, numerical and computational mathematics, and Proceedings of 2011 international conference on Computers, digital communications and computing. World Scientific and Engineering Academy and Society (WSEAS), 27–32.
4. Stefano Secci, Jean-Louis Rougier, and Achille Pattavina. 2008. On the selection of optimal diverse AS-paths for inter-domain IP/(G) MPLS tunnel provisioning. In 2008 4th International Telecommunication Networking Workshop on QoS in Multi-service IP Networks. IEEE, 235–241.
5. Zongyan Xu, Haihua Li, and Ye Guan. 2012. A study on the shortest path problem based on improved genetic algorithm. In 2012 Fourth International Conference on Computational and Information Sciences. IEEE, 325–3

ON NUMERICAL SOLUTION OF TWO POINT BOUNDARY VALUE PROBLEMS WITH LINEAR SHOOTING METHOD

Yücel BALTÜRK [1] Hüseyin DEMİR [2] İnci ÇİLİNGİR SÜNGÜ [3]

[1] Department of Mathematics, Ondokuz Mayıs University, Samsun, Turkey, **yucelbalturk@gmail.com**

[2] Department of Software Engineering, University of Samsun, Samsun, Turkey **huseyin.demir@samsun.edu.tr**

[3] Department of Mathematics and Natural Sciences Education, Ondokuz Mayıs University, Samsun, Turkey, **incicilingir@gmail.com**

Abstract

A practical linear shooting method is applied to the boundary value problem because of the importance of finding the initial value problems with an accurate way in physical application. It is assumed that the shooting method is the best and easiest way to solve boundary values problems, but there are some disadvantages when using the Newton Rapson's method of counting initial values. The solutions that are available in the literature are used to verify the results. Therefore the solution that obtained using linear shooting method is aimed to be fast, and accurate for solving this type of two-point boundary value problems.

Keywords: Linear Shooting Method; Numerical Solution; Boundary Value Problems.

REFERENCES

1. Adam, B. and Hashim, M.H.A. , Shooting Method In Solving Boundary Value Problem, IJRRAS 21 (1), 2014.

2. Edeo, A. , Solution of Second Order Linear and Nonlinear Two Point Boundary Value Problems Using Legendre Operational Matrix of Differentiation, American Scientific Research Journal for Engineering, Technology, and Sciences, 51,1,225-234, 2019.

3. Faragó, István & Filipov, Stefan. The Linearization Methods as a Basis to Derive the Relaxation and the Shooting Methods, 2020.

PROPERLY WEARING FACE MASK DETECTION USING HYBRID DEEP AND CLASSICAL MACHINE LEARNING TECHNIQUES

Sadaf Waziry[1], **Jawad Rasheed**[2], **Mirsat Yesiltepe**[3]

[1] Department of AI & Data Science, Istanbul Aydin University, Istanbul, Turkey
[2] Department of Computer Engineering, Istanbul Aydin University, Istanbul, Turkey
[3] Department of Mathematical Engineering, Yildiz Technical University, Istanbul, Turkey

sadafwaziry@stu.aydin.edu.tr; jawadrasheed@aydin.edu.tr; mirsaty@yildiz.edu.tr;

Abstract

The recent outbreak of COVID-19 around the world has caused global health catastrophe along with economic consequences. As per World Health Organization (WHO) this devastating crisis can be minimized and controlled by wearing mask properly. However, at public places or in chaos, manual check of persons wearing the masks properly or not is a hectic job and can cause panic. For such conditions, an automatic mask wearing system is desired. Therefore, this study analysed various deep learning pre-trained networks and classical machine learning algorithm that can automatically detect either a person is wearing the mask or not. For this, 40000 images are utilized to train and test four different models, namely; EfficientNetB0, InceptionV3, convolutional neural network and random forest in order to recognize facemasks in images. Besides just detecting the mask, the trained models also detect either the person is wearing the mask properly (covering nose and mouth), partially (mouth only), or wearing inappropriately (not covering nose and mouth). Experimental work reveals that InceptionV3 outperformed all three by attaining an overall accuracy of 98.40% and precision, recall, and F1-score of 98.30%.

Keywords: Deep learning; inappropriate wearing; machine learning; mask detection.

REFERENCES

1. M. Tan and Q. V. Le, “EfficientNet: Rethinking model scaling for convolutional neural networks,” 36th Int. Conf. Mach. Learn. ICML 2019, vol. 2019-June, pp. 10691–10700, 2019.
2. 1. C. Szegedy, V. Vanhoucke, S. Ioffe, J. Shlens, and Z. Wojna, “Rethinking the Inception Architecture for Computer Vision,” Proc. IEEE Comput. Soc. Conf. Comput. Vis. Pattern Recognit., vol. 2016-Decem, pp. 2818–2826, 2016, doi: 10.1109/CVPR.2016.308.

AUTOMATIC NOISE DETECTION SYSTEM IN QR CODES

Ahmad Bilal Wardak[1], **Jawad Rasheed**[2], **Mirsat Yesiltepe**[3]

[1] Department of AI & Data Science, Istanbul Aydin University, Istanbul, Turkey
[2] Department of Computer Engineering, Istanbul Aydin University, Istanbul, Turkey
[3] Department of Mathematical Engineering, Yildiz Technical University, Istanbul, Turkey

ahmadwardak@stu.aydin.edu.tr; jawadrasheed@aydin.edu.tr; mirsaty@yildiz.edu.tr;

Abstract

The resuscitation of the quick-response (QR) Code has been made possible in recent years by the expansion of mobile network coverage mixed with an increase in smartphone online content. They have become much more accessible as a result of the integration of a Code reader into smart devices, which has eliminated some inconvenient procedures and provided faster access to critical information. However, due to printer processes and limited printing technology, noise in printed images is unavoidable, which may reduce the quality of a QR code image during digital image collection and transfer, resulting in failure while scanning and extracting actual information. For this, a new dataset containing 20000 photos of original QR codes and noisy QR codes was created. Later, we used two well-known machine-learning classifiers and an advanced deep learning algorithm to separate noisy photos from original QR code images that includes convolutional neural network (CNN), support vector machine (SVM) and logistic regression. The experimental results demonstrate that CNN performed worst, while LG attained an overall accuracy of 97.25%, whereas SVM outperformed with 97.5% accuracy.

Keywords: noisy images, quick-response code, noise identification, CNN, SVM, logistic regression.

REFERENCES

1. Albawi, S., Mohammed, T.A., Al-Zawi, S.: Understanding of a convolutional neural network. In: 2017 International Conference on Engineering and Technology (ICET). pp. 1–6. IEEE (2017). https://doi.org/10.1109/ICEngTechnol.2017.8308186.
2. Peng, C.-Y.J., Lee, K.L., Ingersoll, G.M.: An Introduction to Logistic Regression Analysis and Reporting. J. Educ. Res. 96, 3–14 (2002). https://doi.org/10.1080/00220670209598786.

THE MATHEMATICAL MIRACLE OF AN 80 MINUTE MEMORY

Zeki Taştan[1], Esra Alkan[2]

[1]Department of Turkish Language and Literature, University of Van YYU, Van, Turkey
[2]Department of Turkish Language and Literature Education, University of Gazi, Ankara Turkey
[1]zekitastan@gmail.com,[2]alkanesra.63@gmail.com

Abstract

Born in Okayama in 1962 and actually graduated from Waseda University's Faculty of Letters, Yoko Ogawa is one of the prominent writers of Japanese literature. The author, who has more than thirty novels and stories, publishes a novel called "The Formula That The Professor Falls In Love", which was adapted to Japanese cinema in 2003. Pınar Demircan translated this highly acclaimed novel into Turkish in 2014 as *Housekeeper and the Professor* The novel is also interesting in that it shows Ogawa's mastery of mathematics, who is actually a literary man.

Housekeeper and the Professor is about the interesting and dramatic life of a mathematical genius who has been in a serious car accident and has to live with only the last 80 minutes of his memory. Finally, a maid with a ten-year-old child adapts and starts looking after the professor who is in need of care and does not rely on a maid. One of the most interesting aspects of the professor, who has a memory of 80 minutes and does not remember what he ate even for breakfast, is that his brain works miraculously when it comes to mathematical equations. The novel is also notable for instilling a love of mathematics in the professor, who establishes a warm bond with the ten-year-old boy.

***Keywords**:* Japanese literature, Yoko Ogawa, mathematics, novel, *Housekeeper and the Professor*

.

REFERENCES

1.) Yoko Ogawa, *Housekeeper and the Professor* (Stephen Snyder,) Newyork, 2008.
2.) Yoko Ogawa, *Profesör ve Hizmetçisi*, (Pınar Demircan), Pegasus, İstanbul, 2014

THE GROUP LAWS ON THE ALGEBRAIC CURVES

İsrafil Okumuş[1] **Ercan Çelik**[2]

[1] Erzincan Binali Yıldırım University, Faculty of Art and Science, Department of Mathematics, Erzincan-Turkey

[2]Kyrgyz-Turkish Manas University, Department of Applied Mathematics and Informatics, Bishkek-Kyrgyzstan

iokumus@erzincan.edu.tr, ercan.celik@manas.edu.kg

Abstract

In Algebraic Geometry, The Elliptic Curves is a smooth, projective and plane algebraic curves. The Elliptic Curve Cryptography (ECC) which independently proposeed Neal Koblitz and Victor Miller using the multiplicative group law on the elliptic curves [3-4]. It having been playing important role in public-key cryptogtaphy and traditionally are used in digital signature and in crypto currency. In this study we give properties the adding point on elliptic curve and invetigate the group law on the algebraic curves in plane.

Keywords: Algebraic curves, Elliptic curve, Cryptography.

REFERENCES

1. Neal Koblitz, Elliptic Curve Cryptosystems, Math. Comp. Vol: 48, 203-209, 1987.
2. Joseph H. Silverman, The arithmetic of elliptic curves, Graduate Texts in Mathematics, vol. 106, Springer-Verlag, New York, 1986. MR 817210
3. Neal Koblitz, Introduction to elliptic curves and modular forms, Graduate Texts in Mathematics, vol. 97, Springer-Verlag, New York, 1984. MR 766911
4. V. Miller, "Uses of elliptic curves in cryptography" in Advances in Cryptology Crypto 85, Springer Verlag, vol. 218, pp. 417-426, 1986.
5. A. J. Menezes, P. C. van Oorschot and S. A. Vanstone, Handbook of Applied Cryptography, CRC Press, 1996.

ON A STURM-LIOUVILLE FUZZY PROBLEM

Hülya Gültekin Çitil[1]

[1] Department of Mathematics, Faculty of Arts and Sciences, Giresun University, Giresun, Turkey

hulya.citil@giresun.edu.tr

Abstract

In this paper, we investigate the fuzzy eigenvalues and fuzzy eigenfunctions of a Sturm-Liouville fuzzy problem with fuzzy eigenvalue parameter in the boundary condition. Some properties of the problem are given. Several examples are solved.

Keywords: Sturm-Liouville fuzzy problem; Fuzzy eigenvalue; Fuzzy eigenfunction.

REFERENCES

1. C. T. Fulton, Two point boundary value problems with eigenvalue parameter contained in the boundary conditions, Proc. Soc. Edinburg, 77A, 293-308, 1977.
2. V. Lakshmikantham, R. N. Mohapatra, Theory of Fuzzy Differential Equations and Inclusions, Taylor and Francis, London, New York, 2003.
3. O. Kaleva, A note on fuzzy differential equations, Nonlinear Analysis, 64(5), 895-900, 2006.
4. A. Khastan, J. J. Nieto, A boundary value problem for second order fuzzy differential equations, Nonlinear Analysis, 72, 3583-3593, 2010.
5. H.-K. Liu, Comparison results of two-point fuzzy boundary value problems, International Journal of Computational and Mathematical Sciences, 5(1), 1-7, 2011.
6. H. Gültekin Çitil, N. Altınışık, On the Eigenvalues and the Eigenfunctions of the Sturm-Liouville Fuzzy Boundary Value Problem, Journal of Mathematical and Computational Science, 7(4), 786-805, 2017.

SOME PROPERTIES ON POINTWISE SEMI-SLANT RIEMANNIAN MAPS (PSSRM)

Mehmet Akif Akyol[1] and **Yılmaz Gündüzalp**[2]

[1] Department of Mathematics, University of Bingöl, Bingöl, Turkey

mehmetakifakyol@bingol.edu.tr,

[2]Department of Mathematics, University of Dicle, Diyarbakır, Turkey

ygunduzalp@dicle.edu.tr,

- This work is supported by 1001-Scientific and Technological Research Projects Funding Program of The Scientific and Technological Research Council of Turkey (TUBITAK) with project number 121F277.

Abstract

In this paper, we define and study a new class of Riemannian maps which is called *pointwise semi-slant Riemannian maps* from a Riemannian manifold to an almost Hermitian manifolds as a generalization of slant submanifolds, slant submersions, slant Riemannian maps, semi-slant submanifolds, semi-slant submersions, semi-slant Riemannian maps and pointwise slant Riemannian maps. We give many examples and investigate the geometry of total space, base space and the fibres.

Keywords: Riemannian maps; Semi-slant Riemannian maps; Pointwise semi-slant Riemannian maps; Second fundamental form of a map.

REFERENCES

1. M. A. Akyol, On Pointwise Riemannian Maps in Complex Geometry, International Symposium on Differential Geometry and Its Applications, Maulana Azad National Urdu University, Gachibowli, Hyderabad - 500032, Telangana, India (2022).
2. M. A. Akyol and Y. Gündüzalp, Pointwise slant Riemannian maps, In preprint, 2022.
3. B. Y. Chen and O. J. Garay, Pointwise slant submanifolds in almost Hermitian manifolds, Turk J. Math. 36, (2012), 630-640.
4. A. E. Fischer, Riemannian maps between Riemannian manifolds, Contemp. Math. 132 (1992), 331-366.
5. B. Şahin, Riemannian Submersions, Riemannian Maps in Hermitian Geometry, and their Applications, Elsevier, Academic Press, (2017).
6. K. Yano and M. Kon, Structures on manifolds, World scientific, 1985.

FRACTIONAL INVARIANTS OF CURVES IN LORENTZIAN PLANE

Meltem Öğrenmiş

Department of Mathematics, University of Fırat, Elazig, Turkey

meltemogrenmis@gmail.com,

Abstract

In this study, curves in the Lorentzian plane are discussed. By considering a spacelike (or timelike) curve in the Lorentzian plane, fractional invariants of a spacelike (or timelike) curve are given.

Keywords: Fractional derivative; Lorentzian plane; Frenet-Serret formulas.

REFERENCES

1. K. B. Oldham, J. Spanier, The Fractional Calculus: Theory and Applications of Differentiation and Integration to Arbitrary Order, Dover Publications, New York, 2006.
2. I. Podlubny, Fractional Differential Equations, Academic Press, San Diego, 1999.
3. M. Caputo, F. Mainardi, A new dissipation model based on memory mechanism, Pure and Applied Geophysics, 91(1), 134–147, 1971.
4. I. Grigorenko, E. Grigorenko, Chaotic dynamics of the fractional Lorenz system, Physical Review Letter, 91(3) Article ID 034101, 2003.
5. K. A. Lazopoulos, A. K. Lazopoulos, Fractional differential geometry of curves & surfaces, Progress in Fractional Differentiation and Applications, 2(3), 169–186, 2016.
6. S. I. Vacaru, Fractional nonholonomic Ricci flows, Chaos, Solitons Fractals 45(9-10), 1266–1276, 2012.
7. M. E. Aydın, M. Bektaş, A. O. Öğrenmiş, A. Yokuş, Differential Geometry of Curves in Euclidean 3-Space with Fractional Order, International Electronic Journal of Geometry, 14(1), 132-144, 2021.
8. T. Yajima, S. Oiwa, K. Yamasaki, Geometry of curves with fractional-order tangent vector and Frenet-Serret formulas, Fractional Calculus and Applied Analysis, 21(6), 1493–1505, 2018.

SOME PROPERTIES OF CURVES WITH FRACTIONAL DERIVATIVE IN EUCLIDEAN SPACES

Meltem Öğrenmiş

Department of Mathematics, University of Fırat, Elazig, Turkey

meltemogrenmis@gmail.com,

Abstract

In this presentation, a curve in n-dimensional Euclidean space is taken and the fractional derivative invariants of this curve are investigated. Later, these invariants are compared with the derivative results in the classical sense.

Keywords: Fractional derivative; Euclidean space; Frenet Formulas.

REFERENCES

1. D. Baleanu, K. Diethelm, E. Scalas, J.J. Trujillo, Fractional Calculus: Models and Numerical Methods. World Scientific, New Jersey, 2012
2. I. Podlubny, Fractional Differential Equations, Academic Press, San Diego, 1999.
3. T. Yajima, K. Yamasaki, Geometry of surfaces with Caputo fractional derivatives and applications to incompressible two-dimensional flows. J. Phys. A: Math. Theor. 45, No 6, 2012.
4. I. Grigorenko, E. Grigorenko, Chaotic dynamics of the fractional Lorenz system, Physical Review Letter, 91(3) Article ID 034101, 2003.
5. K. A. Lazopoulos, A. K. Lazopoulos, Fractional differential geometry of curves & surfaces, Progress in Fractional Differentiation and Applications, 2(3), 169–186, 2016.
6. A. Pressley, Elementary Differential Geometry, Springer Verlag, Berlin, 2010.
7. T . Yajima, T., H. Nagahama, Differential geometry of viscoelastic models with fractional-order derivatives. J. Phys. A: Math. Theor. 43, No 38, 2010.

Notes on Surfaces with Constant Mean Curvature along a given Curve in the Lie Group

Zuhal Kucukarslan Yuzbasi[1] **and Sevinc Taze**[2]

[1,2]Department of Mathematics, University of Firat, Elazig, Turkey

zuhal2387@yahoo.com.tr and svncatlla@gmail.com

Abstract

In this study, surfaces as linear combinations of the isoparametric curves and special Smarandache curves and their Frenet frames, respectively, are defined in the 3-dimensional Lie group. Then sufficient conditions when surfaces have constant mean curvature along the given curve are provided. Finally, examples of our results are given.

Keywords: Constant mean curvature; Lie group; Surfaces

REFERENCES

1- Wang G. J., Tang K.,Tai C. L. (2004). Parametric representation of a surface pencil with a common spatial geodesic, *Comput. Aided Des.* **36**, 447–459.
2- Bayram, E. (2022). Construction of surfaces with constant mean curvature along a timelike curve. *Politeknik Dergisi*, 1-1.
3- Yoon, D. W., Yüzbaşi, Z. K., (2019). On constructions of surfaces using a geodesic in Lie group. *J. Geo.,* **110**(2) (2019), 1-10.
4- Yoon, D. W., Yüzbaşi, Z. K., Bektaş, M. (2017). An approach for surfaces using an asymptotic curve in Lie group. *J. Advan. Phys.*, ***6***(4), 586-590.

Big Data Analytics via Penalized Estimation

Bahadir Yuzbasi [1]

[1] Department of Econometrics, University of Inonu, Malatya, Turkey

b.yzb@hotmail.com,

Abstract

We consider efficient estimation and prediction strategies for the classical multiple regression model with High Dimensional data when the dimensions of the parameters are larger than the number of observations. In this case, a number of existing penalized estimation techniques exists. However, they focus on selection variables with different number of predictors. Generally, the least absolute shrinkage and selection operator (Lasso) and Elastic Net approaches produce an over-fitted model compared with its competitors, namely the smoothly clipped absolute deviation (SCAD) method and adaptive Lasso (aLasso). Thus, prediction based only on a sub model selected by such methods will be subject to selection bias. To tackle with this problem, we suggest a shrinkage estimation that is combining two techniques, where one includes more predictors than the other based on relatively aggressive variable selection strategies, to improve the estimation and prediction performance. A Monte Carlo simulation study is carried out using the relative mean squared error (RMSE) criterion to appraise the performance of the listed estimators. The proposed strategy is applied to the analysis of a few real high-dimensional data sets.

Keywords: Sparse Regression Model; Penalized Estimation; High Dimensional Data

REFERENCES

1- Ahmed, S. E., & Yüzbaşı, B. (2016). Big data analytics: integrating penalty strategies. *International Journal of Management Science and Engineering Management*, *11*(2), 105-115.
2- Yüzbaşı, B., Arashi, M., & Ahmed, S. E. (2020). Shrinkage Estimation Strategies in Generalised Ridge Regression Models: Low/High-Dimension Regime. *International Statistical Review*, *88*(1), 229-251.
3- Ahmed, S. E., Kim, H., Yıldırım, G., & Yüzbaşı, B. (2016, June). High-Dimensional Regression Under Correlated Design: An Extensive Simulation Study. In *International Workshop on Matrices and Statistics* (pp. 145-175). Springer, Cham.
4- Yüzbaşı, B., Arashi, M., & Akdeniz, F. (2021). Penalized regression via the restricted bridge estimator. *Soft Computing*, *25*(13), 8401-8416.
5- Ahmed, S. E., & Yüzbaşı, B. (2017). High dimensional data analysis: Integrating submodels. In *Big and Complex Data Analysis* (pp. 285-304). Springer, Cham.
6- Yüzbaşı, B., & Ahmed, S. E. (2015). Shrinkage ridge regression estimators in high-dimensional linear models. In *Proceedings of the Ninth International Conference on Management Science and Engineering Management* (pp. 793-807). Springer, Berlin, Heidelberg.

Numerical Dynamics of Variable Order Fractional Benney-Lin Equation via Finite Element Collocation Method

Berat Karaagac [1], Alaattin Esen [2], Ercan Çelik [3], Edson Pindza [4]

[1] Department of Mathematics Education, University of Adıyaman, Adıyaman, Turkey

[2] Department of Mathematics, University of Inonu, Malatya, Turkey

[3] Kyrgyz-Turkish Manas University, Department of Applied Mathematics and Informatics, Bishkek-Kyrgyzstan

[4] Department of Mathematics and Statistics, University of Tshwane, Pretoria, South Africa.

bkaraagac@adiyaman.edu.tr, alaattin.esen@inonu.edu.tr, ercan.celik@manas.edu.kg, pindzaedson@yahoo.fr

Abstract

The aim of this paper is to focus a numerical study on variable order fractional Benney-Lin equation which specifically describes the propagation of long waves on thin film. The method is based on Finite element collocation method with selection of basis functions as trigonometric quintic B-splines. With the idea of using less order basis, the Benney-Lin equation transformed to couple system via auxiliary variable. The Finite element collocation method through satisfying boundary conditions and continuity conditions, yields the equation convert to an algebraic system of equations whose solutions generates the approximate solution for fractional Benney-Lin equation. The effectiveness and efficient of the method is tested via some illustrative examples.

Keywords: Variable order fractional Benney-Lin equation; Collocation method; Trigonometric quintic B-splines

REFERENCES

1. M. H. Heydari, M. Razzaghi, Jacobi spectral method for variable-order fractional Benney–Lin equation arising in falling film problems, Journal of Computational and Applied Mathematics, Vol*:* 402, 113813, 2022.

2. W. Gao, P. Veeresha, D. G. H. M. Prakasha, Baskonus, New numerical simulation for fractional Benney–Lin equation arising in falling film problems using two novel techniques. Numerical Methods for Partial Differential Equations, Vol:*3*7, No:1, 210-243. 2021.

COMPLEX TRANSFORM METHOD FOR SYSTEM OF FRACTIONAL DIFFERENTIAL EQUATIONS AND APPLICATIONS

Elham Sefidgar[1], Ercan Çelik[2], Babak Shiri[3]

[1] Ataturk University Faculty of Science, Department of Mathematics, Erzurum-Turkey

[2] Kyrgyz-Turkish Manas University, Department of Applied Mathematics and Informatics, Bishkek-Kyrgyzstan

[3] Faculty of mathematical science, University of Tabriz, Tabriz - Iran

e_sefidgar@yahoo.com, ercan.celik@manas.edu.kg, shiri@tabrizu.ac.ir

Abstract

The complex transform method for solving system of the system of fractional differential equations introduced. We applied complex transform method for solving a system appeared in the dynamics of the drug therapy efficiently. Various examples shows that these methods are effective and efficient.

Keywords: Complex Transform method, Fractional calculus, System of fractional differential equations, Riemann-Liouville fractional derivative.

REFERENCES

[1] Agrwal R. P. A propos d'une note de M. Pierre Humbert, C. R. Se'ances Acad. Sci., v.236, n.21, 1953, pp. 2031-2033.

[2] Ahmad B., Nieto J.J. Existence results for a coupled system of nonlinear fractional differential equations with three-point boundary conditions, Computers and Mathematics with Applications, v.58, n.9, 2009, pp. 1838–1843.

[3] Bai Z, L H. Positive solutions for boundary value problem of nonlinear fractional differential equation, Journal of Mathematical Analysis and Applications, v.311, n.2, 2005, pp. 495–505.

[4] Daftardar-Gejji V. Babakhani A. Analysis of a system of fractional differential equations, Journal of Mathematical Analysis and Applications, v.293, n.2, 2004, pp. 511–522.

[5] Daftardar-Gejji V. Positive solutions of a system of non-autonomous fractional differential equations. Journal of Mathematical Analysis and Applications, v.302, n.1, 2005, pp. 56–64.

BLOCKCHAIN-BASED SERVICE NETWORK PORTAL

Hande Yüzel[1], Onur Toklu[1], Senem Şahan Vahaplar[1], M. Fatih Akay[2], Sevtap Erdem[2]

[1] Saha Information Technologies, İstanbul, Turkey

[2] Department of Computer Engineering, Çukurova University, Adana, Turkey

senem.vahaplar@sahabt.com

Abstract

Blockchain-based Service Network (BSN) is a worldwide infrastructure network that provides developers and companies a one-stop-shop solution for blockchain and distributed ledger technology applications, that can operate together within one environment. Red Date Technology, the architect of BSN, signed an agreement with Turkey-China Economic Twinning Center (TUCEM) in order to establish an international BSN portal in Turkey, which is Turkey's first blockchain-based service network. By means of the blockchain-based service network provided by BSN, the users can run and install all kinds of distributed applications and the developers can build and deploy decentralized applications at low cost.

In this study, the new BSN portal developed to offer "blockchain as a service (BaaS)" in Turkey is presented. User Interface is developed with React.js, the backend is developed with .Netcore and PostgreSQL is used for database management.

Keywords: Blockchain; Blockchain-based Service Network; Blockchain as a service (BaaS).

REFERENCES

1. Blockchain-based Service Network User Manual, BSN Foundation, 2019.
2. Wenli Yang, Erfan Aghasian, Saurabh Garg, David Herbert, Leandro Disiuta, Byeong Kang, A Survey on Blockchain-Based Internet Service Architecture: Requirements, Challenges, Trends, and Future, IEEE Special Section on Neural Engineering Informatics, Vol:7, 75845-75872, 2019.
3. Mayra Samaniego, Uurtsaikh Jamsrandorj, Ralph Deters, Blockchain as a Service for IoT, 2016 IEEE International Conference on Internet of Things (iThings) and IEEE Green Computing and Communications (GreenCom) and IEEE Cyber, Physical and Social Computing (CPSCom) and IEEE Smart Data (SmartData), 433-436, 2016.

Relative TL Bi-Ideal, TL Quasi Ideal, TL Interior Ideal

Emine Funda Okumuş[1]

[1] Department of Mathematics, University of Karadeniz Tecnical, Trabzon, Turkey

eminefundaekinci@ktu.edu.tr

Abstract

In this paper, we introduce the notions of $TL-$bi ideals, $TL-$quasi ideals and $TL-$interior ideals in a semigroup, develop some properties of them. We also build the relation between $L-$ideals of a semigroups and TL-ideals of a TL subsemigroups. Furthermore, the lattice structure of TL bi ideal, TL quasi ideal, $TL-$interior ideal of a TL subsemigroups are studied.

Keywords: Semigroup,relative ideal,t-norm ,bi-ideal,quasi ideal.

REFERENCES

1. Ajmal N.,Jahan I.: A Study of normal fuzzy subgroups and characteristic fuzzy subgroups of a fuzzy group. Fuzzy Inf. Eng.2012; 123:143-2
2. Wallace, A. D.: Relative Ideals in Semigroups II. Acta math. seient. himg. 1963; 137:148-14
3. Yu Y. ,Wang Z.:TL-subrings on TL-ideals Part 1. basic concepts . Fuzzy set and Systems 1994;93:103-68.
4. Wallace, A. D:Relative Ideals in Semigroups I. Colloq. math. 1962; 55:61-0
5. Zadeh L.A: Fuzzy sets . Inform. And Control 1965;338:353-8
6. Klement E.P. , Mesiar R. , Pap E.: Triangular norms , Kluwer Academic Publishers, Dordrecht 2000.

A MATHEMATICAL EXAMINATION OF THE FRACTIONAL HIV INFECTION MODEL

Bahar Acay[1]

[1]Department of Mathematics, University of Firat, Elazig, Turkey

bacay@firat.edu.tr,

Abstract

The objective of this study is to investigate the HIV infection model by means of the Caputo fractional differential operator. We present some crucial theoretical results and basic aspects of the non-integer order disease model under consideration. Moreover, the dynamic behavior of the system is analyzed in detail with the help of a convenient numerical scheme based on the Caputo operator. On the other hand, we employ the non-linear least squares curve fitting method in order to estimate the parameters of the suggested model under real-world data of Turkey.

Keywords: Fractional modeling, HIV infection, Mathematical biology, Caputo operator.

REFERENCES

1. Acay, B., Inc, M., Khan, A., & Yusuf, A. (2021). Fractional methicillin-resistant Staphylococcus aureus infection model under Caputo operator. Journal of Applied Mathematics and Computing, 67(1), 755-783.
2. Nagle, R. K., Saff, E. B., & Snider, A. D. (2014). Fundamentals of differential equations. Pearson Education.

A numerical stochastic computing paradigms for the nonlinear singular delay differential problems

Zulqurnain Sabir[1,2], Haci Mehmet Baskonus[3], Fevzi Erdoğan[4]

[1]Department of Mathematical Science, College of Science, United Arab Emirates University, Al Ain, Abu Dhabi, UAE

Email: zulqurnain_s@uaeu.ac.ae

[2]Department of Mathematics and Statistics, Hazara University, Mansehra, Pakistan

Email: zulqurnain_maths@hu.edu.pk

[3]Department of Mathematics and Science Education, Faculty of Education, Harran University, Sanliurfa, Turkey

Email: hmbaskonus@gmail.com

[4]Department of Mathematics, Faculty of Sciences, Yuzuncu Yil University, Van, Turkey

Email: ferdogan@yyu.edu.tr

Abstract

The aim of this work is to present a computational intelligence procedure for solving the nonlinear singular delay differential problems. The stochastic procedures are adopted by using the strength of artificial neural networks (ANNs), together with the optimization schemes of global search genetic algorithm and local search active set programming. An objective function is designed by using the sense of differential singular delay differential model and its boundary conditions. The optimization of this objective function is performed through the hybridization of the global and local search schemes. The objective for presenting this research is to provide the reliable structure of the ANNs to deal with such stiff nonlinear, singular based on the delay differential models.

Keywords: Delay models, singular, nonlinear, active-set programming, genetic algorithm, statistical analysis

References

[1] Adel, W. and Sabir, Z., 2020. Solving a new design of nonlinear second-order Lane–Emden pantograph delay differential model via Bernoulli collocation method. The European Physical Journal Plus, 135(5), pp.1-12.

[2] Sabir, Z., Raja, M.A.Z., Le, D.N. and Aly, A.A., 2021. A neuro-swarming intelligent heuristic for second-order nonlinear Lane–Emden multi-pantograph delay differential system. Complex & Intelligent Systems, pp.1-14.

[3] Guirao, J.L., Sabir, Z. and Saeed, T., 2020. Design and numerical solutions of a novel third-order nonlinear Emden–Fowler delay differential model. Mathematical Problems in Engineering, 2020.

[4] Sabir, Z., Guirao, J.L. and Saeed, T., 2021. Solving a novel designed second order nonlinear Lane–Emden delay differential model using the heuristic techniques. Applied Soft Computing, 102, p.107105.

[5] Sabir, Z., Wahab, H.A., Nguyen, T.G., Altamirano, G.C., Erdoğan, F. and Ali, M.R., 2022. Intelligent computing technique for solving singular multi-pantograph delay differential equation. Soft Computing, pp.1-13.

Artificial intelligence for real-time defect detection and inspection of photovoltaic panels

Saloua SENHAJI1,2 Sanaa FAQUIR1, Mohamed HAMLICH2 , Omar Diouri1, Mohammed Ouazzani Jamil1

1Laboratoire Systèmes et Environnement Durables (SED), Faculté des sciences de l'ingénieur (FSI), Université Privée de Fès (UPF), Fez, Morocco

2Laboratoire Complex Cyber Physical Sysem, ENSAM ,Université Hassan II, Casablanca, Morocco

senhaji@upf.ac.ma

Abstract

Photovoltaic installations can be a lucrative investment. To maximise production and income over decades, however, you need to focus on quality. The solar module, the key element of these installations, must be reliable, producing electricity continuously for years. However, they are confronted with different types of failures. Therefore, the introduction of intelligent devices for real-time inspection and monitoring of photovoltaic panels is a hot topic. In this paper, we propose an intelligent monitoring system based on visual data acquired by thermal cameras connected with microcontrollers equipped with our own convolutional neural network model to detect and process hot spots using Tensorflow and Keras resulting in an intelligent and real-time inspection of large solar farms.

Keywords: Artificial intelligence; convolutional neural network ; photovoltaic panels; fault detection; real –time monitoring.

REFERENCES

1. IEA. Snapshot of global photovoltaic markets. April 2020, https://iea-pvps.org/ wp-content/uploads/2020/04/IEA_PVPS_Snapshot_2020.pdf.
2. Kontges M, Kurtz S, Jahn U, Berger K, Kato K, Friesen T, Liu H, Iseghem MV. Review of failures of photovoltaic modules. IEA PVPS Task; 2014. T13-01. [available at: https://iea-pvps.org/key-topics/review-of-failures-of-photovoltaicmodules-final/.
3. Senhaji, S., Faquir, S., Ouazzani Jamil, M., Towards robotics and artificial intelligence for the prevention of Covid 19 pandemic. E3S Web of Conferencesthis link is disabled, 2021, 229, 01035

The nonlinear fractional generalised Pochhammer-Chree equation arising an electrical field theory is Dynamical Analysis of Caputo-Fabrizio fractional derivative

Ajay Kumar [a,*], Haci Mehmet Baskonus[b]

[a] *Department of Mathematics, Bakhtiyarpur College of Engineering, Champapur, Dedaur, Bakhtiyarpur-803212, India.*

[b] *Department of Mathematics and Science Education, Faculty of Education, Harran University, Sanliurfa, Turkey*

Abstract

In this manuscript, an efficient numerical technique called the fractional homotopy perturbation transform method (FHPTM) is presented for solving the nonlinear fractional generalized Pochhammer-Chree equation. The basic concepts for fractional derivatives are defined in Caputo Fabrizio. An existence and uniqueness analysis of the considered mathematical model is provided. To present the physical representation, several cases are given, and the findings are illustrated by certain surface plots. The findings suggest that the new methodology is prolific, powerful, efficient, and simple to apply, as well as capable of incorporating a wide range of partial fractional differential equations.

Keywords: Homotopy perturbation transform method; Caputo fabrizio derivative; Laplace transform; generalized Pochhammer-Chree equation; Fractional differential equations

References

[1] M. Caputo, Elasticita Dissipazione, Zani-Chelli, 1969.

[2] K. S. Miller, B. Ross, An introduction to fractional calculus and fractional differential equations, A Wiley, New York, 1993.

[3] L.Akinyemi, P.Veeresha, M.Senol, H.Rezazadeh, An efficient technique for generalized conformable Pochhammer-Chree models of longitudinal wave propagation of elastic rod,Indian J Phys, 1-10, 2022.

*kajay9249@gmail.com (A. Kumar)

Preprint submitted to 6th International Conference on Computational Mathematics and Engineering Sciences / 20-22 May. 2022,

EXACT SOLUTIONS OF GOURSAT PROBLEMS INVOLVING MIXED PARTIAL DERIVATIVES VIA HOMOTOPY ANALYSIS METHOD

Mine Babaoglu

Department of Mathematics and Science Education, University of Kahramanmaras Sutcu Imam, Kahramanmaras, Turkey

mnbabaoglu@gmail.com

Abstract

The present research explores, the Homotopy analysis method (HAM) is tested by performing it to obtain the exact solutions for the Goursat problems involving mixed partial derivatives. We carry out the computations, tables and draw the 2D and 3D graphs in this paper with the help of a ready-made package program. Additionaly, gained results and numerical simulations emphasize the forcefulness of the proposed method.

Keywords: Homotopy analysis method; Mixed partial derivatives; Goursat problems; Nonlinear.

REFERENCES

1. S. J. Liao, Beyond Perturbation: Introduction to the Homotopy Analysis Method, Boca Raton.: Champan & Hall/CRC Press, 3-38, 2003.
2. A. M. Wazwaz, Partial Differantial Equations: Methods and Applications, Balkema Publishers, The Netherlands, 2002.
3. S. J. Liao, On the homotopy analysis method for nonlinear problems, Applied Mathematics and Computation, 147(2), 499-513, 2004.
4. J. H. He, Comparison of Homotopy perturbation method and Homotopy analysis method, Applied Mathematics and Computation, 156(2), 527-539, 2004.

ON $\rho-$ STATISTICAL CONVERGENCE OF DOUBLE SEQUENCES OF ORDER α IN TOPOLOGICAL GROUPS

Nazlım Deniz Aral[1], Hacer Şengül Kandemir[2] and Mikail Et[3]

[1]Department of Mathematics, Bitlis Eren University, Bitlis, Turkey

[2]Faculty of Education, Harran University, Şanlıurfa, Turkey

[3]Department of Mathematics, Fırat University, Elazığ, Turkey

ndaral@beu.edu.tr

hacer.sengul@hotmail.com

mikailet68@gmail.com,

Abstract

In this study, by using definition of ρ-statistical convergence which was defined by Çakallı [1], we give some inclusion relations between the concepts of ρ-statistical convergence and statistical convergence of double sequences of order α in topological groups.

Keywords: Topological groups; Statistical convergence; ρ-statistical convergence.

REFERENCES

1. Hüseyin Çakallı, A variation on statistical ward continuity, Bull. Malays. Math. Sci. Soc. Vol:40, 1702-1710, 2017.
2. Ekrem Savaş, Lacunary statistical convergence of double sequences in topological groups, J. Inequal. Appl., Article Number: 480, 2014.
3. Hacer Şengül Kandemir, On I–Deferred Statistical Convergence in Topological Groups, Maltepe Journal of Mathematics, Vol:1, Number:2, 48-55, 2019.
4. Hacer Şengül, Mikail Et, Hüseyin Çakallı, Deferred statistical convergence of order α in topological groups, AIP Conference Proceedings, 2183, 050014, 2019.

DIRECT INTEGRATION OF THE SYSTEMS OF LINEAR DIFFERENTIAL EQUATIONS AND SOME APPLICATIONS IN ECONOMICS

Urdaletova A.B.[1], **Kydyraliev S. K.**[2]

[1] Kyrgyz-Turkish Manas University, Department of Mathematics, Bishkek-Kyrgyzstan

[2]American University of Central Asia (Bishkek), Kyrgyzstan

anarkul.urdaletova@manas.edu.kg, kydyraliev_s@auca.kg

Abstract

Differential equations play an important role in many disciplines, including engineering, physics, economics, and biology, because studying the relationship between quantities and their rates of change, which are expressed by derivatives, is an important scientific task. In the course of ordinary differential equations, when studying linear systems, one usually studies the methods of Leonhard Euler (1707 – 1783) and, sometimes, Jean Léron d'Alembert (1717 - 1783). A combination of these methods, involving the use of characteristic numbers to obtain integrable combinations, as shown in the works of the authors of the report, generates a synergy effect — significantly simplify the solution process. As a result, the direct integration method turns out to be a very convenient tool for studying applied problems. In this paper, the emphasis is on applications in economics.

Keywords: Linear ordinary differential equations, solution of systems of linear equations, characteristic numbers, integrable combinations, integrability by quadrature.

REFERENCES

1. Kydyraliev S.K., Urdaletova A.B. Direct Integration of Systems of Linear Differential and Difference Equations / S.K. Kydyraliev, A.B. Urdaletova // University of Niˇs, Serbia, Filomat 33:5 (2019), p.1453–1461.

2. Kydyraliev S.K., Urdaletova A.B Novyj metod resheniya sistem lineynyh differentsial'nyh i raznostnyh uravneniy / S.K. Kydyraliev, A.B. Urdaletova // Vestnik KAZNU.- Almaty., 2009.- №4 (63). - P.21-29.

3. Guterman, M.M. Differential Equations. A Fist Course. /M.M. Guterman, Z.H. Nitecki. - Philadelphia: Saunders College Publishing, 1988. - 689 p.

4. Bailey D. Mathematics in Economics. /D. Bailey. - Bercshire, England: McGraw-Hill, Inc, 1998. - 485 p.

Some special Smarandache Ruled Surfaces according to Frenet Frame in E^3 -III

[1]Süleyman ŞENYURT, [2]Davut CANLI, [3]Elif ÇAN
[1,2,3]Faculty of Arts and Sciences, Department of Mathematics,Ordu University, Ordu, TURKEY
[1]ssenyurt@odu.edu.tr, [2]davut_canli@hotmail.com, [3]eliff.cann@hotmail.com

Abstract.
In this study, firstly, Smarandache curves were defined using Frenet vectors of a curve and Darboux vector. Then, Gaussian and average curvature of ruled surfaces formed by the vectors obtained from Frenet and Darboux vectors along Smarandache curves are calculated separately and openability conditions are given. Finally, the ruled surfaces obtained by using the Vivian curve were drawn with the maple 17 program.

Key Words: Smarandache ruled surfaces, mean curvature, Gaussian curvature, vivian curve

References

[1] Ali, A.T., Special Smarandache curves in the Euclidean space, International Journal of Mathematical Combinatorics, 2(2010),30-36, 2010

[2]Bektaş, Ö, Yüce,S., Special Smarandache Curves According to Darboux Frame in E^3, Romanian Journal of Mathematics and Computer Science, cilt.3, ss.48-59, 2013

[3]Berk, A., A Structural Basis for Surface Discretization of Free Form Structures: Integration of Geometry, Materials and Fabrication, Ph. D Thesis, Michigan University, ABD, 2012

[4]Çetin , M, Kocayiğit,H., On the Quaternionic Smarandache Curves in Euclidean 3-Space, Int. J. Contemp. Math. Sciences, Vol. 8, 2013, no. 3, 139 – 150

[5]Do-Carmo,P.M., Differential geometry of curvesand surfaces. IMPA, 1976

[6]Fenchel,W., On the Differential Geometry of Closed Space Curves, Bulletin of American Mathematical Society, 57(44-54), 1951

[7]Gray A. Modern differential geometry of curves and surfaces with Mathematica, 1997

[8]Ouarab, S., Smarandache Ruled Surfaces according to Frenet-Serret Frame of a Regular Curve in E3, Hindawi Abstract and Applied Analysis, Vol. 2021, Article ID 5526536, 8 pages

[9]Ouarab, S., Smarandache ruled Surfaces according to Darboux Frame in E^3, Hindawi Journal of Mathematics, Vol. 2021, Article ID 9912624, 10 pages

[10]Ouarab,S., NC-Smarandache ruled surface and NW-Smarandache ruled surface according to alternative moving frame in E^3, HindawiJournal of Mathematics ,Vol. 2021, Article ID 9951434, 6 pages

Machine Learning based Intrusion Detection System using Gray Wolf Optimization for Feature Selection

Oğuzhan Taş[1], Amani Yahyaoui[2]

[1,2] Department of Computer Engineering, Istanbul Sabahattin Zaim University, Istanbul, Turkey

oguzhantas@gmail.com, amani.yahyaoui@izu.edu.tr

Abstract

In the age of information and technology, heterogeneous devices constantly connected to each other share information between different applications and generate large amounts of data traffic. It is important to ensure that this traffic, which occupies an important place in the digital workflow, is not disrupted and that cybersecurity is ensured. Network-based attacks are an important part of cybersecurity issues. Since conventional signature-based intrusion detection systems cannot detect zero-day attacks, machine learning algorithms have been widely used in this field.

In this paper, we propose a model using the AdaBoost ensemble classifier and grey wolf optimization algorithm for feature selection to detect attacks that threaten network security.

As a result of the training and testing studies on the NSL-KDD dataset, an accuracy rate of 89.21% was achieved.

***Keywords**:* Intrusion detection, Network Security, Machine Learning, Grey Wolf Optimization

REFERENCES

1. S. Mirjalili, S. M. Mirjalili, and A. Lewis, “Grey wolf optimizer,” Advances in Engineering Software, vol. 69, pp. 46–61, 2014.
2. Schapire, Robert E. “Explaining adaboost.” In Empirical inference, pp. 37-52. Springer, Berlin, Heidelberg, 2013.
3. Mohammadi, Mokhtar, Tarik A. Rashid, Sarkhel H. Taher Karim, Adil Hussain Mohammed Aldalwie, Quan Thanh Tho, Moazam Bidaki, Amir Masoud Rahmani, and Mehdi Hosseinzadeh. "A comprehensive survey and taxonomy of the SVM-based intrusion detection systems." Journal of Network and Computer Applications 178 (2021): 102983.

EIGENFUNCTION EXPANSIONS FOR A STURM-LIOUVILLE OPERATOR FOR THE CASE OF AN INFINITE INTERVAL

Samet Uytun [1]

[1] Institute of Science and Technology, University of Binali YILDIRIM , Erzincan, Turkey

sametuytun@gmail.com,

Abstract

A Sturm-Liouville problem is said to be regular if the interval [a,b] is finite and the function *q(x)* is summable over this interval. In the contrary case, i.e. if the interval [a,b] is finite or the function *q(x)* is not summable over [a,b] (or both), then Sturm-Liouville problem is said to be singular.

We obtained the expansion theorem for a regular Sturm-Liouville problem by means of a passage to the limit, replacing the differential equation by a finite-differance equation.

A similar idea makes it possible to obtain the expansion theorem for a singular problem, if we consider it as the limit of regular problems.

We will start by considering the case in which the interval is the halfline [0, ∞), and the function *q(x)* is continuous on this interval.

.

Keywords: Expansıons Functıons of Sturm-Lıouville.

REFERENCES

1. The results of this section are due to Weyl .Another prof was given by Titchmarsh.The prof of the Perseval equality presented here was given by Levitan and independently by Levinson and Yoshida.
2. Introduction to Spectral Theory by B.M Levitan and I.S. Sargsjan

Application of a Genetic Algorithm for Generating Regular and Irregular Natural Fractal Patterns

Habiba Akter[1], Rupert Young[2]

Department of Engineering and Design,

School of Engineering and Informatics,

University of Sussex,

United Kingdom

h.akter@sussex.ac.uk, r.c.d.young@sussex.ac.uk

Abstract

This work aims to describe the successful implementation of a Genetic Algorithm (GA) as a search and optimisation tool for the fractal patterns observed in nature. This will involve designing a method for calculating fractal dimensions based on parameter values of the fractal-generating functions. These values will be evolved and optimised using a standard GA. The fact that Genetic Algorithms are closely associated with nature, has been the main motivation behind this work since the natural complex patterns are the most common fractals. However, we aim to optimise both self-similar and irregular fractal patterns using GA.

Keywords: Genetic Algorithm; Fractal Patterns; Optimisation.

REFERENCES

1. Michael F Barnsley and Stephen Demko. 1985. Iterated function systems and the global construction of fractals. Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences 399, 1817 (1985), 243–275.
2. Paul Bourke. 2014. Box counting fractal dimension of volumetric data.
3. Paul Campbell and S Abhyankar. 1978. Fractals, form, chance and dimension.
4. Kenneth Falconer. 2004. Fractal geometry: mathematical foundations and applications. John Wiley & Sons.
5. Manuel Fernández-Martínez and MA Sánchez-Granero. 2014. Fractal dimension for fractal structures. Topology and its Applications 163 (2014), 93–111.
6. Benoit B Mandelbrot and Benoit B Mandelbrot. 1982. The fractal geometry of nature. Vol. 1. WH freeman New York.

6th INTERNATIONAL CONFERENCE ON COMPUTATIONAL MATHEMATICS AND ENGINEERING SCIENCES

www.cmescongress.org