SAKARYA ÜNİVERSİTESİ
FEN BİLİMLERİ ENSTİTÜSÜ DERGİSİ

Sakarya University Journal of Science
SAUJS

ISSN 1301-4048 | e-ISSN 2147-835X | Period Bimonthly | Founded: 1997 | Publisher Sakarya University | http://www.saujs.sakarya.edu.tr/

Title: On the Properties of r-Circulant Matrices Involving Generalized Fermat Numbers

Authors: Bahar KULOĞLU, Engin ESER, Engin ÖZKAN

Recieved: 2023-04-10 00:00:00

Accepted: 2023-06-13 00:00:00

Article Type: Research Article

Volume: 27
Issue: 5
Month: October
Year: 2023
Pages: 956-965

How to cite
Bahar KULOĞLU, Engin ESER, Engin ÖZKAN; (2023), On the Properties of r-Circulant Matrices Involving Generalized Fermat Numbers. Sakarya University Journal of Science, 27(5), 956-965, DOI: 10.16984/saufenbilder.1280572
Access link
https://dergipark.org.tr/tr/journal/1115/issue/80257/1280572

New submission to SAUJS
http://dergipark.gov.tr/journal/1115/submission/start

Sakarya University Journal of Science 27(5), 956-965, 2023

On the Properties of r-Circulant Matrices Involving Generalized Fermat Numbers

Bahar KULOĞLU [*1], Engin ESER[2], Engin ÖZKAN[3]

Abstract

r-circulant matrices have applied in numerical computation, signal processing, coding theory, etc. In this study, our main goal is to investigate the r-circulant matrices of generalized Fermat numbers which are shown by $GR_{a,b,n} = a^{2^n} + b^{2^n}$ We obtain the eigenvalues, determinants, sum identity of matrices. Also we find upper and lower bounds for the spectral norms of generalized Fermat r-circulant matrices. Beside these, we present $GR^*_{a,b,r}$ matrix in the form of the Hadamard product of two matrices as $GR^*_{a,b,r} = A.B$. In addition, we get the right and skew-right circulant matrices for $r = 1$ and $r = -1$. Finally, we examine their different norms (Spectral and Euclidean) and limits for matrix norms.

Keywords: Fermat number, generalized Fermat numbers, norm (spectral and Euclidean), eigenvalues, circulant matrices

1. INTRODUCTION

Special numbers were created by grouping prime numbers within themselves. The best known of these are Pierre de Fermat and Marin Mersenne. Fermat numbers included pseudoprimes. First, Fermat conjectured that all numbers produced by $2^k + 1$ (k is a nonnegative integer) are prime numbers (k must be the power of 2).

It can be easily seen that the first 5 numbers are prime, but when it comes to the 6th number there is a problem. Later, Euler proved that this number has factors. So, it was a composite number. Fermat had made a calculation error. Now an open question: Are there any other numbers like this? Mersenne numbers in the form $2^k - 1$ were studied in antiquity because of their connection with perfect numbers. Euclid-Euler theorem gives this connection.

Later, Francois Proth studied Fermat numbers. Proth gave a generalized version of Fermat numbers in the form $k2^n + 1$ where $2^n > k$ for $n, k \in \mathbb{Z}^+$ and k is an odd number. The Proth numbers are also known

[*]Corresponding author: bahar_kuloglu@hotmail.com (B. KULOGLU)
[1]Hatuniye Hatuniye Vocational and Technical Anatolian High School, Ministry of Education, Mardin, Türkiye
[2]Graduate School of Natural and Applied Sciences, Erzincan Binali Yıldırım University, Erzincan, Türkiye
[3]Department of Mathematics, Faculty of Arts and Sciences, Erzincan Binali Yıldırım University, Erzincan, Türkiye
E-mail: engineser1978@gmail.com, eozkan@erzincan.edu.tr
ORCID: https://orcid.org/0000-0001-7624-8270, https://orcid.org/0000-0001-5965-4162, https://orcid.org/0000-0002-4188-7248

Content of this journal is licensed under a Creative Commons Attribution-Non Commercial No Derivatives 4.0 International License.

as k-Fermat numbers [1]. From this, it can be easily seen that the general forms of Proth numbers without any restrictions are k-Fermat numbers. The first terms of these numbers are as follows:

3, 5, 13, 17, 41, 97, 113, 193, 241, 257, 353, 449, 577, 641, 673, 769, 929, 1153, 1217, 1409, 1601, 2113, 2689, 2753, 3137, 3329, 3457, 4481, 4993, 6529, 7297, 7681, 7937, 9473, 9601, 9857

(Proth numbers are referenced in the On-Line Encyclopedia of Integer Sequences in OEIS as: A080076 [2])

Generalized Fermat numbers [3], denoted by $\{GR_{a,b,n}\}$ is defined by

$$GR_{a,b,n} = a^{2^n} + b^{2^n}, \quad (1)$$

where a and b are any relatively prime integers, $a > b > 0$, $a \geq 2,\ n \geq 0$. The first terms of these numbers as follow:

n	0	1	2	3	…
$GR_{a,b,n}$	$a + b$	$a^2 + b^2$	$a^4 + b^4$	$a^8 + b^8$	…

These matrices are shown by C_r . Their inverse, conjugate, transposes, sums, and multiplications have been calculated in [4].

The algebraic properties of these matrices have been investigated in [5, 6]. For any r complex number except zero, we can define those matrices. We can analyze its eigenvalues, Euclidean norm, spectral norm, determinants and inverse. C_r is determined by its first-row element and r. Given $r = +1$ and $r = -1$ in recurrence relation, we obtain its eigenvalues and determinant for right and skew right circulant matrices.

Circulant matrices and r-circulant matrices including such as Fibonacci, Pell, Pell-Lucas numbers have been great interest. In several studies, eigenvalues, determinants, norms, bounds and inverses for these matrices are found. For instance, Bueno [7] presented eigenvalues and the determinant of the right circulant matrices with Pell numbers and Pell-Lucas numbers.

Solak [8] presented norms for circulant matrices including Fermant and Mersenne numbers. S.-Q., Shen et al. [9, 10] presented the bounds for r-circulant matrices with Fibonacci and Lucas numbers and also presented the spectral norms with k-Fibonacci and k-Lucas numbers. Zheng et al. [11] showed the exact inverse of circulant matrices with Fermat and Mersenne numbers. Bozkurt [12] solved the determinants of these matrices using matrix decomposition. Kumari et al. studied on the properties of the r-circulant matrices involving Mersenne and Fermat numbers [13].

Marin has also discussed examples of concrete usage areas of studies similar to those in [14, 15]. Pucanovic et al. by taking the $r-$circulant matrix and Chebyshev polynomials as the subject, handled the inputs of the r-circulant matrices by constructing them from Chebyshev polynomials [16]. In this paper, we study $r-$circulant matrices with generalized Fermat numbers.

Lemma 1.1 [17, 18] Let $X = (x_{ij})$ be a matrix.

The Frobenius or Euclidean norm of X is defined as

$$\|X\|_{E=F} = \sqrt{\sum_{i=1}^{a}\sum_{j=1}^{b}\left|x_{ij}\right|^2}.$$

The column norm of X is defined as

$$\|X\|_1 = \max_{1 \leq j \leq n} \sum_{j=1}^{n}\left|x_{ij}\right|.$$

The row norm of X is defined as

$$\|X\|_\infty = \max_{1 \le j \le m} \sum_{j=1}^{n} |x_{ij}|.$$

The spectral norm of a matrix X defined as

$$\|X\|_2 = \sqrt{\gamma(X^*X)} = \mu_{max}(X).$$

where $\gamma(X^*X)$ denote the eigenvalues of (X^*X) and X^* is the conjugate transpose of X and $\mu_{max}(X)$ is the largest singular value of X.

Matrix X has the relationship between the norm values given below:

$$\frac{1}{\sqrt{n}}\|X\|_F \le \|X\|_2 \le \|X\|_F \quad (2)$$

Lemma 1.2 [13] For $A = [a_{ij}] \in \mathbb{M}_{m,n}(\mathbb{C})$, $B = [b_{ij}] \in \mathbb{M}_{m,n}(\mathbb{C})$, If C is the Hadamard product of A and B, then we get

$$\|C\|_2 \le m(A)n(B) \quad (3)$$

where $m(A) = max_{1\le i\le m}\sqrt{\sum_{j=1}^{n}|a_{ij}|^2}$ and $n(B) = max_{1\le j\le n}\sqrt{\sum_{j=1}^{m}|b_{ij}|^2}$.

Definition 1.3 [19] For $r \in C - \{0\}$, a matrix C_r is said to be r-circulant matrix if it is of the form and it is denoted by $C_r = Circ(r; \vec{c})$, where $\vec{c} = (c_0, c_1, \dots, c_{n-1})$ is the first row vector. For $r = 1$ and $r = -1$, we get the right and skew-right circulant matrices, respectively

$$C = \begin{bmatrix} c_0 & c_1 & c_2 & \dots & c_{n-2} & c_{n-1} \\ rc_{n-1} & c_0 & c_1 & \dots & c_{n-3} & c_{n-2} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ rc_2 & rc_3 & rc_4 & \dots & c_0 & c_1 \\ rc_1 & rc_2 & rc_3 & \dots & rc_{n1} & c_0 \end{bmatrix}$$

Lemma 1.4 [19] Let C_r be r-circulant matrices then its eigenvalues

$$\mu_i = \sum_{j=0}^{n-1} c_j\,(pw^{-i})^j, \quad i = 0,1,2,\dots,n-1$$

where w is the nth root of unity and p is the nth root of r.

Lemma 1.5 [13] The Euclidean norm of r-circulant matrix C_r is given by

$$\|C_r\|_E = \sqrt{\sum_{j=0}^{n-1}|C_j|^2[n - j(1 - |r|^2)]} \quad (4)$$

Lemma 1.6 [7] For any a and b ,we get

$$\prod_{i=0}^{n-1}(a - bp_i w_{-i}) = a^n - rb^n \quad (5)$$

where w is the n-th root of unity and p is the n-th root of r.

Lemma 1.7 [20] Determinant of the circulant matrix C_n is

$$|C_n| = \prod_{i=0}^{n-1}\left(\sum_{j=0}^{n-1} c_j\left(t_l^j\right)\right)$$

where the entry $\{i, j\}$ is equal to the entry $\{i + l, j + l\}$ for $l = 1,2, \dots$, $t_l = e^{\frac{2\pi l}{n}}$ are the n-th roots of unity and p_i are n^{th} root of of r. Using above lemmas, we can calculate the eigenvalues, the determinant, Euclidean norms, bounds for spectral norms of r-circulant matrices involving generalized Fermat numbers with arithmetic indices. We present many new identities for generalized Fermat numbers. In addition, r-circulant studies for Fermat and k-Fermat are handled in the most general format depending on the numbers a, b and n on these number sequences.

2. GENERALIZED FERMAT r-CIRCULANT MATRIX

Let u and v be non-negative integers and $r \in C - \{0\}$ and show the Generalized Fermat r-circulant matrices with $GR_{a,b,r}$.

Definition 2.1 Generalized Fermat r-circulant matrix is defined as
$GR_{a,b,r} = Circ(a, b, r; \vec{c})$ where first row vector is

$$\vec{c} = (GR_{a,b,u}, GR_{a,b,u+v}, GR_{a,b,u+2v}, \dots, GR_{a,b,u+(n-1)v})$$

ie., matrix of the form:

$$GR_{a,b,r} = \begin{bmatrix} GR_{a,b,u} & GR_{a,b,u+v} & \dots & GR_{a,b,u+(n-1)v} \\ rGR_{a,b,u+(n-1)v} & GR_{a,b,u} & \dots & GR_{a,b,u+(n-2)v} \\ \vdots & \vdots & \ddots & \vdots \\ rGR_{a,b,u+2v} & rGR_{a,b,u+3v} & \dots & GR_{a,b,u+v} \\ rGR_{a,b,u+v} & rGR_{a,b,u+2v} & \dots & GR_{a,b,u} \end{bmatrix}.$$

Let's show eigenvalues of $GR_{a,b,r}$ with $\lambda_i(GR_{a,b,r})$.

Theorem 2.2 The eigenvalues of the generalized Fermat r-circulant matrix $GR_{a,b,r}$ are as follows.

$$\lambda_i(GR_{a,b,r}) = \sum_{j=0}^{n-1} \left(a^{2^{u+jv}} + b^{2^{u+jv}}\right)\left(pw^{-i}\right)^j$$

Proof. If we substitute $a^{2^{u+jv}} + b^{2^{u+jv}}$ for c_j in Lemma 1.4, the proof ends.

Corollary 2.3

$$\sum_{j=0}^{n-1} \left(a^{2^{u+jv}} + b^{2^{u+jv}}\right)\left(pw^{-i}\right)^j = \sum_{j=0}^{n-1} \sqrt[n]{r^j(1)^{-ji}} \left(\left(a^{2^u}\right)^{(2^v)^j} + \left(b^{2^u}\right)^{(2^v)^j}\right)$$

Proof. Since w is the n-th root of unity and p is the n-th root of r, we get $(pw^{-i})^n = p^n(w^{-i})^n = r$.

$$\begin{aligned}
&\sum_{j=0}^{n-1} \left(a^{2^{u+jv}} + b^{2^{u+jv}}\right)\left(pw^{-i}\right)^j \\
&= \left(\left(a^{2^u}\right)^{(2^v)^0} + \left(b^{2^u}\right)^{(2^v)^0}\right) \\
&\quad + pw^{-i}\left(\left(a^{2^u}\right)^{(2^v)} + \left(b^{2^u}\right)^{(2^v)}\right) \\
&\quad + \left(pw^{-i}\right)^2\left(\left(a^{2^u}\right)^{(2^v)^2} + \left(b^{2^u}\right)^{(2^v)^2}\right) + \cdots \\
&\quad + \left(pw^{-i}\right)^{n-2}\left(\left(a^{2^u}\right)^{(2^v)^{n-2}} + \left(b^{2^u}\right)^{(2^v)^{n-2}}\right) \\
&\quad + \left(pw^{-i}\right)^{n-1}\left(\left(a^{2^u}\right)^{(2^v)^{n-1}} + \left(b^{2^u}\right)^{(2^v)^{n-1}}\right) \\
&= \sqrt[n]{r^0(1)^0}\left(\left(a^{2^u}\right)^{(2^v)^0} + \left(b^{2^u}\right)^{(2^v)^0}\right) \\
&\quad + \sqrt[n]{r(1)^{-i}}\left(\left(a^{2^u}\right)^{(2^v)} + \left(b^{2^u}\right)^{(2^v)}\right) \\
&\quad + \sqrt[n]{r^2(1)^{-2i}}\left(\left(a^{2^u}\right)^{(2^v)^2} + \left(b^{2^u}\right)^{(2^v)^2}\right) + \cdots \\
&\quad + \sqrt[n]{r^{n-2}(1)^{-(n-2)i}}\left(\left(a^{2^u}\right)^{(2^v)^{n-2}} + \left(b^{2^u}\right)^{(2^v)^{n-2}}\right) \\
&\quad + \sqrt[n]{r^{n-1}(1)^{-(n-1)i}}\left(\left(a^{2^u}\right)^{(2^v)^{n-1}} + \left(b^{2^u}\right)^{(2^v)^{n-1}}\right) \\
&= \sum_{j=0}^{n-1} \sqrt[n]{r^j(1)^{-ji}}\left(\left(a^{2^u}\right)^{(2^v)^j} + \left(b^{2^u}\right)^{(2^v)^j}\right).
\end{aligned}$$

If we simplify the operations for convenience, we obtain the following equation.

$$\begin{aligned}
&= \left(a^{2^u} + b^{2^u}\right) + pw^{-i}\left(\left(a^{2^u}\right)^{(2^v)} + \left(b^{2^u}\right)^{(2^v)}\right) \\
&+ \left(pw^{-i}\right)^2\left(\left(a^{2^u}\right)^{(2^v)^2} + \left(b^{2^u}\right)^{(2^v)^2}\right) + \cdots \\
&+ \frac{rw^{2i}}{p^2}\left(\left(a^{2^u}\right)^{(2^v)^{n-2}} + \left(b^{2^u}\right)^{(2^v)^{n-2}}\right) \\
&+ \frac{rw^{i}}{p}\left(\left(a^{2^u}\right)^{(2^v)^{n-1}} + \left(b^{2^u}\right)^{(2^v)^{n-1}}\right).
\end{aligned}$$

Corollary 2.4 For $r = 1$ and $r = -1$, we get eigenvalues for the generalized Fermat right circulant and skew-right circulant matrices respectively, so that

$$\begin{aligned}
&\lambda_i(GR_{a,b,1}) = \left(a^{2^u} + b^{2^u}\right) \\
&+ pw^{-i}\left(\left(a^{2^u}\right)^{(2^v)} + \left(b^{2^u}\right)^{(2^v)}\right) \\
&+ \left(pw^{-i}\right)^2\left(\left(a^{2^u}\right)^{(2^v)^2} + \left(b^{2^u}\right)^{(2^v)^2}\right) + \cdots \\
&+ \frac{w^{2i}}{p^2}\left(\left(a^{2^u}\right)^{(2^v)^{n-2}} + \left(b^{2^u}\right)^{(2^v)^{n-2}}\right)
\end{aligned}$$

$$+\frac{w^i}{p}\left(\left(a^{2^u}\right)^{(2^v)^{n-1}}+\left(b^{2^u}\right)^{(2^v)^{n-1}}\right)$$
$$\lambda_i\left(GR_{a,b,-1}\right)=\left(a^{2^u}+b^{2^u}\right)$$
$$+pw^{-i}\left(\left(a^{2^u}\right)^{(2^v)}+\left(b^{2^u}\right)^{(2^v)}\right)$$
$$+\left(pw^{-i}\right)^2\left(a^{2^u}\right)^{(2^v)^2}+((b^{2u})^{2v})^2+\cdots$$
$$-\frac{w^{2i}}{\xi^2}\left(\left(a^{2^u}\right)^{(2^v)^{n-2}}+\left(b^{2^u}\right)^{(2^v)^{n-2}}\right)$$
$$-\frac{w^i}{\xi}\left(\left(a^{2^u}\right)^{(2^v)^{n-1}}+\left(b^{2^u}\right)^{(2^v)^{n-1}}\right)$$

where ξ is the *n*-th root of -1.

Theorem 2.5 For a positive integer *n*, we have

$$\sum_{i=0}^{n-1}\left(a^{2^u}+b^{2^u}\right)$$
$$+pw^{-i}\left(\left(a^{2^u}\right)^{(2^v)}+\left(b^{2^u}\right)^{(2^v)}\right)$$
$$+\left(pw^{-i}\right)^2\left(\left(a^{2^u}\right)^{(2^v)}+\left(b^{2^u}\right)^{(2^v)}\right)+\cdots$$
$$+\frac{rw^{2i}}{p^2}\left(\left(a^{2^u}\right)^{(2^v)^{n-2}}+\left(b^{2^u}\right)^{(2^v)^{n-2}}\right)$$
$$+\frac{rw^i}{p}\left(\left(a^{2^u}\right)^{(2^v)^{n-1}}+\left(b^{2^u}\right)^{(2^v)^{n-1}}\right)$$
$$=n\left(a^{2^u}+b^{2^u}\right).$$

Proof. To get the desired result, we need to know that the trace of any given square matrix is equal to the sum of the eigenvalues of that matrix. In that case

$$\sum_{i=0}^{n-1}\lambda_i\left(GR_{a,b,r}\right)=$$
$$\sum_{i=0}^{n-1}\left(a^{2^u}+b^{2^u}\right)$$
$$+pw^{-i}\left(\left(a^{2^u}\right)^{(2^v)}+\left(b^{2^u}\right)^{(2^v)}\right)$$
$$+\left(pw^{-i}\right)^2\left(\left(a^{2^u}\right)^{(2^v)}+\left(b^{2^u}\right)^{(2^v)}\right)+\cdots$$
$$+\frac{rw^{2i}}{p^2}\left(\left(a^{2^u}\right)^{(2^v)^{n-2}}+\left(b^{2^u}\right)^{(2^v)^{n-2}}\right)$$
$$+\frac{rw^i}{p}\left(\left(a^{2^u}\right)^{(2^v)^{n-1}}+\left(b^{2^u}\right)^{(2^v)^{n-1}}\right)$$

according to the expression given above,

$$\sum_{i=0}^{n-1}\lambda_i\left(GR_{a,b,r}\right)=n.GR_{a,b,u}$$
$$=n\left(a^{2^u}+b^{2^u}\right).$$

The Sum Identity

On setting $i=0, r=1$ and $p=1$ in eigenvalues for generalized Fermat numbers, the following identity is verified for these numbers.

$$\sum_{j=0}^{n-1}GR_{a,b,u+jv}$$
$$=\left(a^{2^u}+b^{2^u}\right)+\left(\left(a^{2^u}\right)^{(2^v)}+\left(b^{2^u}\right)^{(2^v)}\right)$$
$$+\left(\left(a^{2^u}\right)^{(2^v)^2}+\left(b^{2^u}\right)^{(2^v)^2}\right)^2+\cdots$$
$$+\left(\left(a^{2^u}\right)^{(2^v)^{n-2}}+\left(b^{2^u}\right)^{(2^v)^{n-2}}\right)$$
$$+\left(\left(a^{2^u}\right)^{(2^v)^{n-1}}+\left(b^{2^u}\right)^{(2^v)^{n-1}}\right)$$

3. NORM OF GENERALIZED FERMAT *r*-CIRCULANT MATRICES

When we take $u=0$ and $v=1$, we get

$$GR^*_{a,b,r}=\begin{bmatrix} GR_{a,b,0} & GR_{a,b,1} & \dots & GR_{a,b,n-1} \\ rGR_{a,b,n-1} & GR_{a,b,0} & \dots & GR_{a,b,n-2} \\ \vdots & \vdots & \ddots & \vdots \\ rGR_{a,b,2} & rGR_{a,b,3} & \dots & GR_{a,b,1} \\ rGR_{a,b,1} & rGR_{a,b,2} & \dots & GR_{a,b,0} \end{bmatrix},$$

with the help of the lemma given below, the sum of the squares of the generalized Fermat numbers will be used to obtain the norms of different matrices.

Lemma 3.1 The finite sum of squares of the generalized Fermat numbers is given by

$$\sum_{j=0}^{n-1}\left(GR_{a,b,j}\right)^2=\sum_{i=0}^{n-1}(a^{2^i}+b^{2^i})^2$$

Proof.

$$\sum_{j=0}^{n-1}\left(GR_{a,b,j}\right)^2 = \sum_{j=0}^{n-1} a^{2^j} + 2(ab)^{2^j} + b^{2^j}$$
$$= a^{2^1} + a^{2^2} + \cdots + a^{2^n} + 2((ab)^1 + (ab)^2 + \cdots + (ab)^n) + b^{2^1} + b^{2^2} + \cdots + b^{2^n}$$
$$= \sum_{i=0}^{n-1}(a^{2^i} + b^{2^i})^2$$

Theorem 3.2 The Euclidean norm for the generalized Fermat r-circulant matrices is given by

$$\left\|GR^*_{a,b,r}\right\|_E = \sum_{j=0}^{n-1}\left(GR_{a,b,j+1} + 2(ab)^{2^j}\right)[n - j(1 - |r|^2)]$$

Proof. By equation (4) we get

$$\left\|GR^*_{a,b,r}\right\|_E^{\,2} = \sum_{j=0}^{n-1}\left(GR_{a,b,j+1} + 2(ab)^{2^j}\right)[n - j(1 - |r|^2)]$$

$$\sqrt{\sum_{i=0}^{n-1}(a^{2^i} + b^{2^i})^2} \le \left\|GR^*_{a,b,r}\right\|_2 \le \sqrt{a + b + |r|^2\left(\sum_{i=0}^{n-1}(a^{2^i} + b^{2^i})^2\right)}.\sqrt{\sum_{i=0}^{n-1}(a^{2^i} + b^{2^i})^2}$$

and for $|r| < 1$

$$|r|\sqrt{\textstyle\sum_{i=0}^{n-1}(a^{2^i} + b^{2^i})^2} \le \left\|GR^*_{a,b,r}\right\|_2 \le \sqrt{n\textstyle\sum_{i=0}^{n-1}(a^{2^i} + b^{2^i})^2}.$$

Proof. By Eq (4), the Euclidean norm is given as

$$\left\|GR^*_{a,b,r}\right\|_E^{\,2} = \sum_{j=0}^{n-1}\left(GR_{a,b,j}\right)^2[n - j(1 - |r|^2)]$$

State1.

If $|r| \ge 1$, then from Lemma 3.1 we get

Theorem 3.3 The bound for the spectral norm of the generalized Fermat r-circulant matrices is: for $|r| \ge 1$

$$\frac{\left\|GR^*_{a,b,r}\right\|_E}{\sqrt{n}} \ge \sqrt{\sum_{i=0}^{n-1}(a^{2^i} + b^{2^i})^2}$$

and from Eq. (1.2) we get

$$\left\|GR^*_{a,b,r}\right\|_2 \ge \sqrt{\sum_{i=0}^{n-1}(a^{2^i} + b^{2^i})^2}$$

Now to obtain the upper bound for the spectral norm, we write $GR^*_{a,b,r}$ in the form of the Hadamard product of two matrices.

$$A = \begin{bmatrix} GR_{a,b,0} & 1 & \dots & 1 & 1 \\ rGR_{a,b,n-1} & GR_{a,b,0} & \dots & 1 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ rGR_{a,b,2} & rGR_{a,b,3} & \dots & GR_{a,b,0} & 1 \\ rGR_{a,b,1} & rGR_{a,b,2} & \dots & rGR_{a,b,n-1} & GR_{a,b,0} \end{bmatrix}$$

and

$$B = \begin{bmatrix} 1 & GR_{a,b,1} & \dots & GR_{a,b,n-2} & GR_{a,b,n-1} \\ 1 & 1 & \dots & GR_{a,b,n-3} & GR_{a,b,n-2} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & 1 & \dots & 1 & GR_{a,b,1} \\ 1 & 1 & \dots & 1 & 1 \end{bmatrix}$$

then clearly $GR^*_{a,b,r} = A \circ B$, where $\circ$ denotes the Hadamard product. Now,

$$u(A) = \max_{1 \le i \le n} \sqrt{\sum_{j=1}^{n} |a_{ij}|^2}$$

$$= \sqrt{\left(GR_{a,b,0}\right)^2 + |r|^2 \sum_{j=1}^{n-1} \left(GR_{a,b,j}\right)^2}$$

$$= \sqrt{a + b + |r|^2 \sum_{i=0}^{n-1} (a^{2^i} + b^{2^i})^2}$$

$$\sqrt{\sum_{i=0}^{n-1} (a^{2^i} + b^{2^i})^2} \le \left\|GR^*_{a,b,r}\right\|_2$$

$$\le \sqrt{a + b + |r|^2 \sum_{i=0}^{n-1} (a^{2^i} + b^{2^i})^2} \cdot \sqrt{\sum_{i=0}^{n-1} (a^{2^i} + b^{2^i})^2}.$$

State 2. If $|r| < 1$, then from Eq. (4) and Lemma 3.1 we get

$$\left\|GR^*_{a,b,r}\right\|_E^2 \ge \sum_{j=0}^{n-1} (n-j)|r|^2 \left|GR_{a,b,j}\right|^2 + |r|^2 \sum_{j=0}^{n-1} j \left|GR_{a,b,j}\right|^2$$

$$= n|r|^2 \left(\sum_{i=0}^{n-1} (a^{2^i} + b^{2^i})^2 \right)$$

and

$$v(B) = \max_{1 \le j \le n} \sqrt{\sum_{j=1}^{n} |b_{ij}|^2}$$

$$= \sqrt{1 + \sum_{j=1}^{n-1} \left(GR_{a,b,j}\right)^2}$$

$$= \sqrt{1 + \sum_{i=0}^{n-1} (a^{2^i} + b^{2^i})^2}.$$

Thus, by Lemma 1.2, we get

$$\left\|GR^*_{a,b,r}\right\|_2 \le u(A)v(B) = \sqrt{a + b + |r|^2 \sum_{i=0}^{n-1} (a^{2^i} + b^{2^i})^2} \cdot \sqrt{\sum_{i=0}^{n-1} \left(a^{2^i} + b^{2^i}\right)^2}$$

$$\frac{\left\|GR^*_{a,b,r}\right\|_E}{\sqrt{n}} \ge |r| \sqrt{\sum_{i=0}^{n-1} (a^{2^i} + b^{2^i})^2}$$

and from Eq. (2) we get

$$\left\|GR^*_{a,b,r}\right\|_2 \ge |r| \sqrt{\sum_{i=0}^{n-1} (a^{2^i} + b^{2^i})^2}.$$

We calculate the upper bound for the spectral norm of $GR^*_{a,b,r}$.

Let

$$N = \begin{bmatrix} GR_{a,b,0} & GR_{a,b,1} & \dots & GR_{a,b,n-2} & GR_{a,b,n-1} \\ GR_{a,b,n-1} & GR_{a,b,0} & \dots & \dots & GR_{a,b,n-2} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ GR_{a,b,2} & GR_{a,b,3} & \dots & \dots & GR_{a,b,1} \\ GR_{a,b,1} & GR_{a,b,2} & \dots & \dots & GR_{a,b,0} \end{bmatrix}$$

hence, we have

$$M = \begin{bmatrix} 1 & 1 & 1 & \dots & 1 & 1 \\ r & 1 & 1 & \dots & 1 & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ r & r & r & \dots & r & 1 \\ r & r & r & \dots & r & 1 \end{bmatrix}$$

then clearly $GR^*_{a,b,r} = M \circ N$, where $\circ$ denotes the Hadamard product. So,

$$s_1(M) = \max_{1 \le i \le n} \sqrt{\sum_{j=1}^{n} |m_{ij}|^2} = \sqrt{n}$$

$$t_1(N) = \max_{1 \le j \le n} \sqrt{\sum_{j=1}^{n} |n_{ij}|^2}$$

$$= \sqrt{\sum_{j=1}^{n} GR_{a,b,j}{}^{2}} = \sqrt{\sum_{i=0}^{n-1} (a^{2^i} + b^{2^i})^2}$$

hence, by Lemma 1.2, we have

$$\left\| GR^{*}_{a,b,r} \right\|_2 \leq s_1(M) t_1(N)$$
$$= \sqrt{n \sum_{i=0}^{n-1} (a^{2^i} + b^{2^i})^2}$$

thus

$$|r| \sqrt{\sum_{i=0}^{n-1} (a^{2^i} + b^{2^i})^2} \leq \left\| GR^{*}_{a,b,r} \right\|_2$$
$$\leq \sqrt{n \left(\sum_{i=0}^{n-1} (a^{2^i} + b^{2^i})^2 \right)}$$

as desired.

4. CONCLUSION

Based on the generalized Fermat numbers with similar recurrences, which have been studied less, the r-circulant matrices of these sequences were created. Euclidean, row and spectral norms, which are eigenvalues, determinants, and some special norm values, are discussed depending on this matrix. In addition, the lower and upper bounds of these matrices for the spectral norm were examined in closed form. In addition, right circulant and skew-right circulant matrices are examined for 1 and -1 values of r depending on eigenvalues. Finally, some interesting results and sum properties are given.

Funding
The author (s) has no received any financial support for the research, authorship or publication of this study.

Authors' Contribution
The authors contributed equally to the study.

The Declaration of Conflict of Interest/ Common Interest
No conflict of interest or common interest has been declared by the authors.

The Declaration of Ethics Committee Approval
This study does not require ethics committee permission or any special permission.

The Declaration of Research and Publication Ethics
The authors of the paper declare that they comply with the scientific, ethical and quotation rules of SAUJS in all processes of the paper and that they do not make any falsification on the data collected. In addition, they declare that Sakarya University Journal of Science and its editorial board have no responsibility for any ethical violations that may be encountered, and that this study has not been evaluated in any academic publication environment other than Sakarya University Journal of Science.

REFERENCES

[1] W. Weisstein, Eric, "Proth Number", mathworld.wolfram.com, 2019.

[2] N. J. A. Sloane, “A Handbook of Integer Sequences”, Academic Press, New York, https://oeis.org, 1973.

[3] Generalized Fermat numbers, OEIS. Generalized Fermat numbers - OeisWiki, 2022.

[4] B. Fischer and J. Modersitzki, “Fast inversion of matrices arising in image processing”, Number Algorithms vol. 22 no.1, pp.1-11, 1999.

[5] S. Georgiou and C. Kravvaritis, New good quasi-cyclic codes over GF(3), International Journal of Algebra vol.1, no.1, pp.11-24, 2007.

[6] I. Kra and S. R. Simanca, On circulant matrices, Notices AMS vol.59, no.3, pp.368-377, 2012.

[7] A. C. F. Bueno, "On the Eigenvalues and the Determinant of the Right Circulant Matrices with Pell and Pell–Lucas Numbers", International Journal of Mathematics and Scientific Computing, vol.4, no.1, pp.19-20, 2014.

[8] S. Solak, "On the norms of circulant matrices with the Fibonacci and Lucas numbers", Applied Mathematics and Computation, vol.160, no.1, pp.125-132, 2005.

[9] S. Shen, J. Cen, "On the bounds for the norms of r-circulant matrices with the Fibonacci and Lucas numbers", Applied Mathematics and Computation, vol.216, no.10, pp. 2891-2897, 2010.

[10] S. Q. Shen, J. M. Cen, "On the spectral norms of r-circulant matrices with the k-Fibonacci and k-Lucas numbers", International Journal of Contemporary Mathematical Sciences,vol. 5, no. 12, pp. 569-578, 2010.

[11] Y. Zheng, S. Shon, "Exact inverse matrices of Fermat and Mersenne circulant matrix", In Abstract and Applied Analysis, Hindawi, 2015.

[12] D. Bozkurt, T. Y. Tam, "Determinants and inverses of circulant matrices with Jacobsthal and Jacobsthal–Lucas numbers", Applied Mathematics and Computation, vol.219, no. 2, pp.544-551, 2012.

[13] M. Kumaria, K. Prasada, J. Tantib, and E. Özkan, "On the properties of r-circulant matrices involving Mersenne and Fermat numbers", International Journal of Nonlinear Analysis and Applications, vol.1, no.11, 2023.

[14] M. Marin, "Generalized solutions in elasticity of micropolar bodies with voids", Revista de la Academia Canaria de Ciencias, vol.8, no.1, pp. 101-106, 1996

[15] M. Marin, "Contributions on uniqueness in thermoelastodynamics on bodies with voids", Ciencias matemáticas (Havana), vol.16, no.2, pp.101-109, 1998.

[16] Z. Pucanovic, and M. Pesovic "Chebyshev polynomials and r-circulant matrices", Applied Mathematics and Computation, vol.437, no.127521, 2023.

[17] J. M. Blackledget, "Vector and Matrix norm", Digital Signal Processing, pp.208-236, 2006.

[18] Simon Foucart, http://www.math.drexel.edu/foucart/TeachingFiles/F12/M504Lect6.pdf

[19] R. E. Cline, R. J. Plemmons, and G. Worm, "Generalized inverses of certain Toeplitz matrices", Linear Algebra and its Application vol.8, no.1, pp.25-33, 1974.

[20] K. Irwin, R. Santiago, Simanca, "On circulant matrices", ://www.math.colombia.edu/ums/pdf/cir-not5.pdf