
Univers. J. Math. Appl., 8 (1) (2025 1-7 https://doi.org/10.32323/ujma.1566270

UJMA

Universal Journal of Mathematics and Applications

ISSN (Online) 2619-9653
https://dergipark.org.tr/en/pub/ujma



Research Article

A Study on the *k*-Mersenne and *k*-Mersenne-Lucas Sequences

Engin Özkan[1*], **Bayram Şen**[2], **Hakan Akkuş**[3] **and Mine Uysal**[4]

[1]*Department of Mathematics, Faculty of Sciences, Marmara University, İstanbul, Türkiye*
[2,3]*Department of Mathematics, Graduate School of Natural and Applied Sciences, Erzincan Binali Yıldırım University, Erzincan, Türkiye*
[4]*Department of Mathematics, Faculty of Sciences and Arts, Erzincan Binali Yıldırım University, Erzincan, Türkiye*
[*]*Corresponding author*

Article Info

Keywords: *Catalan numbers, Fibonacci numbers, Generating functions, Hankel transform, k-Mersenne sequences, k-Mersenne-Lucas sequences*
2010 AMS: *11B39, 11B83, 11C20*
Received: *13 October 2024*
Accepted: *06 February 2025*
Available online: *23 February 2025*

Abstract

In this study, we examine an application of $k-$Mersenne and $k-$Mersenne-Lucas sequences. We present the Catalan transforms of these sequences and give the properties of these Catalan transforms. Catalan transforms of $k-$Mersenne and $k-$Mersenne-Lucas sequences have terms according to different values of k, and some of them are associated with the sequences in OEIS. We obtain the generating functions of the Catalan transforms of $k-$Mersenne and $k-$Mersenne-Lucas sequences. Moreover, we apply the Hankel transform to the Catalan transforms of these sequences. Finally, the determinants of the created matrices are calculated by applying the Hankel transform to the terms of the Catalan transforms of these sequences.

1. Introduction

Number sequences constitute an important subject area in mathematics. These sequences usually consist of consecutive numbers that follow a certain rule. The Fibonacci sequence is one of the most well-known and frequently encountered number sequences. These sequences have intrigued scientists for a long time. Fibonacci and Lucas numbers and many of their applications have been studied and many properties of these sequences have been investigated [1]. New sequences are obtained by changing the recurrence relation and initial conditions of the generalized Fibonacci sequence.
In [2], Falcon defined, $k-$Fibonacci sequence. This paper is an extension of the work of Falcon [3]. In [3], Falcon gave an application of the Catalan transform to the k-Fibonacci sequences.
Mersenne, Mersenne-Lucas numbers are a generalization of these numbers, similar to the Fibonacci sequence. Many researchers have studied Mersenne number sequences and their different applications have been investigated [4–17].

Definition 1.1. *The Mersenne numbers M_n are defined by the following recurrence relation for $n \in \mathbb{N}$,*

$$M_{n+2} = 3M_{n+1} - 2M_n$$

with the initial conditions $M_0 = 0$, $M_1 = 1$ [16].

Definition 1.2. *The Mersenne-Lucas numbers m_n are defined by the following recurrence relation for $n \in \mathbb{N}$*

$$m_{n+2} = 3m_{n+1} - 2m_n,$$

with the initial conditions $m_0 = 2$, $m_1 = 3$, [17].

Definition 1.3. *[16] The Binet formula for Mersenne numbers is defined by*

$$M_n = 2^n - 1.$$

Definition 1.4. *[17] The Binet formula for Mersenne-Lucas numbers is defined by*

$$m_n = 2^n + 1.$$

***Email addresses and ORCID numbers*: engin.ozkan@marmara.edu.tr, 0000-0002-4188-7248 (E.Özkan), bayram.s29@hotmail.com, 0009-0009-5570-4981 (B.Şen), hakan.akkus@ogr.ebyu.edu.tr, 0000-0001-9716-9424 (H. Akkuş), mine.uysal@erzincan.edu.tr, 0000-0002-2362-3097 (M. Uysal)**
Cite as: E.Özkan, B.Şen, H.Akkuş, M.Uysal *A Study on the k-Mersenne and k-Mersenne-Lucas Sequences*, Univers. J. Math. Appl., 8(1) (2025), 1-7.



Catalan and Hankel transforms of the some number sequences have been investigated by some authors and their properties have been examined [18–24].

In this paper, the Catalan transforms of these sequences are examined in detail. We give, the generating functions for the Catalan transformations of the sequences. Finally, the Hankel transforms of the newly obtained sequences and the determinants of these transforms are discussed.

Definition 1.5. *Let k be any positive number, then the k−Mersenne sequence is recursively given for $n \in \mathbb{N}$ as*

$$M_{k,n+2} = 3kM_{k,n+1} - 2M_{k,n},$$

where $M_{k,0} = 0$, $M_{k,1} = 1$. From now on we will show the sequence as $M_{k,n}$. When $k = 1$, Mersenne sequence is obtained [16]. The characteristic equation of the sequence is as follows

$$x^2 - 3kx + 2 = 0.$$

Its characteristic roots are

$$x_1 = \frac{3k + \sqrt{9k^2 - 8}}{2} \text{ and}$$

$$x_2 = \frac{3k - \sqrt{9k^2 - 8}}{2}.$$

Characteristic roots prove the following properties to be true.

$$x_1 + x_2 = 3k,$$
$$x_1 x_2 = 2,$$
$$x_1 - x_2 = \sqrt{9k^2 - 8}.$$

Binet's formula for $M_{k,n}$ is

$$M_{k,n} = \frac{{x_1}^n - {x_2}^n}{x_1 - x_2}.$$

Definition 1.6. *Let k be any positive number, the k−Mersenne-Lucas number sequence is recursively given for $n \in \mathbb{N}$ as*

$$m_{k,n+2} = 3km_{k,n+1} - 2m_{k,n},$$

where $m_{k,0} = 2, m_{k,1} = 3k$ [9].From now on we will show the sequence as $m_{k,n}$. When $k = 1$, Mersenne-Lucas sequence is obtained. The characteristic equation of the sequence is as follows

$$x^2 - 3kx + 2 = 0.$$

Its characteristic roots are

$$x_1 = \frac{3k + \sqrt{9k^2 - 8}}{2}$$

and

$$x_2 = \frac{3k - \sqrt{9k^2 - 8}}{2}.$$

Binet's formula for $m_{k,n}$ is

$$m_{k,n} = {x_1}^n + {x_2}^n.$$

2. Catalan Transforms for the $k-$Mersenne and $k-$Mersenne Lucas Sequences

In this section, we present the Catalan transform of these sequences and give the properties of these Catalan transforms. Catalan transforms of k-Mersenne and k-Mersenne-Lucas sequences have terms according to different values of k, and some of them are associated with the sequences in OEIS.

Definition 2.1. *The n^{th} Catalan number is introduced by Barry [8]. For $n \geq 0$, it is*

$$C_n = \frac{1}{n+1}\binom{2n}{n}$$

or

$$C_n = \frac{2n!}{(n+1)!n!}.$$

The Catalan number generating function is

$$C(x) = \frac{1 - \sqrt{1 - 4x}}{2x}.$$

Some Catalan numbers are in the order $1, 1, 2, 5, 14, 42, 132, 429, 1430, 4862, \ldots$.

Using the definition of Catalan numbers, the Catalan transform of these sequences can be defined as follows.

Definition 2.2. *Let $k>0, n>0$ and $CM_{k,0}=0$;*

$$CM_{k,n}=\sum_{i=0}^{n}\frac{i}{2n-i}\binom{2n-i}{n-i}M_{k,i}$$

equation is called Catalan transforms in $k-$Mersenne sequences. Based on this definition, the terms of the Catalan transforms of the $k-$Mersenne sequence are as follows.

$$CM_{k,1}=\sum_{i=0}^{1}\frac{i}{2-i}\binom{2-i}{1-i}M_{k,i}=\binom{1}{0}M_{k,1}=1,$$
$$CM_{k,2}=\sum_{i=0}^{2}\frac{i}{4-i}\binom{4-i}{2-i}M_{k,i}=\frac{1}{3}\binom{3}{1}M_{k,1}+\binom{2}{0}M_{k,2}=3k+1,$$
$$CM_{k,3}=\sum_{i=0}^{3}\frac{i}{6-i}\binom{6-i}{3-i}M_{k,i}=9k^2+6k,$$
$$CM_{k,4}=\sum_{i=0}^{4}\frac{i}{8-i}\binom{8-i}{4-i}M_{k,i}=27k^3+27k^2+3k-1,$$
$$CM_{k,5}=\sum_{i=0}^{5}\frac{i}{10-i}\binom{10-i}{5-i}M_{k,i}=81k^4+108k^3+27k^2-6k,$$
$$CM_{k,6}=\sum_{i=0}^{6}\frac{i}{12-i}\binom{12-i}{6-i}M_{k,i}=243k^5+405k^4+162k^3-18k^2-6k+6,$$
$$CM_{k,7}=\sum_{i=0}^{7}\frac{i}{14-i}\binom{14-i}{7-i}M_{k,i}=729k^6+1458k^5+2025k^4-864k^2+36k+84.$$

We can write the found $CM_{k,n}$ sequences as the product of the $nx1$-dimensional $M_{k,n}$ matrix and the lower triangular matrix C.

$$\begin{bmatrix} CM_{k,1}\\ CM_{k,2}\\ CM_{k,3}\\ CM_{k,4}\\ CM_{k,5}\\ CM_{k,6}\\ \vdots \end{bmatrix}=\begin{bmatrix} 1 & & & & & & \\ 1 & 1 & & & & & \\ 2 & 2 & 1 & & & & \\ 5 & 5 & 3 & 1 & & & \\ 14 & 14 & 9 & 4 & 1 & & \\ 42 & 42 & 28 & 14 & 5 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \end{bmatrix}\begin{bmatrix} M_{k,1}\\ M_{k,2}\\ M_{k,3}\\ M_{k,4}\\ M_{k,5}\\ M_{k,6}\\ \vdots \end{bmatrix}$$

In this case, the 1^{st} and 2^{nd} columns of the C matrix are filled with Catalan numbers, and the elements of this matrix validate a recurrence relation. The Table 2.1 shows the coefficients of the terms of the Catalan $k-$Mersenne sequence, which we call the Catalan triangle.

$CM_{k,1}$	1					
$CM_{k,2}$	3	1				
$CM_{k,3}$	9	6				
$CM_{k,4}$	27	27	3	-1		
$CM_{k,5}$	81	108	27	-6		
$CM_{k,6}$	243	405	162	-18	-6	6
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$

Table 2.1: Catalan triangle of $k-$Mersenne sequence

The results of some Catalan transforms are presented below.

$CM_1=\{0,\ 1,\ 4,\ 15,\ 56,\ 210,\ 792,\ \dots\}$ It is indexed as A001791 in OEIS.

$CM_2=\{0,1,\ 7,\ 48,\ 329,\ 2256,\ 15474,\ \dots\}$

$CM_3=\{0,\ 1,\ 10,\ 99,\ 980,\ 9702,\ 96054,\ \dots\}$

$CM_4=\{0,\ 1,\ 13,\ 168,\ 2171,\ 28056,\ 362524,\ \dots\}$

$CM_5=\{0,\ 1,\ 16,\ 255,\ 4064,\ 64770,\ 1032276,\ \dots\}$

$CM_6=\{0,\ 1,\ 19,\ 360,\ 6821,\ 129240,\ 2464882,\ \dots\}$

Definition 2.3. *Let $k>0, n>0$ and $Cm_{k,0}=0$;*

$$Cm_{k,n} = \sum_{i=0}^{n} \frac{i}{2n-i} \binom{2n-i}{n-i} m_{k,i}$$

equation is called Catalan transform in k-Mersenne-Lucas sequences. Based on this definition, the Catalan transform of the k-Mersenne-Lucas sequence are found below.

$$Cm_{k,1} = \sum_{i=0}^{1} \frac{i}{2-i} \binom{2-i}{1-i} m_{k,i} = \binom{1}{0} m_{k,1} = 3k,$$
$$Cm_{k,2} = \sum_{i=0}^{2} \frac{i}{4-i} \binom{4-i}{2-i} m_{k,i} = \frac{1}{3}\binom{3}{1} m_{k,1} + \binom{2}{0} m_{k,2} = 9k^2+3k-4,$$
$$Cm_{k,3} = \sum_{i=0}^{3} \frac{i}{6-i} \binom{6-i}{3-i} m_{k,i} = 27k^3+18k^2-12k-8,$$
$$Cm_{k,4} = \sum_{i=0}^{4} \frac{i}{8-i} \binom{8-i}{4-i} m_{k,i} = 81k^4+81k^3-27k^2-39k-12,$$
$$Cm_{k,5} = \sum_{i=0}^{5} \frac{i}{10-i} \binom{10-i}{5-i} m_{k,i} = 243k^5+324k^4-27k^3-162k^2-60k-24,$$
$$Cm_{k,6} = \sum_{i=0}^{6} \frac{i}{12-i} \binom{12-i}{6-i} m_{k,i} = 729k^6+1215k^5+162k^4-594k^3-306k^2-78k-72,$$
$$Cm_{k,7} = \sum_{i=0}^{7} \frac{i}{14-i} \binom{14-i}{7-i} m_{k,i} = 2187k^7+4374k^6+1458k^5-1944k^4-1458k^3-324k^2-192k-240.$$

We can write the found $Cm_{k,n}$ sequences as the product of the $nx1$-dimensional $m_{k,n}$ matrix and the lower triangular matrix C.

$$\begin{bmatrix} Cm_{k,1} \\ Cm_{k,2} \\ Cm_{k,3} \\ Cm_{k,4} \\ Cm_{k,5} \\ Cm_{k,6} \\ \vdots \end{bmatrix} = \begin{bmatrix} 1 & & & & & & \\ 1 & 1 & & & & & \\ 2 & 2 & 1 & & & & \\ 5 & 5 & 3 & 1 & & & \\ 14 & 14 & 9 & 4 & 1 & & \\ 42 & 42 & 28 & 14 & 5 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \end{bmatrix} \begin{bmatrix} m_{k,1} \\ m_{k,2} \\ m_{k,3} \\ m_{k,4} \\ m_{k,5} \\ m_{k,6} \\ \vdots \end{bmatrix}$$

In this case, the 1^{st} and 2^{nd} columns of the C matrix are filled with Catalan numbers, and the elements of this matrix validate a recurrence relation. The Table 2.2 shows the coefficients of the terms of the Catalan transform of the k-Mersenne sequence, which we call the Catalan triangle.

$Cm_{k,1}$	3						
$Cm_{k,2}$	9	3	-4				
$Cm_{k,3}$	27	18	-12	-8			
$Cm_{k,4}$	81	81	-27	-39	-12		
$Cm_{k,5}$	243	324	-27	-162	-60	-24	
$Cm_{k,6}$	729	1215	162	-594	-306	-78	-72
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$

Table 2.2: Catalan triangle of $k-$Mersenne-Lucas sequence

3. The Generating Functions of the Catalan Transform of the $k-$ Mersenne and $k-$Mersenne-Lucas Sequences

In this section, we obtain the generating functions of the Catalan transforms of k-Mersenne and k-Mersenne-Lucas sequences.

Theorem 3.1. *The generating function of the k-Mersenne sequence is*

$$M(x) = \frac{x}{2x^2-3kx+1}$$

[9].

Theorem 3.2. *The generating function of the Catalan transform of the k-Mersenne sequences is*

$$T(x)=\frac{1-\sqrt{1-4x}}{4-4x-3k+(3k-2)\sqrt{1-4x}}.$$

Proof. The generating function of k-Mersenne and Catalan sequences are as follows, respectively,

$$M(x)=\frac{x}{2x^2-3kx+1}$$

and

$$C(x)=\frac{1-\sqrt{1-4x}}{2x}$$

If this information and the properties of the composition process in the functions are used, the desired result is achieved. Thus, the following equations for the generating function of the Catalan transform of the k-Mersenne sequence are obtained.

$$\begin{aligned}T(x)=M(x)*C=\ T(x*C(x))&=\frac{\frac{1-\sqrt{1-4x}}{2}}{2\left(\frac{1-\sqrt{1-4x}}{2}\right)^2-\frac{3k\left(1-\sqrt{1-4x}\right)}{2}+1}\\&=\frac{\frac{1-\sqrt{1-4x}}{2}}{\frac{1-2\sqrt{1-4x}+1-4x}{2}-\frac{3k-3k\sqrt{1-4x}}{2}+1}\\&=\frac{\frac{1-\sqrt{1-4x}}{2}}{\frac{1-2\sqrt{1-4x}+1-4x-3k+3k\sqrt{1-4x}+2}{2}}\\&=\frac{1-\sqrt{1-4x}}{4-4x-3k+(3k-2)\sqrt{1-4x}}.\end{aligned}$$

□

Theorem 3.3. *The generating function of the k−Mersenne-Lucas sequence is*

$$m(x)=\frac{2-3kx}{2x^2-3kx+1}$$

Theorem 3.4. *The generating function of the Catalan transform of the k-Mersenne-Lucas number is*

$$L(x)=\frac{4-3k+3k\sqrt{1-4x}}{4-4x-3k+(3k-2)\sqrt{1-4x}}.$$

Proof. The generating function of k-Mersenne-Lucas and Catalan sequences are as follows respectively,

$$m(x)=\frac{2-3kx}{2x^2-3kx+1}$$

and

$$C(x)=\frac{1-\sqrt{1-4x}}{2x}.$$

Thus, the following equations for the generating function of the Catalan transform of the k-Mersenne-Lucas sequence is obtained.

$$\begin{aligned}L(x)=m(x)*C=m(x*C(x))&=\frac{2-3k\left(\frac{1-\sqrt{1-4x}}{2}\right)}{2\left(\frac{1-\sqrt{1-4x}}{2}\right)^2-\frac{3k\left(1-\sqrt{1-4x}\right)}{2}+1}\\&=\frac{2-3k\left(\frac{1-\sqrt{1-4x}}{2}\right)}{\frac{1-2\sqrt{1-4x}+1-4x}{2}-\frac{3k-3k\sqrt{1-4x}}{2}+1}\\&=\frac{\frac{4-3k+3k\sqrt{1-4x}}{2}}{\frac{1-2\sqrt{1-4x}+1-4x-3k+3k\sqrt{1-4x}+2}{2}}\\&=\frac{4-3k+3k\sqrt{1-4x}}{4-4x-3k+(3k-2)\sqrt{1-4x}}.\end{aligned}$$

□

4. Hankel Transform

In this section, we apply the Hankel transform to the Catalan transforms of these sequences. In addition, the determinants of the created matrices are calculated by applying the Hankel transform to the terms of the Catalan transforms of these sequences.

Definition 4.1. *Let $R = (r_1, r_2, r_3, r_4, \dots)$ be a sequence of real numbers. The Hankel transform of R is the sequence of determinants $H_n = \det\left[r_{i+j-2}\right]$. That is,*

$$H_n = \begin{vmatrix} r_0 & r_1 & r_2 & r_3 & \dots \\ r_1 & r_2 & r_3 & r_4 & \dots \\ r_2 & r_3 & r_4 & r_5 & \dots \\ r_3 & r_4 & r_5 & r_6 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{vmatrix}$$

the n^{th} order Hankel determinant of R is the upper left nxn subdeterminant of H.

Based on this definition, we apply the Hankel transform to the Catalan transforms of $k-$Mersenne and $k-$ Mersenne-Lucas sequences, respectively, as follows.

$$HCM_{k,1} = |1| = 1,$$

$$HCM_{k,2} = \begin{vmatrix} 1 & 3k+1 \\ 3k+1 & 9k^2+6k \end{vmatrix} = -1,$$

$$HCM_{k,3} = -1,$$

and

$$HCm_{k,1} = det\,|3k| = 3k,$$

$$HCm_{k,2} = \begin{vmatrix} 3k & 9k^2+3k-4 \\ 9k^2+3k-4 & 27k^3+18k^2-12k-8 \end{vmatrix} = 27k^2-16,$$

$$HCm_{k,3} = 270k^4+297k^3-304k^2-240k\ -128.$$

5. Conclusion

In this study, we presented the Catalan transform of $k-$Mersenne and $k-$Mersenne-Lucas sequences and obtained the generating functions of the Catalan transform of these sequences. Catalan transform of $k-$Mersenne and $k-$ Mersenne-Lucas sequences have terms according to different values of k, and some of them associated with the sequences in OEIS. Moreover, we applied the Hankel transform to the Catalan transforms of $k-$Mersenne and $k-$Mersenne-Lucas sequences. Finally, the determinants of the created matrices calculated by applying the Hankel transform to the terms of the Catalan transform of the $k-$Mersenne and $k-$Mersenne-Lucas sequences. If this study is examined, the same results can be found for other sequences. For example, the Catalan and Hankel transforms of Perrin and Padovan sequences.

Article Information

Acknowledgements: The authors would like to express their sincere thanks to the editor and the anonymous reviewers for their helpful comments and suggestions.

Author's Contributions: All authors contributed equally to the writing of this paper.

Conflict of Interest Disclosure: No potential conflict of interest was declared by the authors.

Plagiarism Statement: This article was scanned by the plagiarism program.

References

[1] T. Koshy, *Fibonacci and Lucas Numbers with Applications*, John Wiley & Sons, 2019.
[2] S. Falcon, A. Plaza, *On the Fibonacci k-Numbers*, Chaos Solutions Fractals, **32**(5) (2007), 1615-1624.
[3] S. Falcon, *Catalan transform of the k-Fibonacci sequence*, Communications of the Korean Mathematical Society, **28**(4) (2013), 827–832.
[4] Y. Soykan, *On generalized p-Mersenne numbers*, Earthline Journal of Mathematical Sciences, **8**(1) (2022), 83-120.
[5] R. Frontczak, T. P. Goy, *Mersenne-Horadam identities using generating functions*, Carpathian Math. Publ., **12**(1) (2020), 34-45.
[6] M. Uysal, M. Kumari, B. Kuloğlu, K. Prasad, E. Özkan, *On the hyperbolic k-Mersenne and k-Mersenne-Lucas octonions*, Kragujevac J. Math.,**49**(5) (2025), 765-779.
[7] D. Tasci, *On Gaussian Mersenne numbers*, J. Sci. Arts, **21**(4) (2021), 1021-1028.
[8] P. Barry, *A Catalan transform and related transformations on integer sequences*, J. Integer Seq. J., **8**(4) (2005), 1-24.
[9] M. Chelgham, A. Boussayoud, *On the k-Mersenne-Lucas numbers*, Notes Number Theory Discrete Math., **27**(1) (2021), 7-13.
[10] M. Kumari, K, Prasad, J, Tanti, *On the generalization of Mersenne and Gaussian Mersenne polynomials*, J. Anal., **32**(3) (2024), 931-947.
[11] Y. Soykan, *A study on generalized Mersenne numbers*, Journal of Progressive Research in Mathematics, **18**(3) (2021), 90-112.
[12] S. Demiriz, S. Erdem, *Mersenne matrix operator and its application in p-summable sequence space*, Commun. Adv. Math. Sci., **7**(1)(2024), 42-55.

[13] E. Özkan, M. Uysal, *Mersenne-Lucas hybrid numbers*, Math. Montisnigri, **52**(2021), 17-29.
[14] M. Kumari, K. Prasad, R. Mohanta, *Algebra of quaternions and octonions involving higher order Mersenne numbers*, Proc. Indian Nat. Sci. Acad., (2024), 1-10.
[15] M. Kumari, J. Tanti, K. Prasad, *On some new families of k-Mersenne and generalized k-Gaussian Mersenne numbers and their polynomials*, (2021), arXiv preprint arXiv:2111.09592.
[16] P. Catarino, H. Campos, H., P. Vasco, *On the Mersenne sequence*, Ann. Math. Inform., **46** (2016), 37-53.
[17] N. Saba, A. Boussayoud, K. V. V. Kanuri, *Mersenne Lucas numbers and complete homogeneous symmetric functions*, J. Math. Comput. Sci., **24**(2) (2021), 127- 139.
[18] A. Patra, M. K. Kaabar, *Catalan transform of k-Balancing sequences*, Int. J. Math. Math. Sci., **2021**(1) 2021, 9987314.
[19] S. Kapoor, P. Kumar, *Catalan Transformation of (s; t) Padovan Sequences*, Asian Journal of Pure and Applied Mathematics, **5**(1) (2023), 170-178.
[20] P. J. Larcombe, P. D. Wilson, *On the generating function of the Catalan sequence: A historical perspective*, Congr. Numer., **149** (2001), 97-108.
[21] J. W. Layman, *The Hankel transform and some of its properties*, J. Integer Seq., **4**(1) 2001, 1-11.
[22] P. M. Rajković, M. D. Petković, P. Barry, *The Hankel transform of the sum of consecutive generalized Catalan numbers*, Integral Transforms Spec. Funct., **18**(4) (2007), 285-296.
[23] K. Parmar, V. R. Gorty, *Quaternion Hankel Transform and its Generalization*, Sahand Commun. Math. Anal., **21**(1) (2024), 67-81.
[24] E. Özkan, M. Uysal, B. Kuloğlu, *Catalan transform of the incomplete Jacobsthal numbers and incomplete generalized Jacobsthal polynomials*, Asian-Eur. J. Math., **15**(6) (2022), 2250119.